

Thesis



**Physiological Significance of Ethrel (2-Chloroethyl  
Phosphonic Acid) and Nitrogen in Relation to  
Growth and Metabolism of Mustard under  
Irrigated and Non-Irrigated Conditions**

**ABSTRACT**

**THESIS**

SUBMITTED FOR THE DEGREE OF

**Doctor of Philosophy**

IN

**BOTANY**

**MOHMAD RAMZAN MIR**

DEPARTMENT OF BOTANY  
ALIGARH MUSLIM UNIVERSITY  
ALIGARH (INDIA)

**2002**

3

**Physiological Significance of Ethrel (2-Chloroethyl Phosphonic Acid)  
and Nitrogen in Relation to Growth and Metabolism of Mustard  
under Irrigated and Non-Irrigated Conditions**

**Mohmad Ramzan Mir**

Abstract of the thesis submitted to the Aligarh Muslim University, Aligarh, India for the degree of **Doctor of Philosophy in Botany**.

The present thesis comprises six chapters. Chapter 1 deals with the importance of the problem. Lacunae in the understanding of the problem and justifications for undertaking the present study have been put forth.

Chapter 2 is review of literature. Relevant available literature pertaining to individual as well as combined effect of plant growth regulators especially ethylene sources with nitrogen on crop growth and development has been given in this chapter.

Chapter 3 describes the details of the material used in the study and methods employed in determining observations carried out in the experiments. Relevant information on meteorological and edaphic data has been included.

In Chapter 4, the results obtained in the experiments which were found significant at  $P < 0.05$  have been recorded in detail.

In Chapter 5, significant results have been discussed in the light of earlier reported findings. Possible explanations of the data obtained have also

been given to reach a conclusion. The results of the five field experiments are summarized below.

Experiment 1 (1998-99) was conducted under irrigated conditions to study the response of two cultivars (Alankar and PBM16) of mustard (*Brassica juncea* L.) to leaf-applied ethrel at 60d after sowing (flowering stage). Alankar is a well-established cultivar and accepted by the mustard growers for a decade, whereas PBM16 is a newly released. This was a factorial experiment conducted according to randomized complete block design. The response of the cultivars to ethrel treatments was assessed by determining growth and biochemical characteristics at 80 (pod fill), 100 (pod maturity) and 120d (harvest) after sowing. Physiological characteristics were studied at 80 and 100d after sowing. Yield and quality characteristics were determined at harvest. Growth characteristics were plant height, plant leaf area, leaf area index, specific leaf area, specific leaf weight, plant dry weight and dry weights and per cent dry weight distribution in leaf, stem and pod, leaf fresh weight, leaf turgid weight and leaf relative water content. Physiological characteristics were, rate of photosynthesis, stomatal conductance, internal CO<sub>2</sub> concentration, transpiration rate, carboxylation efficiency, photosynthetic water use efficiency, and plant water use efficiency. Biochemical characteristics included N content and N accumulation in plant. At harvest, yield characteristics studied were, number of pods per plant, number of seeds per pod, 1000 seed weight, seed yield,

biological yield, harvest index and oil yield. The oil was assessed for acid, iodine and saponification values.

Among five concentrations of ethrel (0, 100, 200, 400 and 600 $\mu$ L/L) applied, 200 $\mu$ L/L was found superior over others in increasing the plant characteristics studied. The concentration less than 200 $\mu$ L/L was found less effective, whereas concentration higher than 200 $\mu$ L/L proved inhibitory. Spray of 200 $\mu$ L/L ethrel affected growth of the plants and increased total leaf area per plant. Thus, the increase in photosynthesising surface area led to increase in photosynthesis and CO<sub>2</sub> accumulation, which resulted in increased plant dry weight. The other effects of ethrel spray was noted in improved source-sink relationship, as seen in increase in per cent pod dry mass in ethrel-treated plants. More of the flowers developed into pods, evident from higher pod number and seed yield in ethrel-treated plants compared with water-sprayed control. The oil quality was also improved with 200 $\mu$ L/L ethrel spray.

Comparison of the cultivars showed that Alankar established its superiority over PBM16. Growth, physiological, biochemical, yield and quality characteristics were found superior in Alankar than in PBM16. Interaction effect between cultivar and ethrel spray was non-significant for most of the plant characteristics. This suggests that the two cultivars responded similarly to ethrel spray.

Experiment 2 (1998-99) was a factorial randomized complete block design conducted on the same lines as Experiment 1 but under non-irrigated



conditions. The scheme of the treatments, design of the experiment, ethrel spray treatments and cultivars of mustard were also similar as described for Experiment 1. The observations recorded at different sampling stages were similar to Experiment 1. In this experiment, it was noted that ethrel spray at 200 $\mu$ L/L concentration was more effective than any other concentrations used. The effect of the spray was manifested through changes in various characteristics as described for Experiment 1. Alankar cultivar surpassed PBM16 in performance.

Combined analysis of the two experiments showed that the factors irrigated and non-irrigated were non-significant. Ethrel spray under irrigated and non-irrigated conditions was equally effective. The two cultivars also behaved similarly in the two conditions of irrigation.

Experiment 3 (1999-2000) was a factorial randomized complete block design conducted under irrigated conditions to study the effect of leaf-applied 0, 100 and 200 $\mu$ L/L ethrel on the performance of Alankar cultivar of mustard (the cultivar was selected on the basis of Experiment 1) grown with soil-applied 0, 40, 60 and 80kg N/ha. Ethrel spray application was done at 60d after sowing (flowering stage) and performance of the crop was assessed by determining various plant characteristics at 80 (pod fill), 100 (pod maturity) and 120d (harvest) after sowing. Growth characteristics were those studied in Experiment 1. Among physiological characteristics, 1-aminocyclopropane-1-carboxylic acid content, ACC oxidase and ethylene evolution were also studied

in addition to the characteristics studied in Experiment 1. Among biochemical characteristics, leaf nitrate reductase activity was also studied in addition to the determination of N content and N accumulation in plant. Among yield characteristics, seed N, nitrogen harvest index and nitrogen yield potential were also studied in addition to the yield characteristics studied in Experiment 1. Quality characteristics were similar as in Experiment 1.

Ethrel at 200 $\mu$ L/L concentration and nitrogen at 80kg N/ha registered significantly superior values as compared to other treatments. Ethrel (200 $\mu$ L/L) enhanced growth, physiological, biochemical, yield and quality characteristics. In this experiment, it was found that the effect of 200 $\mu$ L/L ethrel was maximal when plants received soil-applied 80kg N/ha. This combination (200 $\mu$ L/L ethrel and 80kg N/ha) enhanced plant leaf area, photosynthesis, CO<sub>2</sub> accumulation and plant dry weight. Water relations characteristics such as leaf fresh weight, leaf turgid weight and leaf relative water content were also enhanced by 200 $\mu$ L/L ethrel x 80kg N/ha. Pod dry weight was maximal with 200 $\mu$ L/L ethrel x 80kg N/ha, which showed higher translocation of dry matter towards sink (pods). Among physiological characteristics, rate of photosynthesis, internal CO<sub>2</sub> concentration, transpiration rate, photosynthetic water use efficiency, water use efficiency, 1-aminocyclopropane-1-carboxylic acid content and ACC oxidase were maximal in 200 $\mu$ L/L ethrel x 80kg N/ha. Plants grown with sufficient soil-applied N (80kg N/ha) responded to the ethrel (200 $\mu$ L/L). However, if soil-applied N was

less than 80kg N/ha, the effect of ethrel spray was not prominent. The increased vegetative growth due to ethrel (200 $\mu$ L/L) spray made the plant to extract more of the soil N and was reflected in N content and accumulation and increased growth, physiological, biochemical and yield characteristics. The calculated nitrogen uptake efficiency, nitrogen utilization efficiency and nitrogen-use efficiency showed that nitrogen use was better when plants were treated with the 200 $\mu$ L/L ethrel spray. Seed yield, oil yield and nitrogen yield potential were increased and were maximal with 200 $\mu$ L/L ethrel x 80kg N/ha.

Experiment 4 (1999-2000) was a factorial randomized complete block design conducted simultaneously with the Experiment 3 under non-irrigated conditions. The scheme of the treatments, ethrel spray concentrations and soil-applied nitrogen were same as in Experiment 3. The data on growth, physiological, biochemical, yield and quality characteristics recorded were those mentioned for Experiment 3. Individual effects of 200 $\mu$ L/L ethrel spray, soil-applied 80kg N/ha and their interaction proved best for most of the plant characteristics. The response of the plant to ethrel and nitrogen treatments was similar as found in Experiment 3.

Combined analysis of Experiments 3 and 4 showed that the response of the plant to ethrel spray, soil-applied nitrogen and their interaction was uniform irrespective of the irrigation conditions. The data on plant characteristics under irrigated and non-irrigated conditions were non-significant. The data suggests that 200 $\mu$ L/L ethrel spray on plants grown with soil-applied 80kg N/ha may be


used for improving mustard cultivation irrespective of the conditions of irrigation.

Experiment 5 (2000-2001) was a factorial conducted according to randomized complete block design. In this experiment applications of 0 and 200 $\mu$ L/L ethrel or 1mM silver thiosulphate were done as foliar spray at 60d after sowing (flowering stage) on mustard cultivar Alankar grown under irrigated and non-irrigated conditions. Plants were raised with uniform soil application of 80kg N/ha. This experiment was based on the findings of Experiments 3 and 4. The response of the plants to ethrel spray treatment was confirmed in this experiment with the use of silver thiosulphate spray treatment, as silver thiosulphate application inhibits ethylene action. The observations recorded at 80, 100 and 120d after sowing included growth (plant leaf area plant dry weight), physiological, biochemical, yield and quality characteristics were similar to Experiments 3 and 4. Maximum response was noted with 200 $\mu$ L/L ethrel spray treatment. However, silver thiosulphate spray inhibited the ethylene action and ethrel effect was not observed. The interaction effect of spray and irrigation was found non-significant. The results of the experiments suggest that response of the plant to ethrel treatment seen in Experiment 1–4 was manifested through the action of ethylene.

THESIS



Fed in Computer



13 JUN 2005



T5885

THE UNIVERSITY OF MICHIGAN  
LIBRARY  
ANN ARBOR, MI 48106-1500  
TEL: 734 763 7000  
WWW.LIBRARY.MICHIGAN.EDU



**Physiological Significance of Ethrel (2-Chloroethyl  
Phosphonic Acid) and Nitrogen in Relation to  
Growth and Metabolism of Mustard under  
Irrigated and Non-Irrigated Conditions**

**THESIS**

SUBMITTED FOR THE DEGREE OF

**Doctor of Philosophy**

IN

**BOTANY**

**MOHMAD RAMZAN MIR**

DEPARTMENT OF BOTANY  
ALIGARH MUSLIM UNIVERSITY  
ALIGARH (INDIA)

**2002**

11/15/15

*Dedicated to My Late uncle*

*"Who had offered his finger when I attempted to stand  
on my legs"*

*And My Parents*

*"Whose blessings have sustained me through this work"*

THESES

*Nafees A. Khan*  
Ph.D.  
Lecturer




Department of Botany  
Aligarh Muslim University  
Aligarh – 202002, India  
Off: (0571) 702016  
Fax: 91-0571-702016  
e-mail: naf9@lycos.com  
Khan\_na@rediffmail.com

Dated: 10.5.2002

### CERTIFICATE

This is to certify that the thesis entitled **Physiological Significance of Ethrel (2-Chloroethyl Phosphonic Acid) and Nitrogen in Relation to Growth and Metabolism of Mustard under Irrigated and Non-Irrigated Conditions** submitted for the degree of **Doctor of Philosophy in Botany** is a faithful record of bonafide research work carried out at the **Aligarh Muslim University, Aligarh** by Mr. **Mohmad Ramzan Mir** under my guidance and supervision and that no part of it has been submitted for any other degree or diploma.

  
(NAFEES AHMAD KHAN)



## ACKNOWLEDGEMENTS

*My all praise and thanks to the Almighty God, the Lord and the Cherisher of all the world Who taught human-kind everything that they knew not.*

*First and foremost, I must gratefully record my indebtness to all the eminent authorities, scholars and researchers whose works I have consulted and whose ideas and insights have richly contributed to give a shape to this study.*

*I feel pleasure to express my indebtness and gratitude to my respected supervisor, Dr. Nafees Ahmad Khan, Lecturer, Department of Botany, Aligarh Muslim University, Aligarh for his keen interest, enthusiastic support, valuable guidance and constant supervision. Despite his busy schedule, he gave me liberty to encroach upon his valuable time to complete my work.*

*I must register my sincere thanks to Prof. Samiullah, Chairman and Late Prof. M. Wajid Khan, Former Chairman, Department of Botany, Aligarh Muslim University, Aligarh for providing necessary facilities during the work.*

*I would like to place on record my profound thanks to teachers, Drs. Aqil Ahmad, Arif Inam, Firoz Mohammad and M.M.A. Khan for their support.*

*I have no words to express my thanks to Drs. Moinuddin Khan, Shamsul Hayat, Mohd. Mobin, Arshad Hussain; Messers M. Zaki Azam, Qazi Faridudin, Sheikh Javid, Fayaz A. Sheikh, Showkat Hussain, Irfan Ahmad,*

*Perveiz M. Lone, Ms. Afroza Akhtar, Ms. Rotuba Shaheen, Ms. Shahla Saeed, Ms. Lalita Gupta and all other research scholars of Plant Physiology Section for their well timely help and support in every respect, without which the task was impossible.*

*I would miss my duty if I do not make a mention of Dr. Nasir Ahmad Lone for his unforgettable help and encouragement.*

*The acknowledgement will remain incomplete without heartiest thanks to all my friends who provided me motivation, encouragement and moral support in completing this work. Thanks are due to Drs. Anees Ahmad, Jahangeer Iqbal, Mehrajudin and Messers Syed A. Untoo, Mushtaq Ahmad, Ab. Hamid Mir, Aamil Ahmad and also to Dr. Amina Mir and Ms. Ulfat Jabeen, all of them have contributed a lot towards a fruitful completion of the work.*

*I am also grateful to Mr. Rais Ahmad Khan for extending adept services on computer for typing the manuscript.*

*To save the best for last, I thank my parents, brothers, sisters and all the other family members for their love, support and constant encouragement without which this work would have not existed.*

*Mohd Ramzan*  
(Mohmad Ramzan Mir)

## **CONTENTS**

<b>INTRODUCTION</b>	<b>1</b>
<b>REVIEW OF LITERATURE</b>	<b>5</b>
<b>MATERIALS AND METHODS</b>	<b>47</b>
<b>EXPERIMENTAL RESULTS</b>	<b>68</b>
<b>DISCUSSION</b>	<b>100</b>
<b>SUMMARY</b>	<b>117</b>
<b>BIBLIOGRAPHY</b>	<b>123</b>
<b>APPENDIX</b>	<b>i</b>
<b>TABLES</b>	

# ***CHAPTER-1***

## ***INTRODUCTION***

### INTRODUCTION

Agriculture is backbone for the development of Indian economy. It contributes nearly 30 per cent of domestic production and employs about 70 per cent of the people. The Government of India is aimed at achieving and maintaining agricultural growth rate of 4 per cent. Among several crops grown, oilseeds occupy an important place in agricultural produce all over the world, so as in Indian agriculture. Oilseeds hold about 13 per cent of country's gross cropped area, contributing 10 per cent of all agricultural products and 5 per cent to Gross National Productivity. Oilseeds cultivation is carried out across the country in an area of about 26 million hectares. Rapeseed is cultivated in 3.73 million hectares in India, which keeps it to the third position in the world just behind the Canada and China.

Four major crops, soybean, sunflower, rape and palm provide over 70 per cent of the vegetable oil and cottonseed, coconut seed and groundnut supply the remainder. Oilseed *Brassica* accounts for approximately 10 per cent of the total world oilseed production and 14-15 per cent of the total vegetable oil production (Downey and Rimmer, 1993). There are few non-conventional sources of oil, which are employed to increase the resources of vegetable oil. Most important of them are rice bran and maize germ.

The importance of the oilseed is because of fats contained in their seeds, which provide 2.5 times more calories than carbohydrates. Fat is also an important constituent of cell membrane and serves as a precursor for a variety of biologically active compounds collectively known as eicosanoids (e.g. prostaglandins, thromboxanes and leukotrienes). It also helps in transport of fat-soluble vitamins, A, D, E and K in human body (Anonymous, 1993). Most vegetable oil is used directly in human diet as cooking oil and margarine or indirectly processed products such as shortening and

confectionery. Besides the use of oil, oilseeds by-products such as protein meal from seeds and hulls of seeds are used as animal feedstuff and in making of insulation boards. About 20–30 per cent of oil is used for non-edible purposes, e.g. in the production of soaps, detergents, lubricants, plastics, resins, paints, varnishes, cosmetics and as precursors to a wide range of chemicals.

India recorded a spectacular increase both in area and production of oilseeds during the last few decades. The production increased from 11 million tonnes to 22 million tonnes during the last decade in an effort called Yellow Revolution. However, considering the area and production of oilseeds in world's scenario, India produces only about 7 per cent of world's edible oil with about 35 per cent of the world's area under oilseed cultivation. With this limited area and production of oilseeds, it has to feed about 16 per cent of the total world's population. To meet the daily requirements of oils and fats of 18g per day, India has to import edible oil every year on large scale, which causes burden on our limited foreign resources.

The oilseeds production in India is restricted due to several factors. For example: (i) more than 75 per cent of the farmers have small or marginal holdings of less than two hectares (ii) only 15 per cent of area under oilseed is irrigated as compared to 72 per cent under wheat and 44 per cent under rice (iii) absence in advancement of agricultural techniques for high yielding varieties, post harvest technology and proper processing facilities (iv) no genetically modified oilseeds are released for cultivation (v) lower number of flowers (about 68 per cent) develops into pods (vi) pod shattering is a particular problem in oilseed rape, which accounts to an average annual losses of about 20 per cent (Downey and Robbelen, 1989). Seeds may be lost to the extent of 10,000 seeds/m<sup>2</sup> (Lutman, 1993). (vii) hormonal imbalance during sink development, which may be the cause of flower drop and pod shattering and improper source-sink relationship.

With the limitation in increasing the land under cultivation, the focus, thus is on increasing the productivity of available land.

Nitrogen affects favourably growth, development and yield of crops (Marschner, 1986; Khan *et al.*, 1990; Samiullah *et al.*, 1990; Khan *et al.*, 2000; Dodd, 2001; Khan *et al.*, 2001) as this nutrient forms the building blocks of macromolecules in cell (Salisbury and Ross, 1994). Nitrogen application brings about increase in water use efficiency through increase in soil moisture extraction at sub soil surface and increase in growth at upper ground level (Saran and Giri, 1990; Vyas *et al.*, 1995; Thakral *et al.*, 1997; Zaman and Choudhri, 1998; Dodd, 2001). However, continuous application of nitrogen has environmental as well as economic limitations. For example, about half of the nitrogen unavailable to the crops (Greenwood, 1982) remains in the soil and washed away with water to increase the biological productivity of the water bodies causing a well-known eutrophication phenomenon. Moreover, excessive use of nitrogen exerts a pressure on its production and involves possibility of its high prices. Besides subsidy on nitrogen by the Government, a majority of small crop growers (75 per cent) still face difficulty in procuring the nitrogenous fertilizers at appropriate time. It is, therefore, a challenge before the research scientists to find out options, which reduce the input of nitrogen without affecting seed yield. In this regard, methods for better management of applied nitrogen through higher N-use efficiency may be an answer. If physiological basis of nitrogen utilization efficiency with increase in the sink strength and better source-sink relationship is developed, then problems discussed earlier would find solution and physiological traits may be developed as a biochemical tool. In the preceding pages, it was mentioned that lower number of flowers (about 68 per cent) develops into pods due to hormonal imbalance. The use of plant hormones as foliar spray at appropriate stages has shown to affect flowering and pod formation (Wareing and Phillips, 1981; Cheema *et al.*, 1987; Rajala and Peltonen-Sainio, 2001). Further, researches

carried out in the author's laboratory have shown that plant growth regulator, gibberellic acid, has potential of utilization of nitrogen, phosphorus, potassium and sulphur from the soil, increasing the growth and crop productivity (Ansari, 1996, Khan, 1996a, Khan *et al* , 1996, Khan, 1997, Khan *et al* , 1997, 1998, 1999, Mobin, 1999, Khan *et al* , 2001) Few studies on ethrel have also shown that it brings about increase in photosynthetic capacity and yield of mustard (Khan, 1996b, 1998, Khan *et al* , 2000) Other workers have also documented role and regulation of ethylene in plant growth and development (Abeles *et al* , 1992, Mattoo and White, 1991, Mattoo *et al* , 1997) In the present thesis, efforts were made for extensive examination of the role of ethylene (applied as ethrel) in physiology and metabolism of nitrogen in mustard (*Brassica juncea* L Czern & Coss ) Five field experiments were conducted to accomplish the following objectives

- 1 To study the effect of leaf-applied ethrel on growth, physiology, yield and quality characteristics under irrigated and non-irrigated conditions
- 2 To study the effect of leaf-applied ethrel on growth and physiology of mustard grown with nitrogen levels
- 3 To study the relationship of nitrogen utilization efficiency of mustard with ethylene evolution
- 4 To test the hypothesis that ethylene has a central role in mediating plant responses with the use of ethylene action inhibitor silver thiosulphate



***CHAPTER-2***  
***REVIEW OF LITERATURE***

## CONTENTS

### REVIEW OF LITERATURE

2 1	Introduction	5
2 2	Plant Growth Regulators	10
2 3	Ethylene as Plant Growth Regulator	11
2 3 1	Ethylene	12
2 4	Crop Response to Ethylene Sources	14
2 4 1	Ethylene sources and growth parameters	14
2 4 1 1	Plant height	14
2 4 1 2	Leaf number	15
2 4 1 3	Leaf area	15
2 4 1 4	Leaf area index	15
2 4 1 5	Dry matter	16
2 4 2	Ethylene sources and photosynthetic parameters	16
2 4 2 1	Chlorophyll	16
2 4 2 2	Photosynthesis	16
2 4 2 3	Photosynthetically active radiation	17
2 4 3	Ethylene sources and nutrient uptake	17
2 4 4	Ethylene sources and yield parameters	18
2 4 4 1	Pod number	18
2 4 4 2	Seed number	19
2 4 4 3	1000 seed weight	19
2 4 4 4	Seed yield	19
2 4 4 5	Biological yield	20
2 4 4 6	Harvest index	20
2 4 4 7	Oil yield	21
2 4 5	Ethylene sources and quality parameters	21
2 4 5 1	Oil content	21
2 4 5 2	Amino acid content	21
2 4 5 3	Protein content	21
2 4 5 4	Fatty acid	22
2 5	Mineral Nutrition	22
2 5 1	Nitrogen	23
2 5 1 1	Acquisition and assimilation	23
2 5 1 2	Nitrogen source	24
2 5 1 3	Mineralization	25
2 5 1 4	Nitrogen immobilization	26
2 5 1 5	Reasons for nitrogen deficiency	26
2 5 1 6	Denitrification loss	27
2 5 1 7	Gaseous loss of nitrogen	27
2 5 1 8	Control of mineral fertilizer N loss in soil	28

2 5 1 9	Consequences of nitrogen deficiency	29
2 6	Crop Response to Nitrogen	30
2 6 1	Nitrogen and growth parameters	30
2 6 1 1	Plant height	30
2 6 1 2	Leaf number	31
2 6 1 3	Leaf area	31
2 6 1 4	Leaf area index	31
2 6 1 5	Dry matter	32
2 6 1 6	Crop growth rate	33
2 6 1 7	Leaf area duration	33
2 6 1 8	Relative growth rate	34
2 6 1 9	Net assimilation rate	34
2 6 2	Nitrogen and photosynthetic parameters	34
2 6 2 1	Chlorophyll	34
2 6 2 2	Photosynthesis	34
2 6 3	Nitrogen and nutrient uptake	35
2 6 4	Nitrogen and yield parameters	37
2 6 4 1	Pod number	37
2 6 4 2	Seed number	37
2 6 4 3	1000 seed weight	38
2 6 4 4	Seed yield	38
2 6 4 5	Harvest index	39
2 6 4 6	Oil yield	39
2 7 5	Nitrogen and quality parameters	40
2 7 5 1	Oil content	40
2 7 5 2	Aminoacids	41
2 7 5 3	Protein content	41
2 7 5 4	Fatty acids and glucosinolate	42
2 7 5 5	Acidity of oil/iodine number	43
2 8	Crop Response to Interaction between Ethylene Sources and Nitrogen	43
2 8 1	Interaction effect on growth parameters	43
2 8 2	Interaction effect on photosynthetic parameters	44
2 8 2 1	Chlorophyll content	44
2 8 2 2	Photosynthesis	44
2 8 2 3	Photosynthetic active radiation	45
2 8 3	Interaction effect on nutrient uptake	45
2 8 4	Interaction effect on yield parameters	45
2 9	Concluding Remarks	46

## **REVIEW OF LITERATURE**

### **2.1 Introduction**

Indian agriculture has progressed a long way to become a significant exporter of a diversified basket of agricultural commodities. Persistent efforts have been made to harvest a larger portion of the land to agricultural purpose and for induction of new technology. In post-independent India, agriculture has been growing at a rate of about three per cent per annum, while food grains production is increasing by 1.67 per cent, which falls short of population growth of 1.80 per cent.

A spectacular rise in food grains production has been observed since 1949-50. It is estimated that the production of cereal has gone to 206 million tonnes during 1999-2000. If acreage production of cereal crops is considered, rice ranks first. With an area of 44 million hectare that produces 88 million, this crop is a major contributor amongst cereal crops. Wheat and other cereal crops occupy lower position in terms of production.

The Indian oilseed scenario has also undergone a significant transformation due to contributions of oilseeds production technology, expansion in cultivated area, price support policy and institutional support. In particular, the set up of the Technology Mission on Oilseed in 1986 played a major role. The role of technology in fostering and sustaining the oilseeds production is vast. The self-sufficiency in oilseed has been possible only due to economically viable and sustainable technology. Since oilseeds are considered as hardy crops, they are mostly grown under moisture and nutrient scarce conditions that may limit the crop productivity. In India, agro-ecological conditions are such which favour growth of nine oilseeds crops. These are seven edible oilseeds, namely groundnut, rapeseed-mustard, soybean,

sunflower, safflower, sesame and niger and two non-edible castor and linseed, apart from a wide range of other minor oilseeds and oil bearing tree species.

With a production of 24 million tonnes during 1996-97, the country accounted for 9.7 per cent of the global oilseeds production (about 250 million tonnes). Indian vegetable oil industry achieved domestic turnover of Rs. 35,000 crores compared to international trade in import of vegetable oil and exports of oil meals and castor oil, which accounts to Rs. 65,000 crores.

The expansion in area under oilseeds has been main reason of increase in oilseeds production, but further scope of area expansion is meagre. Therefore, the productivity growth should take a lead in bridging the demand and supply gap. However, there are several constraints in increasing the productivity of oilseeds. These may be biotic and abiotic, on-farm and off-farm, technological and non-technological. With about 25 per cent of the area under irrigation, oilseeds are subjected to vagaries of monsoon resulting in lower yields compared to irrigated wheat and rice. This warrants limited and protective irrigation at least during critical stages of crop growth. Besides, several other constraints like continuous cultivation of oilseeds without proper crop rotation, low level of management adopted by small and marginal farmers, poor post-harvest technology and inadequate marketing support and weak technology transfer add to the problems of low crop productivity. Researches are done to overcome most of the constraints related to technology, in term of genetic improvement, efficient agronomic management, pest management, and post-harvest technology and to improve oilseeds crop production. As per study based on the data available from the Frontline Demonstration Project under the Oilseeds Production Programme, the mean yield of improved technology of oilseeds realizable on farmers' field is 1545 kg/ha, while the national average yield is of the order of 794 kg/ha. There exists a realizable yield gap of 751 kg/ha, which accounts for nearly about 95 per cent of the existing national average yield. In other words, an additional production of the order 16.27

million tonnes need to be added to the current oilseeds production in the country from the existing area under oilseeds. Oilseeds remained almost neglected during the green revolution phase in India in terms of total output

The languid production of oilseeds has converged the attention of agricultural scientists to innovate and implement the improvised methods for boosting the yield through proper nutrient management. In this context, *inorganic nutrients (concerned with acquisition of mineral elements from the soil)* found to be immensely remarkable. The green revolution of India owes much to this form of nutrition. Now, when we are on the verge of entering in to the era of Ever Green Revolution, the investigations regarding application of nutrients need to be thoughtfully considered, which has largely been ignored

It may be emphasized that work carried out in relation to nutrient is voluminous and it is known that application of only few essential nutrients like, N, P and K alter the availability and uptake of other nutrients. Among them nitrogen is a major nutrient and its availability in the rhizosphere has been found to enhance the yield and quality of seed crops (Ansari, 1990; Jeschke *et al.*, 1992, Awasthi and Surajbhan, 1994, Sarkar *et al.*, 1999; Khan *et al.*, 2000, Dodd, 2001)

Water plays an important role in physiological processes of plants and its inadequacy results in depressed crop productivity around the world (Ludlow and Muchow, 1990, Nilsen and Orcutt, 1996, Singh *et al.*, 1996). Oilseed *Brassicas* (rapeseed and mustard) are generally grown on conserved soil moisture from monsoon rain, which is progressively depleted with the advancement of growing season. Decline in relative water content because of reduced water availability brings about changes in morphological plant characteristics (Kumar and Bharti, 1988; Nilsen and Orcutt, 1996). A decrease in the rate of photosynthesis in leaves has been attributed to both stomatal and non-stomatal limitations (Graan and Boyer, 1990; Lauer and Boyer, 1992). Decreased photosynthetic rate is associated with a reduction in protoplast

and/or chloroplast volume (Santakumari and Berkowitz, 1990). Under low plant water status leaf chlorophyll content and stomatal conductance decreases causing reduction in photosynthetic rate. The net rate of CO<sub>2</sub> assimilation decreased as water stress developed in wheat (*T. aestivum*) (Lu and Zhang, 1999). Decrease in photosynthetic capacity during water stress in *Brassica juncea* could be ascribed to restricted CO<sub>2</sub> availability as a consequence of stomatal resistance to gas exchange (Sawhney *et al.*, 1996).

The area of the leaves developed to lesser extent when plants are exposed to low water status (Lu and Neumann, 1998; Pankovic *et al.*, 1999). Kumar and Elston (1993) reported that leaf area was reduced in both species of rape and Indian mustard under low water regime. Similar results were reported by Burke *et al.* (1985), Kumar *et al.* (1994), Sardadevi *et al.* (1996). Leaf area index, crop cycle length and phenology determine the total dry matter production, yield and yield components under drier environments. Shoot is found to more sensitive to water than root growth in *Brassica juncea* (Pannu *et al.*, 1992; Sharma *et al.*, 1994).

Water availability affects the process of cell expansion via physical and metabolic changes (Taiz, 1984; Nonami *et al.*, 1997). Reduced rates of new cell production may make additional contributions to the inhibition of growth (Silk, 1992; Durand *et al.*, 1995; Lecoeur *et al.*, 1995). Decreased water availability reduced leaf area index, relative water content, xylem potential, seed yield and harvest index in mustard (Sharma, 1992).

Nutrient uptake of plants decreased under water stress condition due to reduced transpiration. A decline in soil moisture has also been associated with a decrease in nutrient uptake (Tanguilig *et al.*, 1987) primarily caused by the physiological impairment of the active nutrient absorption and transport mechanism of the roots.

Amino acid accumulation has been shown to occur in a variety of monocots and dicots. One of the best-characterized osmoregulatory responses

is the accumulation of proline. Water stress enhanced the proline content in mustard, wheat and *Plantago*. To maintain uptake of water and nutrients under limiting water conditions, continuous proliferation of roots into new soil layer is important at the cell and tissue level (Nilsen and Orcutt, 1996).

Drought stress resulted in reduction in seed yield and was due to differential responses of yield contributing traits in species of rapeseed-mustard (*Brassica campestris* and *Brassica napus*) (Mathur and Wattal, 1996). Kumar *et al* (1994) reported that the seed yield reduction by water stress was because of decrease in the number of pods per plant and number of seeds per pod.

Fertilizer application has been found to bring about marked improvement in water use efficiency with increasing fertility levels (Singh and Srivastava, 1996, Thakral *et al*, 1997). The role of N in photosynthetic capacity has long been recognized. Nitrogen ensures the build up of chlorophyll (Evans and Terashima, 1988, Sage *et al*, 1990, Liu and Dickman, 1992) and increases the amount and activity of enzymes for leaf nitrogen content. Sugiharto *et al* (1992) found a significant positive correlation between the photosynthetic capacity of leaves and their leaf nitrogen used for synthesis of components of photosynthetic apparatus.

Nitrogen application enhanced soil moisture extraction and increased water use efficiency during larger leaf area without affecting water use during reproductive phase in mustard under different levels of stored soil moisture (Saran and Giri, 1990, Vyas *et al* 1995).

Water potential, relative water content, photosynthesis and concentration of photosynthetic pigments are known to be influenced by growth regulators under stress conditions (Kumar and Bharti, 1988, Cliquet *et al*, 1991, Sairam *et al*, 1991, Dayal *et al*, 1993). Growth regulators application under moisture stress also helps in preserving water balance of the plant by closing the stomata and canopy modification and thereby reducing the rate of transpiration which ultimately results in protecting the enzyme proteins and cell organelles for



biochemical activities (Leitch and Kuat, 1999, Sanvicente *et al* , 1999) Along with the proper nutrition and water management, hormonal balance if made at particular developmental phase this would be an additional benefit for crop yield A group of chemicals that modify hormonal balance is thought to be plant growth regulators

Keeping in view the specific nature of the problem of the thesis, a general account of the plant growth regulators has been given with special emphasis of ethylene and impact of ethylene releasing compounds on aspects of growth and development of plants

## 2.2 Plant Growth Regulators

The definition of phytohormones accepted by most plant physiologists is similar to that developed for animal hormones Pincus and Thimann (1948) defined it as *a plant hormone is an organic compound synthesized in one part of a plant and translocated to another part, wherein very low concentrations it causes a physiological response* and to distinguish it from animal hormone it was named phytohormone and the substances which regulate growth are called 'growth regulators' (Sircar, 1971) Hormones are effective at internal concentrations of about  $1\mu\text{M}$ , whereas other metabolites necessary for growth and development are usually present at concentration 1 to  $50\mu\text{M}$  The precise location of synthesis of phytohormones is uncertain but actively growing leaves, fruits and developing seeds are thought to be active sites of synthesis of phytohormones However, it appears that all tissues have potential to produce any of the phytohormones, which are transported via the xylem or phloem and by diffusion such as in the case of ethylene They occur in plants as free and conjugated forms The latter are conjugates of sugars, amino acids and possibly peptides The free forms are generally considered to be biologically active, while conjugates are viewed as functioning in controlling levels of the more active forms of transport and storage

The plant hormones are extremely important agent in the integration of developmental activities. Environmental factors often exert inductive efforts by evoking changes in hormone metabolism and distribution within the plant. Apart from this, they also regulate expression of the intrinsic genetic potential of plants. The mechanism of action of phytohormones is poorly understood. However, several mechanisms or combinations may be operative. Control of genetic expression has been demonstrated for the phytohormones at the transcription and translational levels. Also, hormones receptors binding proteins have been identified on membrane surface that are specific for some phytohormones. The type and abundance of these proteins appear to be important in determining the sensitivity of the tissues to phytohormones differing in concentrations.

The chemical control of plant growth and through the use of a plant growth hormone is a common practice to make a plant more commercially acceptable. A number of synthetic compounds control shoot growth in higher plants without being phytotoxic or causing malformation or damage (Cathey, 1964; Salisbury and Ross, 1994). Some of these substances have found application in agricultural practice (Nickell, 1982; Salisbury and Ross, 1994). They have been shown to be involved in the regulation of photosynthesis and the movement of the photosynthetic products from their site of synthesis in the leaf to their site of accumulation (Patrick, 1982; Thomas, 1986; Patrick and Steains, 1987; Pereto and Beltran, 1987; Khan, 1996a, b; Khan *et al.*, 1997; 1998; 2000; 2001).


### **2.3 Ethylene as Plant Growth Regulator**

The most commonly used and best understood group of plant growth hormones consist of those which regulate the production of ethylene (Yang and Hoffman, 1984; Abeles *et al.*, 1992). Ethylene is believed to control every aspect of plant growth and development.

### 2.3.1 Ethylene

The Russian botanist Neljubow (1901) was the first to recognize the growth regulatory properties of ethylene. Later on in 1930 it was recognized to have a wide variety of effects on plants and it was until 1934, when Gane in England first obtained positive proof that ethylene was a natural plant product.

Ethylene (ethane according to the IUPAC system of nomenclature) has the formula  $\text{H}_2\text{C} = \text{CH}_2$  (MW 28.05). It is a gas (m.p =  $-169.15^\circ\text{C}$ ; b.p =  $-103.71^\circ\text{C}$ ). It is colourless with ether like smell and is lighter than air (density of ethylene at  $0^\circ\text{C}$  and 1 atmosphere =  $1.260\text{g L}^{-1}$ ; density of dry air at  $0^\circ\text{C}$  and 1 atmosphere =  $1.293\text{g L}^{-1}$ ). It is highly flammable. It is more soluble in water than  $\text{O}_2$  or  $\text{N}_2$  but less than  $\text{CO}_2$ .




*Structural formula of ethylene*

Ethylene is the simplest olefin, which exists in the gaseous state under normal physiological conditions. Its effect on various physiological processes at different stages of plant growth and development has been documented (Yang and Hoffmann, 1984; Reid, 1987; Abeles *et al.*, 1992; Pua and Chi, 1993; Khan, 1996b and Khan *et al.*, 2000). Ethylene is known to exert its effects by altering gene expression both at transcriptional and post-transcriptional phases (Lincoln and Fischer, 1988). Dependant on the plant material and state of development, promoting or inhibiting effects of ethylene on internode growth has been observed (Raskin and Kende, 1984; Sisler and Yang, 1984). Physiological conditions like water stress or drought also promote the ethylene synthesis in plants (Tudela and Primo-Millo, 1992; Bergner and Teichmann, 1993; Michelozzi *et al.*, 1995).

Ethylene generating commercial chemical is ethephon or ethrel (2-chloroethyl phosphonic acid –  $\text{ClCH}_2\text{CH}_2\text{PO}_3\text{H}_2$ ) (Warner and Leopold, 1969).

Ethrel is most important and versatile ethylene- releasing agent marketed and registered for more than 20 crops. It is a synthetic plant growth regulator that undergoes chemical biodegradation in cell cytoplasm at pH greater than 4.1 to release ethylene (Urwiller and Stutte, 1986; Kasele *et al.*, 1995).



*2-chloroethyl phosphonic acid decomposes spontaneously in plant tissues to yield ethylene and phosphonic acid*

The use of ethrel as a growth regulator has been shown for controlling growth and productivity of cereals and grain crops (Davis and Curry, 1991). Ethrel has also been found to impart tolerance against water stress and increasing the productivity of oilseed crop (Khan *et al.*, 2000). The other ethylene releasing compounds are etaelasil, holoethylesulfonic acids, 2-Hydroethylhydrazine (Palmer *et al.*, 1967; Dowlet and Kumanoto, 1972; Lurssen, 1982; Arteca, 1997).

Ethylene is a gaseous plant growth regulator involved in a diverse array of cellular developmental and stress related processes in plants. Prevention of ethylene accumulation in the atmosphere and inhibition of its effects by lowering the temperature and increasing the CO<sub>2</sub> concentration are wide spread storage practices. Silver ion also inhibits the ethylene action, acting at the receptor level (Veen, 1986), is a useful laboratory tool (Yang, 1987; Smalle *et al.*, 1997). Cobalt has been found to inhibit ethylene production and reversed

the effects of ethylene in soybean (Samimy, 1978). Mohan Ram and Rina (1982) reported the antagonistic properties of  $\text{AgNO}_3$  and  $\text{COCl}_2$  on ethylene and application of these compounds reverse the effects of ethylene in *Canabis sativa*. Mhatre *et al.* (1998) also reported the similar results. Silver thiosulphate is an inhibitor of ethylene action (Saniewske and Ludhika, 1989) and aminoethoxy vinylglycine inhibits the biosynthesis of ethylene (Child *et al.*, 1998; Kushad and Pooviah, 1984; Koritsas, 1988). Cyclopropenes and 2,5-norbornadiene have also been found to be effective antagonists of the ethylene response (Sisler and Yang, 1984; Sisler *et al.*, 1996a, b).

## **2.4 Crop Response to Ethylene Sources**

### **2.4.1 Ethylene sources and growth parameters**

#### **2.4.1.1 Plant height**

Growth regulators have been found to control plant height (Sauerbrey *et al.*, 1988; Guruprasad and Guruprasad, 1988; Dijkstra and Kuiper, 1989; Krishnamoorthy, 1993).

Depending on the plant material and state of development, promoting or inhibiting effects of ethylene on internode growth have been observed (Raskin and Kende, 1984; Sisler and Yang, 1984). Ethephon appreciably reduced the shoot length of sunflower plants and the internode elongation (Sauerbrey *et al.*, 1988). Slife and Earley (1970) applied ethrel to flowering soybean plants at 0.56 to 2.24 kg/ha rates and all the treatments caused a decrease in plant height. Foliar application of ethrel at a rate of 500, 1000 and 1500  $\mu\text{L/L}$  reduced plant height in *Brassica napus* (Grewal *et al.*, 1993). Ethephon has been found to reduce plant height in barley (Bulman and Smith, 1993a; Sanvicente *et al.*, 1999), sunflower (Sauerbrey *et al.*, 1988), winter rape (Wareing and Philips, 1981), rice (Nafziger *et al.*, 1986), winter wheat (Van Sanford *et al.*, 1989), lupin (Ortuno *et al.*, 1993), linseed (Leitch and Kuat, 1999), radish (Vreugdenhil and Harro, 1989) and arabidopsis seedlings (Smalle *et al.*, 1997).

Contrarily, Jana and Kabir (1991) reported that application of ethrel to cauliflower cv. Dania significantly increased plant height at 300  $\mu\text{L/L}$  concentration. However, growth was adversely affected at higher concentration.

#### **2.4.1.2 Leaf number**

Ramos *et al.* (1989) observed that ethephon application at tillering increased both number of ears per plant and per pot in spring barley (*Hordeum vulgare*). Ethephon was found to be beneficial for increasing the number of leaves per plant of onion (*Allium cepa* L.).

Researchers have observed controlling effects of ethrel on leaf expansion and production of plants with darker green foliage (Shanahan and Nielsen, 1987; Davis *et al.*, 1988; Butler *et al.*, 1989; Sairam *et al.*, 1989; Reddy *et al.*, 1996; Zhou and Xi, 1993; Kulkarni *et al.*, 1995; Zhou and Ye, 1996; Lee and Reid, 1997; Hussain *et al.*, 1999).

#### **2.4.1.3 Leaf area**

Ethylene has been shown to influence leaf expansion by suppressing cell enlargement rather than division (Kieber *et al.*, 1993; Rodriguez-Pousida *et al.*, 1993). Ethephon treatment reduced leaf area as compared to control plants in *Zea mays* (Kasele *et al.*, 1995). However, ethephon application promoted expansion of primary leaves of *Helianthus annuus* while higher concentrations reduced it (Lee and Reid, 1997).

#### **2.4.1.4 Leaf area index**

Singh *et al.* (1987) working on soybean and Grewal and Kolar (1990) on mustard reported an increase in LAI in soybean by the application of ethrel. Flag leaf area index was greater in wheat treated with ethephon over control but plant leaf area index was not affected by ethephon application (Van Sanford *et al.*, 1989). Khan (1996b) and Khan *et al.* (2000) also reported an increase in leaf area index in response to ethrel spray in *Brassica juncea* L. under irrigated and non-irrigated conditions.

#### 2.4.1.5 Dry matter

The plant growth regulator ethephon influenced the dry matter significantly in winter wheat (Nafziger *et al.*, 1986; Van Sanford *et al.*, 1989), in barley (Simmons *et al.*, 1988) and in mustard (Khan, 1996b; Khan *et al.*, 2000). The dry weights of the main stem and root of ethrel treated mungbean plants were significantly higher (Panwar *et al.*, 1988). Contrary to these reports, Urwiller and Stutte (1986) noted decrease in dry weight of soybean plants due to ethephon treatment.

#### 2.4.2 Ethylene sources and photosynthetic parameters

##### 2.4.2.1 Chlorophyll

Favourable effects of growth regulators on chlorophyll content have been reported by number of workers. Ethrel at 500  $\mu\text{L/L}$  concentration significantly increased the chlorophyll content in leaves of *Brassica napus*. However, at higher doses of ethrel (1000 and 1500  $\mu\text{L/L}$ ) showed detrimental effect (Grewal and Kolar, 1990; Grewal *et al.*, 1993). The exogenous application of ethylene in mustard (*Sinapis alba* L.) seedlings enhanced chlorophyll synthesis considerably (Buehler *et al.*, 1978). The ethylene producers (Camposan M, Ethephon and Dektrel) did not cause chlorosis and reduced green pigment level to a lesser degree in developing raspberry leaves.

##### 2.4.2.2 Photosynthesis

Growth regulators may be employed to improve the physiological efficiency of plants by modifying the balance between photosynthesis and respiration (Arteca and Dong, 1981; Zerbe and Wild, 1981; Makeev *et al.*, 1992). The photosynthetic response of crop to the agro-chemicals varies from one extreme to the other. Increased rates of photosynthesis per unit leaf area have been observed after the application of growth regulators on different plant species (Child *et al.*, 1985; Liu *et al.*, 1993; Yang *et al.*, 1994). Plant growth regulators can affect photosynthetic  $\text{CO}_2$  uptake either by effecting stomatal

aperture or by affecting the activity of photosynthetic enzymes (Foroutan-Pour *et al.*, 1997).

Foliar application of the ethephon to spring barley caused an increase in penultimate leaf photosynthetic rate (Pua and Chi, 1993). In one of the study of Subrahmanyam and Rathore (1992a), they found that ethrel application had no significant effect on photosynthesis in Indian mustard. However, photosynthesis in upper leaves and to a lesser extent in lower leaves was lowered by ethrel application. Exogenous application of ethrel has also been found to cause 12-18% reduction in photosynthesis (Subrahmanyam and Rathore, 1992a, b).

Studies conducted in the author's laboratory have shown that exogenous application of ethrel enhanced photosynthesis of *Brassica juncea* in irrigated and unirrigated conditions (Khan, 1996b; Khan, 1998; Khan *et al.*, 2000).

#### **2.4.2.3 Photosynthetically active radiation**

Photosynthetically active radiation (PAR) is a measure of radiation available for photosynthesis. It is well known that plants vary in response of the photosynthetic apparatus to radiations of different wavelengths within the canopy. Changes in radiation quality are largely due to the spectral properties of leaf pigments leading to a reduction in the red/far red ratio as light penetrates the canopy (Holmes, 1981; Ballare *et al.*, 1989; Guiamet *et al.*, 1989). In this respect there are several evidences that potentiate the chains of canopy change under the influence of growth regulators (Nickell, 1982), which bring about a desirable modifications in photosynthetically active radiation (Mathias *et al.*, 1989). Grewal and Kolar (1990) in their experiment on *Brassica napus* reported that application of ethrel (500, 1000 and 1500  $\mu\text{L/L}$ ) had negative impact on photosynthetically active radiation interception.

#### **2.4.3 Ethylene sources and nutrient uptake**

Plant growth regulators are known to influence transport and transport functions. Growth regulators have been affiliated with refinement of



assimilated translocation in established sink-source system (Thomas, 1986; Patrick and Steains, 1987). Desirable increase in the produce of field crops was probably due to alteration in the trends of assimilated distribution (Ado Quaye *et al.*, 1986). The allocation of newly fixed carbon in different metabolic products influenced the partitioning of carbon and growth activity of whole plant (Champigny, 1985).

Ethephon showed a strong effect on N-use efficiency and in particular on the role of N-uptake efficiency in winter wheat (Van Sanford *et al.*, 1989). Use of ethephon resulted in an increased N uptake in barley (Bulman and Smith, 1993a).

Contrarily, Dhakal and Erdi (1986) found that ethelene had no influence on K and Na levels neither at lower nor at higher concentrations in wheat.

In a field trial on mustard under irrigated conditions (Khan, 1998) and under non-irrigated conditions, Khan *et al.* (2000) reported that ethrel spray plants accumulated higher plant N and seed N content, and enhanced nitrogen harvest index and nitrogen yield merit (Khan, 1998). Under non-irrigated conditions ethrel-sprayed plants utilized N from the soil more efficiently and showed increased nitrogen harvest index and nitrogen yield merit.

#### **2.4.4 Ethylene sources and yield parameters**

Yield components like pod number, seeds per pod and seed specific weight not only depend on nutritional factor but also on hormonal concentration (Morgan, 1980; Crosby *et al.*, 1981; Carlson *et al.*, 1987; De-Bouille *et al.*, 1989; Paulpandi *et al.*, 1998).

##### **2.4.4.1 Pod number**

Foliar application of ethrel at a rate of 200  $\mu\text{L/L}$  at flower- initiation stage improved the number of pods per plant in soybean (Singh *et al.*, 1987). Foliar spray of ethrel increased the number of mature pods per plant of peanut (*Arachis hypogea*) positively with sequential spray treatment over no spray in all the varieties (Saini *et al.*, 1984).

Urwiller and Stutte (1986) reported increased number of one-seeded pods in ethephon treated soybean plants. However, exogenous application of ethrel resulted in detrimental effect on pods per plant in *Brassica napus* (Grewal *et al.*, 1993). Results reported from authors' laboratory have confirmed the beneficial effects of ethrel on pod number of mustard (*Brassica juncea* L.) under irrigated (Khan, 1996b; 1998) and non-irrigated (Khan *et al.*, 2000) conditions.

#### 2.4.4.2 Seed number

Pod number and seed number per pod are determined early after flowering (Pechan and Morgan, 1983) and though nutritional factors play an important role (Allen and Morgan, 1972; Tayo and Morgan, 1978), there is evidence that hormones may also operate on these yield components. Foliar application of ethrel at flowering and pegging stages increased number of seeds and size of seeds in groundnut (Mishra *et al.*, 1984).

#### 2.4.4.3 1000 seed weight

Foliar application of ethrel increased 1000-grain weight in soybean (Singh *et al.*, 1987). Spray at flowering stage on Indian mustard also increased seed weight (Singh and Kumar, 1991). Ethephon increased 1000 seed weight of onion (*Allium cepa* L.) compared to control (Singh *et al.*, 1995). However, Grewal *et al.* (1993) observed that ethrel treatment resulted in detrimental effect on 1000 seed weight in *Brassica napus*, while Ramos *et al.* (1989) in barley (*Hordeum vulgare* L.) Khan *et al.* (2000) in mustard (*Brassica juncea* L.) Coffelt and Howell (1986) in peanut (*Arachis hypogea* L.) failed to observe any increase in 1000 seed weight in response to the foliar spray of ethrel.

#### 2.4.4.4 Seed yield

Improvement in seed yield of different crops in response to growth substances was observed by Gopalkrishnan and Srinivasan (1975), Ries *et al.* (1977) and Menon and Srivastva (1984). The growth regulators have been found to engage in assimilate translocation towards reproductive parts of plants

(Pando and Srivastava, 1985; Khan *et al.*, 1996). Differential responses of ethephon were observed and several investigators reported about the beneficial effect on grain yields of winter wheat (Dahnous *et al.*, 1982; Leary and Oplinger, 1983; Wiersma *et al.*, 1986), while others claimed reduction with the use of ethephon (Nafziger *et al.*, 1986; Simmons *et al.*, 1988). Increase in seed yield of mustard in response to ethrel has been reported by Grewal *et al.* (1993), Khan (1996b; 1998), Singh and Kumar (1991) and Khan *et al.* (2000). Joshi *et al.* (1987) also reported increase in seed yield in groundnut (*Arachis hypogea* L.) when treated with ethrel. Foliar application of ethrel on cauliflower increased seed yield per plant at 300 mg/L while it was adversely affected at higher concentration (Jana and Kabir, 1991).

Ethephon treatment increased grain yield in corn (Kasele *et al.*, 1995) and barley (Bulman and Smith, 1993b). Contrarily, Slife and Earley (1970) applied ethrel to flowering soybean plants at 0.56 to 2.24 kg/ha, the treatments decreased yield of seed per hectare. Grewal *et al.* (1993) also observed substantial decrease in seed yield in *Brassica napus* with the use of 1000 and 1500 mg/L ethrel. Application of ethrel to foliage enhanced seed yield in *Brassica juncea* was reported by Khan *et al.* (2000). Early application of ethephon resulted in significant reduction in seed yield of linseed (Leitch and Kuat, 1999).

#### **2.4.4.5 Biological yield**

Biological yield was enhanced in ethylene treated mustard plants (Khan, 1996b; 1998; Khan *et al.*, 2000).

#### **2.4.4.6 Harvest index**

Seeds of groundnut (*Arachis hypogea* L.) were when treated with ethrel gave higher harvest index than water-soaked seeds (Joshi *et al.*, 1987), while Van Sanford *et al.* (1989) observed equivalent harvest index in both ethephon treated and control plants in winter wheat.

#### **2.4.4.7 Oil yield**

Khan (1996b) reported impressive increase in the oil yield of mustard in response to ethrel application.

#### **2.4.5 Ethylene sources and quality parameters**

##### **2.4.5.1 Oil content**

Exogenous application of ethrel at the rate of 250 mg/L reduced the essential oil content of peppermint and slightly increased the essential oil content of sage (*Sahia officinalis*) (El-Keltawi and Rodney, 1986). Farooqi and Sharma (1988) also reported an increase in oil content of *Rosa damascena* Mill by the application of 0.02 and 0.06% concentrations of ethrel. However, Khan (1996b) showed an increase in oil content in seeds of mustard. Grewal *et al.* (1993) in *B. napus* and Leitch and Kuat (1999) in linseed reported that ethrel application had no influence on the seed oil content of the plants.

##### **2.4.5.2 Amino acid content**

Sharma *et al.* (1982) reported that free amino acids contents increased during the pod development stage on application of ethrel on groundnut (*Arachis hypogea*).

##### **2.4.5.3 Protein content**

Plant growth regulators have been found to influence significantly the protein content in crop plants (Lurie *et al.*, 1994; Liu *et al.*, 1993; Yang *et al.*, 1994; Kulkarni *et al.*, 1995). They have also been implicated in the control of protein allocation among plant organs and accumulation in developing cereal grains (Oritant and Yoshida, 1971).

Grain protein concentration increased (Morris *et al.*, 1989; Van Sanford *et al.*, 1989) or remained unaffected (Pearson *et al.*, 1989) by ethephon treatment. Soluble proteins increased with application of ethrel during initial stages of pod development but declined later in groundnut (*Arachis hypogea*) (Sharma *et al.*, 1982). Ethephon treatment increased protein content per grain and grain protein concentration in barley (Bulman and Smith, 1993b; Ma *et al.*,

1994b). Ethrel treatment has been found to increase protein content and efficient incorporation of amino acids into proteins in wheat (Sekhon and Singh, 1994).

#### **2.4.5.4 Fatty acid**

Ethephon has been reported to change the relative proportions of fatty acids, reducing the content of linolenic acid and increasing the oleic acid content (Leitch and Kuat, 1999).

Ethrel has also showed a promotive effect on conversion of lipid into sugars through glyoxylate cycle under water stress conditions in soybean seeds (Sharma *et al* , 1986).

### **2.5 Mineral Nutrition**

Mineral nutrition includes the supply, absorption, utilization of essential nutrients for growth and yield of crop plants. This is not known with certainty when humans first incorporated organic substances, manures or wood ashes as fertilizers in the soil to stimulate plant growth. However, it is documented that as early as 2500 B.C., humans recognized the richness and fertility of alluvial soils in valleys of the Tigris and Euphrates rivers (Hewitt, 1963; Tisdale *et al.*, 1985).

Early progress in the development of understanding of soil fertility and nutrition concept was slow, although the Greeks and Romans made significant contributions in the years 800 to 200 B.C. (Marschner, 1986; Westerman and Tucker, 1987). It was only to the credit of Justus Von Leibig (1803-1873) that the scattered information concerning the importance of mineral nutrients for plant growth was collected and summarized the mineral nutrition of plants and was established as a scientific discipline (Marschner, 1983). Plants contain small amounts of about more than hundred elements but only 17 elements are known to be essential (Epstein, 1972; Fageria, 1984; Salisbury and Ross, 1994). The exclusive requirements of inorganic nutrients in higher plants basically distinguish them from man, animals and a number of microorganisms,

which additionally need organic compounds. Keeping in view the specific nature of the problem of thesis, the following pages deal with nutrient nitrogen, its properties and effect on oil crops with special emphasis on mustard.

### **2.5.1 Nitrogen**

Nitrogen is one of the essential elements required for crop growth in higher concentrations than any other element. Nitrogen is the most important element for plants growth and productivity During seed germination hydrolysis products of storage proteins provide the nitrogen for initial growth. After the seedlings become photosynthetic, roots take up nitrogen from a soil, which is assimilated into amino acids and proteins in the vegetative tissues. The need for nitrogen increases steadily with multiplication and expansion of the developing vegetative organs and culminates during the reproductive stage with the synthesis of new seed storage proteins (Pate, 1980).

#### **2.5.1.1 Acquisition and assimilation**

Plant acquires nitrogen from two principal resources (1) the soil through commercial fertilizers, manure and/or mineralization of indigenous organic matter and (2) the atmosphere through symbiotic  $N_2$  fixation (Vance, 1990). The availability of nitrogen in the rhizosphere regulates crop growth and development through optimum source and sink development during ontogeny. The uptake of N into plant roots is active energy requiring process (Oak and Hirel, 1985) making the roots a barrier to N uptake. The main sources of inorganic nitrogen for plant metabolism are nitrate and ammonium from the soil solution which act as substrates for reduction and subsequent incorporation into organic substances utilized by plant cell (Kleinhoffs and Warner, 1990; Lea *et al.*, 1990; Solomonson and Barber, 1990). Since ammonia is rapidly nitrified in soil (Schmidt, 1982), nitrate is a dominant form of mineral N available to higher plants except under acidic, cold and anaerobic soil conditions. The uptake and transport differ for nitrate and ammonium. The nitrate ions are transported to leaves largely as nitrates whereas ammonium

ions are assimilated in roots and transported as amino acids and amides to leaves (Arnozis and Findenegg, 1986; Shelp, 1987). Since  $\text{NH}_4^+$  is toxic, it must be rapidly assimilated into non-toxic metabolites (Lea *et al.*, 1990). The type and concentrations of nitrogen in growth media exerts a considerable influence on growth and mineral composition of crop plants (Kurvitis and Kirkby, 1980; Gashaw and Mugwira, 1981; Ansari, 1990; Jeschke *et al.*, 1992). The form of N ion supplied to roots also influences the absorption and transport of other nutrients including cations  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Na}^+$  and anions  $\text{SO}_4^{2-}$ ,  $\text{PO}_4^{3-}$  and  $\text{Cl}^-$  (Pilbean and Kirkby, 1992). As nitrate reduction is a process involving high-energy consumption, the direct utilization of ammonium could bring about a higher plant growth. However, more often the reverse is observed (Salsac *et al.*, 1987). Ammonium ions on entering the roots of plants are mediated by the GS-GOGAT system to form Gln and glutamate (Amanico and Santosh, 1992). The assimilation of  $\text{NO}_3^-$  by plants require the uptake of  $\text{NO}_3^-$ , reduction of  $\text{NO}_3^-$ , conservation of  $\text{NO}_3^-$  to  $\text{NH}_4^+$ , and the incorporation of  $\text{NH}_4^+$  to organic compounds (Migge and Becker, 1996; Sivasankar and Oak, 1996).

Subsequently nitrogenous organic compounds such as amino acids are utilized through primary metabolic pathways for various aspects of the life cycle of plants (Richards *et al.*, 1994).

#### 2.5.1.2 Nitrogen source

On a global scale terrestrial plants assimilate about 1.4 gigatons of nitrogen annually, approximately 90 to 95% of the total is from mineral nitrogen and the remaining from symbiotic dinitrogen fixation (Paul and Clark, 1988). Since dinitrogen fixation by living organisms (except for some symbiotic  $\text{N}_2$  fixers) is generally quite limited (Tisdale and Nelson, 1967), therefore nitrogen is added to soils for most cropping system to achieve high yields. The two major nitrogen fertilizers in current use are ammonium

sulphate and urea with latter becoming more common specially in the regions of high rainfall and on heavy clay soils (Shand, 1996).

### 2.5.1.3 Mineralization

It is generally assumed that N fertility in soil in natural agro-ecosystems reaches at maximum equilibrium level under specific environmental conditions where organic matter inputs equal losses (Tate, 1992). However, the intervention of man in agro-ecosystem changed this equilibrium through effects of cultivation, which accelerates the oxidation of soil organic matter at or near the surface (Campbell and Souster, 1982). This results in loss of potentially mineralizable N fraction of organic matter that serves as a reservoir of N available to crops. Mineralization and immobilization processes occur simultaneously in wetland soils and depend on soil properties and environmental factors. Mineralization of organic N (native or added organic N sources) is the most important process in N nutrition of wetland crops (Broadbent, 1978). The mineralization of organic nitrogen compounds takes place in essentially three step-by-step reactions: aminization, ammonification and nitrification.

Aminization involves the breakdown of proteins in neutral and alkaline environments by bacteria, fungi and actinomycetes. One of the final stages in the decomposition of nitrogenous materials is the hydrolytic decomposition of proteins and release of amines and amino acids. This step is termed as aminization. The amines and amino acids so released are further utilized by the still other groups of heterotrophs including both aerobic and anaerobic microorganisms with the release of ammonical compounds. This step is termed ammonification.

Nitrification is generally referred to as biological oxidation of ammonium to nitrate via nitrite, affected by *Nitrosomanas* and *Nitrobacter* species of nitrifying bacteria (Sahrawat, 1978; 1980a, b, 1989; 1996). The nitrification process is generally controlled by ammonium availability



(Robertson, 1982). Additionally, soil pH and temperature regulate the rates of ammonification and nitrification (Francis, 1982; Schmidt, 1982; Gilmour, 1984). When nitrification inhibitors are used for controlling nitrification, the process is not arrested completely resulting in a preponderance of ammonium over nitrate in the soil which affects the persistence of applied N in the soil as well as plant N metabolism and N nutrition (Sahrawat and Keeney, 1984). Nitrification inhibitors also affect nitrogen transformation other than nitrification in soils, such as ammonium fixation and release, mineralization and immobilization, nitrous oxide production and ammonia volatilisation, which affect N persistence in the soil and its subsequent availability to plants (Sahrawat, 1989; 1996).

#### **2.5.1.4 Nitrogen immobilization**

Immobilization of nitrogen is the reverse of mineralization and occurs when large quantities of low-nitrogen crops residue such as cereal crop straw begin decomposing in soil. The high amounts of carbohydrate in such residues cause the population of soil microflora to build up quickly. As new cells are formed, nitrogen and other essential elements are used to build up protoplasm. This invariably leads to decrease in the level of inorganic nitrogen for crops.

#### **2.5.1.5 Reasons for nitrogen deficiency**

The utilization efficiency of soil applied N rarely exceeds 50 per cent in most cultivated crops (Greenwood, 1982). The remainder may be immobilized in soil as organic matter or it may be lost from soil causing potentially economic and environmental consequences. Leaching, surface runoff, ammonia volatilisation and denitrification comprise the major ways for N-loss, the actual relevance of these depends on the climatic and soil conditions (Vlek and Byrnes, 1986).

Nitrogen leaching takes place mainly from nitrate but also from urea molecules. As nitrate is not absorbed to the soil complex, it tends to move downward at the same rate as water. The amount of nitrate leaching depends on

rainfall rate and soil conditions (especially texture and organic matter content). Except in few situations, the loss of N through leaching has been found non-significant (Katyal *et al.*, 1985). In general, the risk of surface transport of N is directly proportional to the amount of N applied as fertilizer. The nitrogen losses caused by leaching due to excess of rainfall can be very high. Losses were measured upto 90 kg/ha/year and were certainly one of the expectations for the low nitrogen use efficiency in cereal crops in Mediterranean regions. Reducing the risk of leaching losses would increase nitrogen use efficiency and reduce nitrogen fertilization (Carvalho and Basch, 1996).

#### **2.5.1.6 Denitrification loss**

Denitrification is the process by which nitrogenous oxides principally  $\text{NO}_3^-$  and  $\text{NO}_2^-$  are reduced to dinitrogen gases. The process is generally considered to be carried out by facultative anaerobes – organisms that normally use oxygen for respiration but in its absence use N-oxides as electron acceptors (Groffman, 1995). There is evidence that some organisms can simultaneously respire  $\text{O}_2$  and  $\text{NO}_3^-$  in what has been termed “anaerobic denitrification” (Lloyd, 1993, Robertson and Kuenen, 1991).

Nitrification in oxidized soil zones and flood water converts ammoniacal N form by ammonification and hydrolysis of urea into  $\text{NO}_3\text{-N}$ . Thus,  $\text{NO}_3\text{-N}$  can thereafter move to reduce soil zone where it is readily denitrified to dinitrogen and nitrous oxide (Reddy and Patrick, 1986). Denitrification in waste treatments can remove excess nitrate, decrease nitrate contamination of ground waters, affects atmospheric composition through the production and consumption of  $\text{N}_2\text{O}$  and thus has impact on climate. It can produce toxic intermediates  $\text{NO}$  and  $\text{NO}_2^-$  the latter can lead to carcinogenic nitrosamines (Tiedje, 1988).

#### **2.5.1.7 Gaseous loss of nitrogen**

The increase in use of N fertilizer worldwide coupled with a trend towards more extensive use of anhydrous ammonia and urea has increased the

potential for  $\text{NH}_3$  loss from fertilizers (Sharpe and Harper, 1995). Urea has become the most used N fertilizer in the world, accounting for approximately 40% of the total synthesized N supply. Its market share is increasing because it is the least expensive form of solid N fertilizer available and its high N content (46%) offers transportation advantages (Byrnes and Freney, 1995). Although urea has cost advantage over other products, but disadvantage with it is that in the considerable loss of N occurs if the urea is not incorporated in the soil soon after application. The loss occurs through ammonia volatilisation after the urea is hydrolysed at the soil surface by reaction with the enzyme urease (Mulvaney and Bremner, 1981; Ladd and Jackson, 1982). Gaseous loss of urea fertilizer N as  $\text{NH}_3$  is of particular concern because it can exceed 50% of the applied amount (Terman, 1979). Soil factors that affect volatilisation include cation exchange capacity, soil pH and moisture. Reducing  $\text{NH}_3$  loss has been found to be associated with a high cation exchange capacity i.e. there exists a negative correlation between CEC and  $\text{NH}_3$  loss (Lyster *et al.*, 1980; Ryan *et al.*, 1981; Ferguson *et al.*, 1984; O'Toole *et al.*, 1985). Ammonia losses increase with higher soil pH because of increased dissociation of  $\text{NH}_4$  to  $\text{NH}_3$ , thus increasing the potential for volatilisation (Ryan *et al.*, 1981; Chai and Hou, 1975; More and Varade, 1978). Environmental factors, which affect  $\text{NH}_3$  volatilisation, include air, temperature, wind speed, rainfall and atmospheric  $\text{NH}_3$  concentration (Denmead *et al.*, 1982; Freney *et al.*, 1983).

#### **2.5.1.8 Control of mineral fertilizer N loss in soil**

Nitrogen losses can be controlled either agrotechnically or chemically. The goal is to conserve mineral fertilizer N to a form, which is not sensitive to leaching or volatilisation. In this respect application of required amount of fertilizer to the specific requirement of the crop is recommended. This optimum nitrogen content of a plant species is the amount sufficient to maintain maximum growth rate. Any amount more than optimum nitrogen does not increase the growth rate but increases the chances of loss. Deep placement of

urea-based fertilizers can minimize volatilisation and immobilization losses by reducing fertilizer contact with dead and decaying plant residue (Doran, 1980; Jansson and Persson, 1982). When compared with surface broadcast application, deep placement has been shown to increase fertilizer N-use efficiency to about 20% (Soper *et al.*, 1971; Towes and Soper, 1978; Tomar and Soper, 1981).

#### **2.5.1.9 Consequences of nitrogen deficiency**

Nitrogen is considered one of the major limiting nutrients in plant growth and development (Richards *et al.*, 1994). The deficiency of the essential nutrients results in a significant change in the development of various organs and the plant as a whole (Mishra *et al.*, 1985). Nitrogen deficiency not only affects cell number but also size in plant species of various crops. Nitrogen stress affected leaf size through both cell production and cell expansion. Nitrogen deficits altered source-sink relations, strengthened root growth and reduced growth of whole plant and dry matter accumulation. Nitrogen metabolism and photosynthesis were unpaired, plant nitrate content and all free amino acids and proteins decreased. Nitrogen deficiency reduced plant height, leaf area, leaf number, dry matter, harvest index and also reduced yield by reducing the number of seeds per plant and by decreasing seed weight. Chweya (1990) reported that nitrogen deficiency leads to poor vegetative growth and therefore leads to accumulation of glucosinolates in the leaves of kale plant. In general, limited N supply results in plant with a reduced shoot/root dry weight ratio (Ingestad, 1980; Kerr *et al.*, 1986). The plants with no nitrogen are usually pale and yellow in colour, which would signal to a grower to apply more N (Paparozzi *et al.*, 1994). Nitrogen used in chlorophyll synthesis aids in visual diagnosis of N deficiency by influencing the plant colour, as total chlorophyll reduces in low nitrogen supply (Longstreth and Nobel, 1980). Upon decrease in nitrogen supply the rate of photosynthesis per unit leaf area decreased whereas the photosynthetic nitrogen use efficiency increased (Van der Werf *et al.*,

1993). Uhart and Andrade (1995a, b) reported that nitrogen deficiency reduced leaf area duration, leaf area index, aerial dry matter, crop growth rate, intercepted photosynthetic active radiation and water use efficiency. Results of similar nature on N deficiency were observed by number of researchers including Novoa and Loomis (1981), Lemcoff and Loomis (1986), Mai *et al.* (1987), Sinclair and Horie (1989) and Connor *et al.* (1993) and Dodd (2001).

## **2.6 Crop Response to Nitrogen**

There is a great pressure to achieve breakthrough in human nutrition by introducing new foods into diets by developing new cultivars of staple crops that have high protein contents and at the same time produce high yields. The probability of accomplishing either goal on sustained basis is still eluding. But strategic use of fertilizers and manures may change the scenario.

### **2.6.1 Nitrogen and growth parameters**

#### **2.6.1.1 Plant height**

Nitrogen promotes growth of stem by way of better cell division and their elongation (Awasthi and Surajbhan, 1994; Pradhan *et al.*, 1994). Each successive increase in N level from 0 to 8 kg/ha significantly increased the growth characters in Indian mustard (Sharma *et al.*, 1994), while increasing nitrogen levels resulted in significant increase in plant height under rainfed conditions in Indian mustard. Application of nitrogen showed significant positive effect on mean plant height of *B. napus* cv. Canola and *B. compestris* L. cv. Rapeseed (Al-Jaloud *et al.*, 1996). At both vegetative and reproductive stages, N doses showed a linear increase in plant height of fennel (*Foeniculum vulgare* Mill.) with 90 kg/ha giving maximum height increase at both stages (Afridi *et al.*, 1983). Nitrogen enhanced vegetative growth of plants has also been reported by Chweya (1990) and Paparozzi *et al.* (1994). The height of *Brassica juncea* increased significantly with increasing N from 0 to 90 kg/ha at every growth stage (Tomar *et al.*, 1996). Khanpara *et al.* (1993) have also reported the similar results.

### 2.6.1.2 Leaf number

Nitrogen supply can be very limiting to productivity mainly through its effects on number and size of leaves and roots and on the photosynthetic rate of single leaf. Increase in N levels in rainfed cotton significantly increased the number of leaves (Perumal, 1999). Application of nitrogen at a rate of 30 kg/ha increased number of leaves of mustard (*B. juncea*) (Prasad and Shukla, 1993).

### 2.6.1.3 Leaf area

The essential role of nitrogen on leaf growth has long been recognized. Nitrogen fertilization results in increased leaf area (Pregtizer *et al.*, 1990; Ericsson *et al.*, 1992; Liu and Dickman, 1992; Heilman and Fu-Guang, 1993; 1994; Perumal, 1999). Leaf area can be viewed as the result of the rate and duration of leaf expansion. Leaf expansion rate is very responsive to nitrogen supply both under controlled and field conditions (Steer *et al.*, 1984; 1986; Connor *et al.*, 1993; Dodd, 2001). Rood and Major (1984) showed leaf growth to be particularly responsive to large amounts of N in Brassica crops. Nitrogen fertilizer application increased per unit leaf area of oil seed rape (*Brassica napus* L.) (Asare and Scarisbrick, 1995). Increased allocation of dry matter to shoot was reflected in increased rates of leaf expansion and elongation in high N-plants (Huber *et al.*, 1989). Leaf area expansion was higher in wheat cultivar Oligoculum when nitrogen was applied (Madhulety and Prakash, 1988). Certainly, it is clear that if N supply rate to the crop for new leaf growth is restricted, the amount of leaf area produced is proportionally restricted (Nova and Loomis, 1981; Lemcoff and Loomis, 1986; Prasad and Shukla, 1993).

### 2.6.1.4 Leaf area index

Increasing level of N increased the leaf area index progressively. Higher leaf area index helps the crop to put forth high production (Sarkar *et al.*, 1999).

Nitrogen application to *Brassica juncea* significantly increased LAI and was considerably higher at all stages at 100 kg N/ha than 50 kg N/ha while in absence of N, leaf area index was lesser (Grewal and Kolar, 1990). Khan and

Agarwal (1988) reported that leaf area index increased with increasing N rates in the range of 0-80 kg/ha but did not show further enhancement with 120 kg N/ha in mustard (*Brassica juncea*). Leaf area index increased with increase in nitrogen level, however there was no significant increase after 40 kg N/ha in groundnut (Patra *et al.*, 1995). Perumal (1999) also reported that leaf area index increased significantly with increasing N levels on rainfed grown cotton. Pradhan *et al.* (1994) and Dey *et al.* (1989a, b) also reported similar results for leaf area index in response to nitrogen.

#### 2.6.1.5 Dry matter

Dry matter in general increased significantly with N doses applied (Cajuste *et al.*, 1996; Polley *et al.*, 1999). Nitrogen application increased biomass production favourably from an early stage and was significantly higher with 80 kg N/ha at advance stage of growth due to vigorous growth in sunflower (*Helianthus annuus* L.) (Sarkar *et al.*, 1999). Dry matter production in groundnut increased at all growth stages with increase in nitrogen level, however beyond 40 kg/ha no significant increase took place (Patra *et al.*, 1995). Sawan *et al.* (1998) reported that seedling fresh and dry weight increased with increasing N rate (161 kg N/ha) in cotton plants. Increase in N rate, increased dry matter production under water scarce condition (Vyas *et al.*, 1995). Dry matter production has also been reported to increase with the application of nitrogen (Dubey and Khan, 1993). Khan and Agarwal (1988) reported that dry matter accumulation per plant increased with increasing N rates in the range of 0-80 kg/ha in mustard. However, it does not increase further with 120 kg N/ha. Application of 60 kg N/ha significantly increased dry matter in Indian rape (*Brassica campestris* L.) genotypes (Satyavan *et al.*, 1999).

Nitrogen produced linear increase in fresh weight and dry weight of fennel (*Foeniculum vulgare* Mill.). Nitrogen fertilized plants of linseed (*Linum usitatissimum* L.) produced appreciably more dry matter throughout the growing

season than unfertilised plants (Singh and Mishra, 1994) Raghuvanshi *et al.* (1987) and Singh and Saran (1987) reported the similar results in linseed. Dry matter of maize (Mishra and Srivastava, 1985; Huber *et al.*, 1989) wheat (Madhulety and Prakash, 1988), oilseed rape (Asare and Scarisbrick, 1995; McGrath and Zhao, 1996), poinsettia (Paparozzi *et al.*, 1994), *B. juncea* (Tomer *et al.*, 1996, Khanpara *et al.*, 1993; Samiullah *et al.*, 1990), castor (Bheemaiah *et al.*, 1998) significantly increased with increase in the rates of N application. Nitrogen progressively and significantly increased the dry matter production of rainfed as well as irrigated sesame (*Sesamum indicum* L.) (Vyas *et al.*, 1999)

#### **2.6.1.6 Crop growth rate**

Nitrogen application increased crop growth rate of castor (Bheemaiah *et al.*, 1998) Pradhan *et al.* (1994) also reported increased crop growth rate by N application. Increasing level of nitrogen from 0 to 60 kg/ha increased the crop growth rate of barley genotypes under moisture stress conditions (Awasthi and Surajbhan, 1994) Post flowering application of urea solution significantly and positively influenced crop growth rate of groundnut (Reddy *et al.*, 1991). Patra *et al.* (1995) also reported increase in crop growth rate with increase in nitrogen level, however there was no significant increase after 40 kg/ha in groundnut. The crop fertilized with 80 kg N/ha showed higher crop growth rate values over 40 kg N/ha due to potential role of N in metabolic activities and on vegetative growth of *Helianthus annuus* plants (Sarkar *et al.*, 1999). Khan and Agrawal (1988) reported that crop growth rate increased with increase in N rates in mustard (*B. juncea*) but did not further increase with 120 kg N/ha.

#### **2.6.1.7 Leaf area duration**

Leaf area duration increased with increasing N rates in the range of 0-80 kg/ha but did not increased further with 120 kg N/ha in mustard (Khan and Agarwal, 1988) Patra *et al.* (1995) also reported increase in leaf area duration with increase in nitrogen level.



#### 2.6.1.8 Relative growth rate

Nitrogen application showed progressively higher relative growth rate than respective level of N upto 45-60 days but at later stages the increase was inconsistent and even decreased in certain cases with increase in N level in sunflower (*Helianthus annuus* L.) (Sarkar *et al.*, 1999).

#### 2.6.1.9 Net assimilation rate

Glimskar and Ericsson (1999) reported that decrease in net assimilation rate at nitrogen limitation was very small but net assimilation rate was slightly positively correlated with relative growth rate while the plant nitrogen content was significantly and positively correlated with relative growth rate. Increasing level of N upto 80 kg/ha increased net assimilation rate mainly due to increase in crop growth rate and relative growth rate in sunflower (Sarkar *et al.*, 1999).

### 2.6.2 Nitrogen and photosynthetic parameters

#### 2.6.2.1 Chlorophyll

The role of nitrogen in the build up of chlorophyll molecules has long been recognized (Evans and Terrshima, 1988; Sage *et al.*, 1990; Liu and Dickman, 1992). Chlorophyll content of fresh leaves, a measure of photosynthetic activity increased with N application in *Brassica juncea* (Grewal and Kolar, 1990). An increase in N rate increased levels of total chlorophyll in Indian mustard grown under different level of stored moisture (Vyas *et al.*, 1995). Leaf chlorophyll content in mustard (*Brassica juncea*) increased upto 90 kg N/ha and was highest with three irrigations (Chouhan *et al.*, 1994). Grewal *et al.* (1993) reported that the chlorophyll content in fresh leaves of *Brassica napus* L. increased with an increase in N upto 100 kg/ha.

#### 2.6.2.2 Photosynthesis

Nitrogen has a profound role in metabolic processes resulting in increasing production of photosynthates (Dhoble, 1998). Nitrogen supply can affect plant growth and productivity by altering both leaf area and

photosynthetic capacity (Bolton and Brown, 1980; Novoa and Loomis, 1981; Wong *et al.*, 1985; Sinclair, 1990).

Photosynthesis in the leaves of soybean increased with nitrate supply (Rigaud, 1981; Chi *et al.*, 1983). The increase in the rate of photosynthesis has been noted to 43% higher than that of untreated plant (Varade *et al.*, 1995). CO<sub>2</sub> assimilation is generally higher at higher N concentrations (Sage *et al.*, 1987; Evans, 1989), as the nitrogen is involved in the enzymes of the photosynthetic carbon reduction cycle and thylakoid proteins constitute the majority of leaf nitrogen (Evans and Seemann, 1989). Leaves with high nitrogen content can utilize high photon flux densities more effectively for photosynthesis than those with low nitrogen content (Field, 1983; DeJong and Doyle, 1985). With the increase in N supply there was increase in photosynthesis due to increased leaf size in groundnut (Patra *et al.*, 1995). Increasing N from 0 to 50 and from 50 to 100 kg/ha helped the crop canopy of *Brassica juncea* to trap more radiation for photosynthesis (Grewal and Kolar, 1990). Grewal *et al.* (1993) also reported that interception of PAR by the crop canopy of *Brassica napus* improved significantly with increase in N rates from 0 to 100 kg/ha.

### 2.6.3 Nitrogen and nutrient uptake

The type and concentration of nitrogen in growth media exert a considerable influence not only on the growth and mineral composition of crop plants (Kurvitis and Kirkby, 1980; Gashaw and Mugwira, 1981; Ansari, 1990; Jeschke *et al.*, 1992) but also affect the relative uptake of cations and anions (Kirkby, 1981; Lovatt, 1986). Increasing levels of nitrogen increased the uptake of N regularly irrespective of growth stage in linseed (*Linum usitatissimum* L.) (Singh and Mishra, 1994). Reddy and Reddy (1986) have also reported the similar results. The concentration of some nutrients like phosphorus, potassium and calcium in certain organs of the oilseed rape plant has been found to be influenced by the nitrogen nutrient e.g. leaf P

concentration in oilseed rape was not influenced by N supply but P concentrations in the root, stem and axillary branches increased in response to higher N nutrition (Kullimann *et al.*, 1989). Similar observations were reported by Cornertill and Steele (1981) and Amoruwa *et al.* (1987) in field grown maize.

Applied N affected amount of leaf N content such that less N was present in leaves if no N was applied in poinsettia (*Euphorbia pulcheinma* wild) (Paparozzi *et al.*, 1994). The content of organic nitrogen in leaves of maize seedlings increased with increase in the supply of inorganic nitrogen in the nutrient medium (Mishra and Srivastava, 1985). Muchow and Sinclair (1994) also reported that canopy leaf N increased in response to increased N fertilizer treatments in maize (*Zea mays* L.).

Nitrogen application increased its uptake and the effect was more pronounced under irrigated conditions in sesame (Vyas *et al.*, 1999). Pinkerton (1991) observed that critical P concentration depended on plant age and N in oilseed rape and Indian mustard. Bulman and Smith (1993a) found that higher rates of nitrogen fertilizer increased the plant N concentration and total plant accumulation in spring barley N and P uptake increased with increase in N rate in Indian mustard (Vyas *et al.*, 1995). N, P and K uptake increased with N rate in mustard varieties (*Brassica juncea* L. Czern & Coss.) (Tripathi and Singh, 1992). Compared with unfertilized control uptake of N, P, K, Ca and Mg increased with increase in N fertilizer rate in mustard. Application of 60 kg N/ha significantly increased N uptake by seed, whereas uptake of N by stalk as well as total increase was upto 90 kg/ha of N applied to Indian rape (*B. campestris* L.) genotypes (Satyavan *et al.*, 1999). Increase in the rates of N uptake with increased nitrogen application in rapeseed mustard has also been observed by Patil and Bhargava (1987), Kumar *et al.* (1989), Dubey and Khan (1993).

## 2.6.4 Nitrogen and yield parameters

### 2.6.4.1 Pod number

Nitrogen application increased mean number of pods per plant in oilseed rape (*Brassica napus* L.) under drought conditions (Asare and Scarisbrick, 1995). The number of pods per plant in Indian mustard increased by nitrogen application significantly upto 80 kg N/ha under rainfed conditions (Thakuria and Gogoi, 1996). Increase in the number of siliqua per plant with increasing levels of N upto 60 kg/ha in Indian mustard (*Brassica juncea*) has also been reported by Rathore and Manohar (1989), Khanpara *et al.* (1993), while Tomar *et al.* (1996) and Dubey *et al.* (1993) reported the increase upto 90 kg/ha. Grewal and Kolar (1990) observed that the number of pods per plant in *B. juncea* increased significantly with increase in N from 0 to 100 kg/ha. Post flowering application of urea solution significantly and positively influenced pods per plant of groundnut (Patra *et al.*, 1995).

### 2.6.4.2 Seed number

Seeds per siliquae in *Brassica juncea* significantly increased due to each increment in level of N from 0 to 90 kg/ha (Tomar *et al.*, 1996) while Khanpara *et al.* (1993) reported the increase in seeds per siliquae with increasing level of N upto 60 kg/ha in Indian mustard.

Increase in seeds per siliquae with increase in application of N fertilizers has also been reported in *B. juncea* by Reddy and Sinha (1988), Rathore and Manohar (1989), Dubey *et al.* (1993). The number of seeds per pod of *B. juncea* increased significantly with the increase in N from 0 to 100 kg/ha (Grewal and Kolar, 1990). Thakuria and Gogoi (1996) reported that nitrogen application increased seeds/siliquae upto 80 kg N/ha under rainfed conditions in Indian mustard. Contrarily, Asare and Scarisbrick (1995) could not find any significant effect of nitrogen application on number of seeds per pod in *Brassica napus* under drought conditions.

#### 2.6.4.3 1000 seed weight

Seed weight of *B. napus* was increased by nitrogen (Asare and Scarisbrick, 1995). 1000-seed weight in Indian mustard (*Brassica juncea*) increased with increasing levels of applied N upto 60 kg/ha (Khanpara *et al.*, 1993). Similar results were also recorded by Reddy and Sinha (1988), Rathore and Manohar (1989). Patra *et al.*, (1995) reported that 1000-kernel weight significantly and positively increased by post-flowering application of urea solution to groundnut.

Tomar *et al.* (1996) reported that 1000-seed weight was significantly increased due to each increment in level of N from 0 to 90 kg/ha in *Brassica juncea*. Seed weight in *B. juncea* increased significantly with increase in N level from 0 to 50 kg/ha but decreased with the further increase of N from 50 to 100 kg/ha (Grewal and Kolar, 1990). Thakuria and Gogoi (1996) also reported significant increase in 1000-seed weight in Indian mustard. Increase in 1000-seed weight in Indian mustard under rainfed conditions with increase in nitrogen upto 60 kg N/ha has also been reported by Singh and Kumar (1991). Contrary to above results, Chaudhary *et al.* (1992) reported that 1000-seed weight was unaffected by N application. Weight per seed was not affected by N treatment applied either through soil or to the peduncle in barley (Foroutan-Pour *et al.*, 1997; Ma and Smith, 1992; Ma *et al.*, 1994a).

#### 2.6.4.4 Seed yield

For agricultural crops yield has been found increasing with increase in nitrogen fertilizer application (Bandel *et al.*, 1980; Howard and Tyler, 1989; Stecker *et al.*, 1993). Reddy (1983) noted that higher fertilizer rate increased seed yield of linseed. Application of 80 kg N/ha gave significantly higher seed yield in sunflower over no nitrogen application (Sarkar *et al.*, 1999). Increase in seed yield of sunflower has also been reported by Zaman and Chaudhari (1998), Dhoble (1998) and Singh and Bansal (1999). Awasthi and Surajbhan (1994) reported that at 60 kg N/ha, growth and yield attributes were found

maximum over other levels of nitrogen in barley. Increase in the seed yield of cotton due to applied nitrogen has been reported by Sawan *et al.* (1998) and Perumal (1999). Nitrogen had little effect on the grain yield of triticale below 90 kg N/ha, but increased with higher levels of nitrogen and declined at 180 kg N/ha (Naylor and Stephan, 1993). Increase in the seed yield of *Brassica juncea* with the increase in applied nitrogen has been reported by Reddy and Sinha (1988), Rathore and Manohar (1989), Grewal and Kolar (1990), Prasad and Shukla (1992), Khanpara *et al.* (1993), Sharma (1994), Ghosh *et al.* (1995) and Thakuria and Gogoi (1996). A similar effect on *Brassica napus* has been reported by Gendy and Marquard (1989), Grewal *et al.* (1993) and Asare Scarisbrick (1995). Application of nitrogen showed significant positive effect on mean seed yield of *Brassica napus* cv. Canola and *B. campestris* L. cv. rapessed (Al-Jaloud *et al.*, 1996). Bheemaiah *et al.* (1998) also reported increase in seed yield of castor with increased nitrogen application.

#### 2.6.4.5 Harvest index

Application of 40 kg N/ha significantly increased harvest index in mustard but was reduced at 60 kg N/ha (Ghosh *et al.*, 1995). Dey *et al.* (1989b) also reported that harvest index increased upto 60 kg N/ha in rice (*Oryza sativa*).

#### 2.6.4.6 Oil yield

A linear increase in oil yield was observed with increasing N application from 0 to 100 kg N/ha (Arthamwar *et al.*, 1996). Application of 40 kg N/ha increased seed yield significantly but was reduced at 60 kg N/ha in rainfed rapeseed (Ghosh *et al.*, 1995). With the application of nitrogen, oil yield increased in different *Brassica* genotypes (Kumar *et al.*, 1995). Singh *et al.* (1994) reported that oil yield of *Brassica* species increased upto 80 kg N/ha. Rana *et al.* (1991) reported that the oil yield in *Brassica juncea* increased with irrigation and N rate. Increase in N rates in the range of 90-180 kg/ha increased

oil yields compared with lower N rates or no N in mustard (Rathore and Manohar, 1989)

## 2.7.5 Nitrogen and quality parameters

### 2.7.5.1 Oil content

Smith *et al* (1988) described an inverse linear response between oil content and seed N concentration in *Brassica napus*. Nitrogen application affected adversely the seed oil content of oilseed rape (*B. napus* L.) (Asare and Scarisbrick, 1995; Pinkerton, 1991, Gendy and Marquard, 1989) while Grewal *et al* (1993) reported that oil content of seeds in *B. napus* significantly increased with the application of 50 kg N/ha over no nitrogen application. However, increasing the dose of N from 50 to 100 kg/ha resulted in a decline in the oil content of seeds. Oil content of rainfed mustard increased significantly upto 40 kg N/ha and there was reduction in oil content at 60 kg N/ha (Ghosh *et al*, 1995). Oil percentage of *Brassica juncea* seeds increased significantly with the application of 50 kg N/ha but application of 100 kg N/ha resulted in significant decline (Grewal and Kolar, 1990). Increase in the oil content with N application has also been reported by Ghatak *et al*. (1992), Arora *et al*. (1994), Chouhan *et al* (1994) and Patra *et al* (1995). Application of nitrogen showed significant positive effect on mean oil content of *Brassica napus* L. cv Canola and *Brassica campestris* L. cv. Rapeseed (Al-Jaloud *et al*, 1996). However, Satyavan *et al* (1999) reported that N application at all rates decreased oil content of Indian rape (*B. campestris* L.) genotypes when compared with control. The adverse effect of nitrogen on oil content in different Brassica genotypes has been reported by Rana *et al*. (1991), Tripathi and Singh (1992), Dubey and Khan (1993), Singh *et al*. (1994), Shukla and Kumar (1994), Kumar *et al*. (1995) and Tomar *et al*. (1997). While Rathore and Manohar (1989), Saran and Giri (1990) and Chaudhary *et al*. (1992) reported that seed oil content in *Brassica juncea* was unaffected by the applied nitrogen rate.

### 2.7.5.2 Aminoacids

As nitrogen application to triticale increased over the range of 0-80 kg/ha there was a significant decrease in the proportions of alanine and glycine. However, significant increase in histidine and phenylalanine was observed. Serine showed no significant response to the supply of nitrogen. Other amino acids generally showed a curvilinear response to proportion of nitrogen in the grain protein (Naylor and Stephen, 1993). An increase in amino acids with a simultaneous decrease in carbohydrate synthesis has been reported when ammonium (Platt *et al.*, 1977) or nitrate (VanQuay *et al.*, 1991; Champigny *et al.*, 1992) was supplied. Vyas *et al.* (1995) reported that as N rate increased, the free amino acids increased in Indian mustard despite low water potential.

### 2.7.5.3 Protein content

Under field conditions grain protein concentration increased as soil nitrogen fertility increases in barley (Bulman and Smith, 1993b). Increasing the amount of nitrogen applied usually caused an increase in the deposition of storage proteins (which has higher contents of glutamic acid and proline) and thereby reducing the proportion of proteins typical of the embryo and aleurone layer (e.g. aspartic acid, arginine and lysine) in triticale (Naylor and Stephen, 1993). Patra *et al.* (1995) reported that with the increase in N supply, the quantity of soluble proteins increased in groundnut. Protein content of leaves in maize seedlings increased with the increase in the supply of inorganic nitrogen in the nutrient medium (Mishra and Srivastava, 1985). Nitrogen levels significantly affected the protein content with each increase in the dose of N upto 90 kg/ha in Indian mustard (*Brassica juncea*). As the N supply increased the formation of protein also increased (Dubey *et al.*, 1994). Tandon (1986) and Singh and Saran (1987) also found similar results. Asare and Scarisbrick (1995) also reported inverse relationship between applied nitrogen and seed protein of *Brassica napus* L. seeds. However, Gendy and Marquard (1989) reported that protein content of *Brassica napus* L. seeds increased with increase



in nitrogen supply. Application of nitrogen showed significant positive effect on mean protein content of *Brassica napus* L. cv. Canola and *Brassica campestris* L. cv. Rapeseed under irrigated conditions (Al-Jaloud *et al.*, 1996). Increase in the protein content in rapeseed mustard with increasing rate of nitrogen applied has also been reported by Patil and Bhargava (1987) and Satyavan *et al.* (1999).

#### 2.7.5.4 Fatty acids and glucosinolate

Seed glucosinolate content responded variably to use of N fertilizer application. It increased significantly with N application in some cultivars but not in other cultivars of *Brassica napus* L. (Asare and Scarisbrick, 1995).

In contrast, Ramans (1989) reported no significant effect of nitrogen fertilizer application on glucosinolate content in *Brassica napus*. The allyl-isothiocyanate content increased in the oil of Indian mustard (*Brassica juncea* L. Czern & Coss.) with higher N rates (Narang *et al.*, 1985). Seed allyl-isothiocyanate content was increased by N application in mustard (Arora *et al.*, 1994). Thakral *et al.* (1996) reported that glucosinolate contents of *Brassica carinata* and *Brassica napus* increased with increase in N fertilizer. While Singh *et al.* (1994) reported that the sinigrin glucosinolate content of *Brassica* species increased upto 20 kg N/ha.

Fatty acid composition in seeds of *Brassica* species was affected significantly with increase in levels of N fertility (Thakral *et al.*, 1995). Allyl-isothiocyanate and fatty acid contents were increased by applied N (0-90 kg N/ha) in *Brassica juncea*. Application of nitrogen increased the erucic acid content but decreased linolenic acid linoleic acid contents in *Brassica juncea*. Application of N on white mustard increased percentage of oleic acid but decreased linolenic acid in the oil. Total saturated acids increased with N. Contrarily, Gendy and Marquard (1989) reported that the pattern of fatty acids in the oil of *Brassica napus* L. was not influenced by nitrogen level.

#### 2.7.5.5 Acidity of oil/iodine number

The iodine value of *Brassica* species oil increased upto 20 kg N/ha (Singh *et al.*, 1994). Arora *et al.* (1994) also reported that iodine value of mustard oil was increased by N application. Iodine value of *Brassica carinata* and *Brassica napus* oil increased with increase in N (Thakral *et al.*, 1996). Application of nitrogen at a rate of 0-90 kg/ha increased iodine value of *Brassica juncea* oil. Narang *et al.* (1985) also reported that iodine number of *Brassica juncea* oil increased with higher N rates.

### 2.8 Crop Response to Interaction between Ethylene Sources and Nitrogen

Plant hormones and inorganic nutrients share a common physiological function i.e. both of these growth factors influence the growth and development of plants. Plant growth regulators have been found to be engaged in enhancing growth and productivity of crop plants. There are many instances which suggest that growth regulators and nutrients can interact in a variety of ways. Deficient and toxic levels of nutrients can affect the concentration of specific hormones, and in turn, hormones have the capacity to direct the translocation and accumulation of nutrients in plants (Kuiper, 1988; Kuiper *et al.*, 1989). Actually the nutrient status of a plant influences its metabolism and growth and can affect the synthesis and distribution of growth substances (Haru *et al.*, 1982; Green, 1983). Considering the complex interactions of plant hormones and multiplicity of plant functions they control, the impact on nutrients on hormones is an important issue (Whenham *et al.*, 1989; Thoresteinsen and Eliasson, 1990; Arshad and Frankenberger, 1991; Cao *et al.*, 1993).

In the following pages, attempt has been made to cover aspects of interactions of nutrients with ethylene sources application.

#### 2.8.1 Interaction effect on growth parameters

Growth of plant was found to be influenced by the interaction of hormones and nutrients. Retardation in the plant height of gobhi sarson (*Brassica napus* L.), with the foliar application of ethrel at all the levels of

nitrogen was noted and the reduction was more pronounced in the absence of nitrogen (Grewal *et al.*, 1993). Flower number of different crops was found to be positively responsive towards exogenous application of growth substances (Morgan *et al.*, 1983; Friends, 1985). Grewal *et al.* (1993) noted that in a field grown *Brassica napus* L. foliar spray of 500  $\mu\text{L/L}$  ethrel with basal 50 kg N/ha, resulted in more LAI, whereas at 100 kg N/ha, 1000  $\mu\text{L/L}$  ethrel proved to be more beneficial. The findings are in close conformity with Bengal *et al.* (1982) and Singh *et al.* (1987). Ethrel spray (400 or 600  $\mu\text{L/L}$ ) along with basal (45 or 60 kg/ha) and foliar application of (0 or 10 kg/ha) of N, increased leaf area index and dry mass of mustard (Khan, 1996b). In a study on mustard, Khan *et al.* (2000) reported that at 0 or 40 kg N/ha, ethrel did not produce any significant effect, but at basal 80 kg N/ha, ethrel affected growth parameters like LAI and total dry matter.

## **2.8.2 Interaction effect on photosynthetic parameters**

### **2.8.2.1 Chlorophyll content**

Ethrel at 500  $\mu\text{L/L}$  significantly improved the chlorophyll content in leaves of *Brassica napus* when 50 and 100 kg N/ha was applied. However, under no nitrogen application, higher dose of ethrel (1000 and 1500  $\mu\text{L/L}$ ) showed detrimental effects (Grewal *et al.*, 1993).

Grewal and Kolar (1990) reported that application of 500, 1000 and 1500  $\mu\text{L/L}$  of ethrel reduced chlorophyll content when no N was applied but the reduction was only significant for 1500  $\mu\text{L/L}$  ethrel at 0 and 50 kg N/ha. However, 500 and 1000  $\mu\text{L/L}$  of ethrel at 100 kg N/ha significantly increased the chlorophyll content of leaves in *Brassica juncea* compared with water sprayed plants.

### **2.8.2.2 Photosynthesis**

Grewal *et al.* (1993) observed indirect evidence of photosynthetic activities by enhancing chlorophyll content and retaining higher LAI with nitrogen (50 and 100 kg N/ha) and spray of ethrel (500  $\mu\text{L/L}$ ) in *Brassica*

*napus*. Khan *et al.* (2000) also reported that at basal 80 kg N/ha, ethrel spray improved LAI, thus resulting in more solar radiation being retained and enhanced net photosynthetic rate in Indian mustard (*Brassica juncea* L.)

#### **2.8.2.3 Photosynthetic active radiation**

Since the growth and crop productivity of crop species are governed to a great extent by its surrounding environment, hence any change in the quality of solar radiation would certainly influence the growth and productivity of several crop species in various agro-climatic zones (Parry, 1992; Sinha, 1992; Singh, 1997). The rate of photosynthesis and photosynthetically active radiation are highly influenced with leaf development and canopy structure (Ramanujan and Mohan Naidu, 1995).

In absence of N, ethrel at 1000 and 1500  $\mu\text{L/L}$  significantly reduced the interception of PAR, while increase in nitrogen application (50 and 100 kg N/ha) resulted in improvement in interception of PAR (Grewal *et al.*, 1993). Similarly, in another study, Grewal and Kolar (1990) observed that increase in N (from 0 to 50 and 50 to 100 kg/ha) helped the crop canopy to trap more radiation. Ethrel at 500  $\mu\text{L/L}$  concentration improved the radiation interception at 50 and 100 kg N/ha. At basal 60 kg N/ha and foliar 10 kg N/ha ethrel improved photosynthetic activities, improving crop canopy and retaining higher LAI during development phase of *Brassica juncea* (Khan, 1996b).

#### **2.8.3 Interaction effect on nutrient uptake**

Salama and Bazas (1987) reported the significant interaction effect between growth regulators and N, P and K fertilizers on the copper content of sunflower on calcareous sandy loam soils. Erdei and Dhakal (1980) reported that ethrel stimulated K uptake in wheat. In *Brassica juncea* increase in N uptake with ethrel and N application has been reported by Khan *et al.* (2000).

#### **2.8.4 Interaction effect on yield parameters**

Enhanced seed yield in *Brassica napus* in response to N application was more pronounced with spray of ethrel. Application of 50 kg N/ha and spray of

ethrel (500  $\mu\text{L/L}$ ) significantly improved the grain yield and further improvement was observed with 100 kg N/ha and 1000  $\mu\text{L/L}$  ethrel (Grewal and Kolar, 1993). Khan (1996b) reported that ethrel in association with nitrogen significantly increased pods per plant, seed per pod, 1000 seed weight, seed yield, oil content and oil yield in mustard. However, ethrel proved less effective at lower nitrogen dose because of insufficient availability of photosynthates. The number of pods per plant and number of seeds per pod increased sufficiently with 500 and 1000  $\mu\text{L/L}$  ethrel only at 50 and 100 kg N/ha in *Brassica juncea*. At 50 kg N/ha, ethrel at 500  $\mu\text{L/L}$  significantly improved seed yield and resulted in an increase in seed weight (Grewal and Kolar, 1990). Khan *et al.* (2000) reported that ethrel enhanced pods per plant, seed yield and seed yield merit of Indian mustard. However, the response of mustard was greater with the application of 80 kg N/ha than 0 and 40 kg N/ha.

## 2.9 Concluding Remarks

The foregoing review clearly established the fact that plant growth regulators are potent chemicals for enhancing performance of many crops. They exhibit pronounced interaction effect with nutrients. However, only few studies regarding the interaction effect of ethylene sources with nitrogen in mustard has been conducted. These studies do not provide in depth insight on the role of ethylene sources on nitrogen metabolism and involvement of ethylene sources on physiological aspects of plant growth and development. The work on the interaction effects of ethylene sources like, ethrel with nitrogen seems to have been neglected particularly in mustard. Therefore, an in depth study on the physiological response of mustard to ethrel and nitrogen is highly desirable. The research work reported in the subsequent chapter is related to this aspect in order to fill the existing lacuna in the studies on ethrel with nitrogen interaction for better understanding of role played by ethrel in determining growth and productivity of mustard grown with nitrogen doses.

***CHAPTER-3***  
***MATERIALS AND***  
***METHODS***

# CONTENTS

## MATERIAL AND METHODS

3 1	Experimental Material	47
3.1 1	Nomenclature	47
3 1 2	Botanical description	47
3.2	Experimental Site	48
3 3	Agro-Climatic Conditions	48
3 3 1	Topography	48
3 3 2	Climate	48
3 3 3	Temperature	48
3 3 4	Rainfall	49
3 4	Meteorological Inputs	49
3 5	Soil Characteristics	49
3 6	Cultural Operations	49
3 6 1	Preparatory tillage	49
3 6.2	Sowing	50
3 6 3	Thinning	50
3 6 4	Crop protection	50
3 6.5	Irrigation	50
3.6 6	Plant sampling	50
3.7	Application of N Fertilizer	50
3 8	Application of Ethrel (2-Chloroethyl phosphonic acid)	51
3 9	Application of Silver Thiosulphate	51
3 10	Experimentation	52
3 10.1	Experiment 1	52
3 10 2	Experiment 2	52
3 10.3	Experiment 3	53
3 10 4	Experiment 4	53
3 10.5	Experiment 5	54
3 11	Biometric Observation	55
3 12	Growth Characteristics	55
3 12.1	Plant leaf area	56
3 12.2	Leaf area index	56
3.12 3	Specific leaf area	56
3 12 4	Specific leaf weight	56
3 12.5	Dry weight	57
3.12.6	Leaf fresh weight and turgid weight	57
3 12 7	Leaf relative water content	57
3 13	Physiological Characteristics	57
3.13 1	Rate of photosynthesis	58
3 13.2	Carboxylation efficiency	58

3.13.3	Photosynthetic water use efficiency	58
3.13.4	Plant water use efficiency	59
3.13.5	Estimation of 1-aminocyclopropane-1-carboxylic acid (ACC)	59
3.13.6	Assay of ACC oxidase activity	59
3.13.7	Ethylene evolution	59
3.14	Biochemical Characteristics	60
3.14.1	Assay of nitrate reductase activity	60
3.14.1.1	Colour development	61
3.14.1.2	Standard curve for NRA	61
3.14.2	Nitrogen content	61
3.14.2.1	Digestion of plant samples for N content	61
3.14.2.2	Estimation of nitrogen	62
3.14.2.2.1	Standard curve for nitrogen	62
3.14.3	N <sup>+</sup> accumulation	63
3.15	Yield Characteristics	63
3.15.1	Number of pods per plant	63
3.15.2	Number of seeds per pod	63
3.15.3	1000 seed weight	63
3.15.4	Seed yield	64
3.15.5	Biological yield	64
3.15.6	Harvest index	64
3.15.7	Seed N content per plant	64
3.15.8	Nitrogen harvest index	64
3.15.9	Nitrogen yield potential	64
3.15.10	Oil yield	65
3.16	Quality Characteristics	65
3.16.1	Determination of oil content	65
3.16.2	Determination of acid value	65
3.16.3	Determination of iodine value	66
3.16.4	Determination of saponification value	67
3.17	Statistical Analysis	67



## MATERIALS AND METHODS

A comprehensive detail of material used and methodology adopted during the course of the present investigation are presented in this chapter.

### 3.1 Experimental Material

The term rapeseed is used to cover sarson, toria and taramira, whereas the term mustard refers only to 'rai'. Seeds of mustard (*Brassica juncea* L. Czern & Coss.) of two cultivars, namely Alankar and PBM16 were used as experimental material.

Seeds of the above cultivars were obtained from the Indian Agricultural Research Institute, New Delhi, India, and are recommended equally well for cultivation under drier and wet conditions.

#### 3.1.1 Nomenclature

Indian *Oleiferous Brassicae* are divided into four groups.

1. Brown mustard, commonly called 'rai' (raya or laha) *Brassica juncea* (L.) Czern & Coss.
2. Sarson (i) Yellow Sarson – *Brassica campestris* L. var. Sarson Prain.  
(ii) Brown Sarson – *Brassica campestris* L. var. Dichotoma Watt.
3. Toria (lahi or maghi lahi) *Brassica campestris* L. var. Toria Duth
4. Taramira or tara (*Eruca sativa* Mill.)

In addition, there are two other species, namely *Brassica nigra* Koch (Banarasi rai) and *Brassica juncea* var. Rugosa (pahadi rai) which do not fall under any of the four groups, and are grown to a limited extent.

#### 3.1.2 Botanical description

Rape and mustard are annual herbs. Roots, in general, are long and tapering. Toria is more or less a surface feeder but brown mustard has long roots, with limited lateral spreads, enabling its successful cultivation under drier conditions. Yellow sarson has both extensive and lateral spreads. The

height of the stem varies from 0.45m (in some cultivars of toria) to 1.90m (in yellow sarson). In toria and brown sarson, the branches arise at an angle of 30° to 40°. In yellow sarson, the branches arise laterally at an angle of about 10° to 20° and give the plant a narrow and pyramidal shape. The inflorescence is a corymbose raceme. In the case of yellow sarson, the four petals are spread apart, whereas in brown sarson and toria, the petals overlap or may be placed apart, depending upon the cultivar. The flowers bear a hypogynous syncarpous ovary. In brown sarson and toria, the ovary is bicarpellary, whereas in case of yellow sarson, it may be tri or tetra-carpellary.

The fruit is a siliqua. The pods are two, three or four valved, depending on number of carpels in the ovary. The flowers begin to open from 8 a.m. and continue upto 12 noons.

### **3.2 Experimental Site**

Five field experiments were conducted at the Experimental field of the Aligarh Muslim University, Aligarh, India.

### **3.3 Agro-Climatic Conditions**

#### **3.3.1 Topography**

Aligarh has an area of 5,024 sq. km and is situated at 27°52'N latitude, 78°51'E longitude and 187.45m altitude above sea level.

#### **3.3.2 Climate**

It has a semi arid and subtropical weather with severest hot dry summers and intense cold winters.

#### **3.3.3 Temperature**

The winter stretches from middle of the October till the end of March. A gradual decrease in the temperature in December and January is observed, reaching as low as 15°C and 13°C, and lowest recorded for any single day is 2°C and 0.5°C respectively. The summer season extends from April to the end of June. In this season, a gradual increase in temperature is recorded, which attains its maximum, sometimes in the month of June upto 46°C (Fig. 1).




Fig. 1. Monthly temperature variation at Aligarh

Temp  
min max

### 3.3.4 Rainfall

The mean annual rainfall is about 847.3mm. More than 85% of the total downpour is derived during a short span of four months from June to September. The remaining rain drops are received during winter. These showers are very useful for winter crops. However, they are sometimes accompanied with high wind velocity and hailstorm (Figs. 2-3).

### 3.4 Meteorological Inputs

Meteorological data for the present study were documented at the Meteorological Observatory, Department of Physics, Aligarh Muslim University, Aligarh, India.

### 3.5 Soil Characteristics

Random soil samples were collected from various chosen spots, spread over the entire experimental crops upto depth of 15 cm, and analysed for physico-chemical characteristics of the soil. Data obtained on chemical characteristics and physical constant for soil are presented in Table 1. The soil was also analysed for moisture content at different growth stages of the crop (Table 1a).

### 3.6 Cultural Operations

The field experiments were laid out in randomized complete block design with three replicates for each treatment. The individual plot size was 10 sq m (2m X 5m).

#### 3.6.1 Preparatory tillage

Before each trial, diligent ploughing of field was done to turn the soil for maximum aeration and weed eradication. The plots were made with proper boundaries along with necessary irrigation channels and were irrigated lightly before sowing to maintain proper moisture in the sub-surface of the soil.




Fig. 2-3. Variation in relative humidity and rainfall during experimental period at Aligarh

Table 1 Physico-chemical characteristics of soil of the field used for experiments

	Experiment 1 (1998–1999)	Experiment 2 (1998–1999)	Experiment 3 (1999–2000)	Experiment 4 (1999–2000)	Experiment 5 (2000–2001)
Texture	Sandy loam	Sandy loam	Sandy loam	Sandy loam	Sandy loam
pH (1 2)	<u>8 0</u>	<u>8 4</u>	7 8	7 9	7 6
EC (1 2) (m mhos/cm)	0 46	<u>0 41</u>	0 45	0 43	0 45
Available nitrogen (N) (kg/ha)	205	190	210	215	215

Table 1a Moisture content (%) of the soil sample during various growth stages of experimental period

Sampling stages	Experiment 1	Experiment 2	Experiment 3	Experiment 4	Experiment 5	
	(1998–1999)	(1998–1999)	(1999–2000)	(1999–2000)	Irrigated	Non-irrigated
Sowing (0d)	12.6	12.5	12.7	12.3	12.5	12.1
Flowering (60d)	11.5	8.6	11.4	8.2	11.6	8.3
Pod fill (80d)	9.6	6.6	9.4	6.4	9.6	6.3
Pod maturity (100d)	8.2	4.2	8.3	3.9	8.0	4.0
Harvest (120d)	6.0	3.1	6.2	2.9	6.3	2.9



### 3.6.2 Sowing

The seeds were sown by the usual behind the plough method at a rate of 10 kg/ha. A distance of 30 cm between rows and 15 cm between plants in each row was maintained.

### 3.6.3 Thinning

After establishment of the crop i.e. after 12 days of sowing, seedlings were thinned to maintain the uniformity (12 plants/m<sup>2</sup>) of the plant population.

### 3.6.4 Crop protection

In order to check the aphid contagion, if any, insecticidal spray of Dimecron-100 was done. Hand weeding was done twice during the entire crop season to keep the experimental field free of weeds.

### 3.6.5 Irrigation

Before sowing 20 litre/m<sup>2</sup> irrigation was given to all plots maintained for irrigated and non-irrigated experiments. Thereafter this amount of irrigation was done only to the plots used for experiments ~~done~~ under irrigated conditions at 50d after sowing.

### 3.6.6 Plant sampling

Plants were sampled to record observations on various growth (Section 3.10), physiological (Section 3.11) and biochemical (Section 3.12) characteristics at different stages of plant growth. At maturity, plants were harvested to record yield characteristics (Section 3.13) by cutting at the ground level and were allowed for sun drying. After drying, threshing was done to clear the seeds for seed yield. Seeds were analysed for quality characteristics (Section 3.14).

## 3.7 Application of N Fertilizer

In Experiments 1 and 2, a recommended uniform soil application of 80 kg/ha was done to all plots, and in other experiments nitrogen (as urea) was applied according to the treatment and design of the experiments.



### 3.8 Application of Ethrel (2-Chloroethyl phosphonic acid)

Concentrations of ethrel used in experiments were sprayed as a.i. ( $\mu\text{L/L}$ ) on leaves at 60d after sowing (flowering stage) at 600 litre /ha (600 mL/pot) using a hand sprayer (Hindustan Sprayers, New Delhi).

In mustard, flowering stage has been found as appropriate stage for spray (Khan, 1998; Khan *et al.* 2000). Early application of PGRs is not recommended because treatment effects are compensated for by prolonged duration of the crop photosynthesis and equal or increased assimilate partitioning to the seeds. This may result in non-significant difference in control and treated plants. The other reason for spray at flowering is that there exists internal hormonal imbalance during sink development and due to this, only 68 % flowers develop into pods. Therefore, spray at flowering stage brings hormonal status to suitable concentration and restricts flowers and pods abortion. Moreover, plants may be benefited additionally by spray at this stage by increasing the capacity of photoassimilate assimilation and sink strength since plants inherent capacity of these started declining.

### 3.9 Application of Silver Thiosulphate

Silver thiosulphate (STS) at 1mM concentration was sprayed at 60d after sowing (flowering stage) at 600 litre/ha (600 mL/plot). Application of ethrel was used to promote ethylene evolution, while the action of this gaseous plant growth regulator was blocked using silver ions applied as silver thiosulphate (STS). The objective was to test the hypothesis that ethylene has a central role in mediating plant responses. Silver thiosulphate solution was used to block the action of ethylene, as silver ions have been proposed to reduce the capacity of ethylene to interact with its receptors (Beyer, 1976). Silver thiosulphate is readily absorbed and transported by plants (Morgan *et al.*, 1993).

### 3.10 Experimentation

#### 3.10.1 Experiment 1

This experiment was a factorial performed according to randomized complete block design during the winter season of 1998-99. The experiment was performed to assess the effect of leaf-applied 0, 100, 200, 400 and 600  $\mu\text{L/L}$  of ethrel (2-chloroethyl phosphonic acid, 99.9 % a.i. CDH Bombay). The ethrel was applied on Alankar and PBM 16 cultivars of mustard (*Brassica juncea* L.) at 60 d after sowing (flowering stage). Alankar is a well adapted cultivar grown in the region, whereas PBM16 is a newly released cultivar. Ethrel was sprayed at a rate of 600 litre/ha (600 mL/plot) together with 0.5% teepol (a surfactant). In control group of plants, equal amount of de-ionized water with 0.5% teepol was sprayed.

The size of each plot was  $10\text{m}^2$  (2m X 5m). The seeds were sown by the usual behind the plough method at a rate of 10 kg/ha. Each treatment was replicated thrice. Irrigation was done once during the entire season of the crop. Spray of an insecticide (Dimecron-100) was done to check aphid contagion, if any.

The scheme of the treatments is summarized in the Table 2 and ANOVA is given in the Table 2a.

At 20 days interval i.e. 80 (pod fill), 100 (pod maturity) and 120 d (harvest) after sowing, five plants from each plot were taken out with the help of hand hoe and various growth characteristics (Section 3.10) and biochemical characteristics (Section 3.12) were determined.

Physiological characteristics (Section 3.11) were studied at 80 and 100 d after sowing. At harvest (120 d after sowing), yield characteristics (Section 3.13) and quality characteristics (Section 3.14) were recorded.

#### 3.10.2 Experiment 2

This experiment was also a factorial conducted simultaneously with Experiment 1 according to randomized complete block design, but the

Table 2. Scheme of treatments for Experiment 1 (1998–99)

Cultivars	Ethrel ( $\mu\text{L/L}$ )				
	0	100	200	400	600
Alankar	+	+	+	+	+
PBM16	+	+	+	+	+
Crop	Mustard ( <i>Brassica juncea</i> L. Czern & Coss.)				
	grown under irrigated conditions				
Treatment	Spray at 60d after sowing (flowering stage) <i>and 20 days</i> <i>interval thereafter</i>				
Design	Randomized complete block design				

Table 2a. Model of analysis of variance (ANOVA) of data from a 5x2 factorial experiment in randomized complete block design (Experiment 1)

Source of variation	d.f.	S.S.	M.S.S.	F. value
Replication	2			
Treatment	9			
Spray (S)	4			
Cultivar (C)	1			
Interaction (S x C)	4			
Error	18			
Total	29			

experiment was carried out under non-irrigated conditions. The scheme of the treatments and other details of observations are same as in Experiment 1. The scheme of treatments is given in Table 3 and ANOVA in Table 3a.

### 3.10.3 Experiment 3

A factorial experiment conducted according to randomized complete block design during winter season of 1999-2000 under irrigated conditions. The aim of the experiment was to investigate the effect of leaf-applied 0, 100 or 200  $\mu\text{L/L}$  ethrel (selected on the basis of Experiment 1) at 60d after sowing (flowering stage) on growth, physiological, biochemical, yield and quality characteristics of mustard (*Brassica juncea* L.) cultivar Alankar. The cultivar Alankar was selected based on its better performance than PBM 16 in Experiment 1. The plants were grown with 0, 40, 60 and 80 kg N/ha. The N doses were selected on the basis of recommendation for cultivation of mustard. Basal 80 kg N/ha is recommended dose and the other doses 60 and 40 kg N/ha are two sub-optimal doses.

The scheme of the treatments is summarized in the Table 4 and ANOVA is given in Table 4a. All other plant growing cultivation practices, including size of plots, sowing method, seed rate, number of irrigation, weeding and pest control operations were kept same as in Experiment 1.

Sampling was done at 80, 100 and 120 d after sowing to record growth, physiological, biochemical, yield and quality characteristics as described for Experiment 1. Among biochemical characteristics, nitrate reductase activity in leaves was also determined in this experiment.

### 3.10.4 Experiment 4

This experiment was carried out simultaneously with Experiment 3, but under non-irrigated conditions in the winter season of 1999-2000. The aim of this factorial randomized complete block design experiment was to study the effect of leaf-applied 0, 100 or 200  $\mu\text{L/L}$  ethrel at 60 d after sowing (flowering stage) on mustard (*Brassica juncea* L.) cultivar Alankar grown with basal 0,

Table 3. Scheme of treatments for Experiment 2 (1998–99)

Cultivars	Ethrel ( $\mu\text{L/L}$ )				
	0	100	200	400	600
Alankar	+	+	+	+	+
PBM16	+	+	+	+	+
Crop	Mustard ( <i>Brassica juncea</i> L. Czern & Coss.)				
	grown under non-irrigated conditions				
Treatment	Spray at 60d after sowing (flowering stage)				
Design	Randomized complete block design				

Table 3a. Model of analysis of variance (ANOVA) of data from a 5x2 factorial experiment in randomized complete block design (Experiment 2)

Source of variation	d.f.	S.S.	M.S.S.	F. value
Replication	2			
Treatment	9			
Spray (S)	4			
Cultivar (C)	1			
Interaction (S x C)	4			
Error	18			
Total	29			

Table 4 Scheme of treatments for Experiment 3 (1999–2000)

Basal nitrogen levels (kg N/ha)	Ethrel (μL/L)		
	0	100	200
N <sub>0</sub>	+	+	+
N <sub>40</sub>	+	+	+
N <sub>60</sub>	+	+	+
N <sub>80</sub>	+	+	+
Crop	Mustard ( <i>Brassica juncea</i> L Czern & Coss )		
	grown under irrigated conditions		
Treatment	f, ct Spray at 60d after sowing (flowering stage)		
Design	Randomized complete block design		

Table 4a Model of analysis of variance (ANOVA) of data from a 3x4 factorial experiment in randomized complete block design (Experiment 3)

Source of variation	d f	S S	M S S	F value
Replication	2			
Treatment	11			
Spray (S)	2			
Nitrogen (N)	3			
Interaction (S x N)	6			
Error	11			
Total	35			

40, 60 and 80 kg N/ha on the plant characteristics described for Experiment 3. All practices of cultivation and observation details recorded at various growth stages are similar to Experiment 3. The scheme of treatments is summarized in the Table 5 and ANOVA is given in Table 5a.

Experiment 1 and 2 were conducted separately in one season in view of easy handling of the experiments, but for comparison of effect of ethrel spray treatments under irrigated and non-irrigated conditions analysis of some important data as group comparison was done. Similar analysis was carried out for Experiment 3 and 4.

### 3.10.5 Experiment 5

This factorial experiment was performed according to randomized complete block design during winter season of 2000-2001. Based on the findings of Experiment 3 and 4, application of 0 and 200  $\mu\text{L/L}$  ethrel or 1mM silver thiosulphate (STS) on mustard (*Brassica juncea* L.) cultivar Alankar grown under irrigated and non-irrigated conditions was done at 60 d after sowing (flowering stage). A uniform basal application of 80 kg  $\text{N/ha}$  was given. The aim of this experiment was to confirm the findings on effects of ethrel on plant characteristics as found in Experiment 3 and 4. Because of this reason silver thiosulphate, which inhibits physiological action of ethylene was used in the experiment along with ethrel. Ethrel is a source of ethylene and its effects are manifested through physiological action of ethylene. The scheme of the treatment is summarized in the Table 6 and ANOVA is given in Table 6a.

All other plant growing cultivation practices, including size of plots, sowing methods, seed rate, number of irrigation, weeding and pest control operations were kept same as in Experiment 3 and 4.

Sampling was done at 80, 100, 120 d after sowing to assess the growth performance in terms of leaf area and dry weight per plant, physiological, yield and quality characteristics were studied. The details of the characteristics studied are given in the following pages.

Table 5 Scheme of treatments for Experiment 4 (1999–2000)

Basal nitrogen levels (kg N/ha)	Ethrel (μL/L)		
	0	100	200
N <sub>0</sub>	+	+	+
N <sub>40</sub>	+	+	+
N <sub>60</sub>	+	+	+
N <sub>80</sub>	+	+	+
Crop	Mustard ( <i>Brassica juncea</i> L. Czern & Coss.)		
	grown under non-irrigated conditions		
Treatment	Spray at 60d after sowing (flowering stage)		
Design	Randomized complete block design		

Table 5a Model of analysis of variance (ANOVA) of data from a 3x4 factorial experiment in randomized block design (Experiment 4)

Source of variation	d f	S.S	M S S	F. value
Replication	2			
Treatment	11			
Spray (S)	2			
Nitrogen (N)	3			
Interaction (S x N)	6			
Error	11			
Total	35			



Table 6. Scheme of treatments for Experiment 5 (2000–2001)

Irrigation	Spray treatment		
	0 Water	200 $\mu$ L/L Ethrel	1mM Silverthiosulphate
Irrigated	+	+	+
Non-irrigated	+	+	+
Crop :	Mustard ( <i>Brassica juncea</i> L. Czern & Coss.) cv. Alankar grown under irrigated and non-irrigated conditions		
Treatment :	Spray at 60d after sowing (flowering stage) 8 litres a/ha		
Design :	Randomized complete block design		

Table 6a. Model of analysis of variance (ANOVA) of data from a 3x2 factorial experiment in randomized complete block design  
Experiment 5

Source of variation	d.f.	S.S.	M.S.S.	F. value
Replication	2			
Treatment	5			
Spray (S)	2			
Irrigation (I)	1			
Interaction (S x I)	2			
Error	10			
Total	17			


### 3.11 Biometric Observation

The observations were made at 20 days interval after spray treatment i.e., at 80 (pod fill), 100 (pod maturity) and 120 (harvest) d after sowing. At each sampling five plants from each plot were cut at the soil level, washed with water and dried with blotting paper to record growth, physiological and biochemical characteristics. At harvest, 12 plants, equivalent to 1m<sup>2</sup> land area were removed to record the yield characteristics.

### 3.12 Growth Characteristics

The following growth characteristics were studied at 80, 100 and 120 d after sowing in Experiments 1-4.

1. Plant height
2. Plant leaf area
3. Leaf area index
4. Specific leaf area
5. Specific leaf weight
6. Plant dry weight
7. Dry weight of different plant parts
  - (i) Leaf dry weight
  - (ii) Stem dry weight
  - (iii) Pod dry weight
8. Per cent distribution of dry weight
  - (i) Leaf
  - (ii) Stem
  - (iii) Pod
9. Leaf fresh weight
10. Leaf turgid weight
11. Relative water content in leaf



In Experiment 5, the following growth characteristics were studied.

1. Plant leaf area

## 2 Plant dry weight

### 3.12.1 Plant leaf area

Leaf area was determined by gravimetric method. Leaf area of about 10 % of total leaves from each treatment was determined by tracing on a graph sheet and dry weight for these leaves was recorded. Leaf area/plant was computed by using leaf dry weight/plant and dry weight of those leaves for which the area was estimated (Watson, 1958) using the following formula

$$LA = \frac{LA_1 \times W_2}{W_1}$$

$LA_1$  = Leaf area of the leaves traced on graph paper

$W_1$  = Dry weight of the leaves for which area was traced on graph paper

$W_2$  = Total leaf dry weight/plant

### 3.12.2 Leaf area index

Leaf area index (LAI) was calculated on the formula suggested by Watson (1958)

$$LAI = \frac{\text{Leaf area}}{\text{Ground area}}$$

### 3.12.3 Specific leaf area

Specific leaf area (SLA) represents the leaf area of unit amount of leaf biomass. This was calculated as leaf area divided by leaf dry weight

$$SLA = \frac{\text{Leaf area}}{\text{Leaf weight}}$$

### 3.12.4 Specific leaf weight

Specific leaf weight (SLW) is the measurement of allocation of leaf dry weight per unit of leaf area. It was calculated as leaf dry weight divided by leaf area

$$SLW = \frac{\text{Leaf weight}}{\text{Leaf area}}$$

### 3.12.5 Dry weight

Dry weight of different plant parts was recorded after drying them in an oven for 48 h at 80 °C.

### 3.12.6 Leaf fresh weight and turgid weight

The sampled plants were divided into different parts like leaf, stem and pod corresponding to different sampling stages. Leaf fresh weight was recorded and samples were placed in water and their turgid weight was recorded accordingly.

### 3.12.7 Leaf relative water content

Relative water content (RWC) of leaves was expressed as percentage of the water content of the fully turgid leaves and calculated as follows.

$$\text{Relative water content (\%)} = \frac{W_F - W_d}{W_t - W_d} \times 100$$

Here,  $W_t$  = Weight of the fully turgid leaves

$W_F$  = Fresh weight of the leaves

$W_d$  = Dry weight of the leaves

## 3.13 Physiological Characteristics

Following physiological parameters were studied at 80 and 100 d after sowing in all experiments.

1. Rate of photosynthesis
2. Stomatal conductance
3. Internal CO<sub>2</sub> concentration
4. Transpiration rate
5. Carboxylation efficiency
6. Photosynthetic water use efficiency
7. Plant water use efficiency

In Experiments 3-5, in addition to the above mentioned parameters following characteristics were also studied.

1. 1-Aminocyclopropane carboxylic acid (ACC) content
2. ACC oxidase
3. Ethylene evolution

### 3.13.1 Rate of photosynthesis

The data on rate of photosynthesis, stomatal conductance, internal CO<sub>2</sub> concentration and transpiration rate were measured in fully expanded top leaf of each main axis of plant using the Li COR-6200 Portable photosynthesis system (Nebraska, USA) at 1250  $\mu\text{mol m}^{-2} \text{s}^{-1}$  photosynthetically active radiation at 1100-1200 hours (temperature 23°C, relative humidity 72 %). The care was taken to use leaves of the same age for measurement of photosynthesis in control and treated plants. Each observation was replicated twice for control and treated plants and average of these was taken as a replicate.

### 3.13.2 Carboxylation efficiency

Carboxylation efficiency was computed by dividing the photosynthesis by internal CO<sub>2</sub> concentration and expressed in percentage.

$$\text{Carboxylation efficiency (\%)} = \frac{\text{Photo}}{C_{\text{int}}} \times 100$$

### 3.13.3 Photosynthetic water use efficiency

Photosynthetic water use efficiency is a measure of CO<sub>2</sub> assimilation and has direct relation with rubisco activity (Vanden Boogard *et al.*, 1996). It was expressed as the ratio of photosynthesis rate to stomatal conductance (A/g<sub>s</sub>) to avoid effects of small differences in vapour pressure between measurements (Von Cammerer and Farquhar, 1981).

#### 3.13.4 Plant water use efficiency

Water use efficiency was calculated as the ratio of biomass to cumulative transpiration (Vanden Boogard *et al.*, 1996).

$$\text{Plant water use efficiency} = \frac{\text{Biomass}}{\text{Transpiration}}$$

#### 3.13.5 Estimation of 1-aminocyclopropane-1-carboxylic acid (ACC)

At sampling times, 1 g of leaf sample was homogenized in 10 ml of 80 % ethanol and then extracted under reflux with boiling ethanol for 30 min. It was then filtered through four layers of fine gauze and evaporated to dryness. The process was repeated with the addition of 0.5 ml chloroform. The residue was suspended in 2 ml double-distilled water, and then centrifuged for 20 min at 27000 g (Mc Keon *et al.* 1982). The ACC content in the aqueous extract was determined by its chemical conversion to ethylene after the addition of NaOCl (Lizada and Yang 1979). ACC content in the extract was estimated assuming that the percentage conversion of ACC to ethylene in the extract was identical to that of ACC added as an internal standard.

#### 3.13.6 Assay of ACC oxidase activity

ACC oxidase activity was measured as ability of leaves to convert exogenous ACC to ethylene. Leaf sample (0.5 g) was cut into small pieces and incubated with 0.5 mL of 5 mM ACC. After flushing with air, the tubes were capped and incubated in light for 1 h under the same conditions used for plant growth (Vioque and Castellano, 1994). The ethylene evolved during incubation was determined on gas chromatograph (Nucon GLC 5700, India), as described in Section 3.11.3

#### 3.13.7 Ethylene evolution

For ethylene measurement leaf material was trimmed to small pieces, weighed and placed in 30 ml tubes, which were stoppered with rubber secure cap and placed in light for 2 h under the same conditions as used for plant

growth. Ethylene content in the gas phase of tubes was determined from 1 ml samples were removed from the tubes and injected into a Nucon GLC 5700 gas chromatograph fitted with a flame ionisation detector and 1.8 m x 4 mm glass column packed with 80-100 mesh porapak-N. The oven temperature was 100<sup>o</sup>/h C. The flow rates of nitrogen and hydrogen were 30 ml min<sup>-1</sup>, and of oxygen was 300 ml min<sup>-1</sup>. Ethylene identification was based on the retention time compared with a pure ethylene standard

### 3.14 Biochemical Characteristics


Following parameters were studied:

1. Nitrate reductase activity
2. N content
3. N accumulation

Nitrate reductase activity in fresh leaves was estimated and dried plant material, collected at different sampling stages was used for the estimation of N content and its accumulation. The details of the estimation procedure are given in the following pages.

#### 3.14.1 Assay of nitrate reductase activity

Enzyme nitrate reductase catalyses the reduction of nitrate to nitrite.



Leaf nitrate reductase activity (NRA) was estimated *in vivo* by the method of Jaworski (1971), which is based on the reduction of nitrate to nitrite. The nitrite formed was then determined spectrophotometrically.

Fresh leaf sample (200 mg) was transferred to polythene vials, containing 2.5 mL of phosphate buffer (pH 7.0) and 0.5 mL of 0.2 M potassium nitrate solution (Appendix) was added followed by addition of 2.5 mL of 5 % isopropanol (Appendix). Finally, 2 drops of 0.5 % chloramphenicol solution (Appendix) was added to avoid bacterial growth in the medium. These vials were incubated for 2 hours in dark at 30°C.

#### **3.14.1.1 Colour development**

Incubated mixture (0.4 mL) was taken in a test tube to which 0.3 mL of 1 % sulphanilamide (Appendix) and 0.02 % naphthylethylenediamine hydrochloride (NED-HCl) were added. The test tube was left for 20 minutes for maximum colour development. The mixture was diluted again to 5 mL with sufficient amount of double distilled water. Then absorbance was read at 540 nm using a blank on spectrophotometer (SL171 Elico, Hyderabad, India).

#### **3.14.1.2 Standard curve for NRA**

30 mg of sodium nitrite ( $\text{NaNO}_2$ ) was dissolved in 100 mL double distilled water. From this solution, various amounts was taken in ten different test tubes, viz., 0.2, 0.4, 0.6, 0.8, 1.0, 1.2, 1.4, 1.6, 1.8, and 2.0 mL. To these 0.3 mL of 1 % sulphanilamide and 0.02 % NED-HCl were added. The solutions were diluted to 5 mL with double distilled water and the absorbance was read at 540 nm, using a blank on spectrophotometer (SL171 Elico, Hyderabad, India).

#### **3.14.2 Nitrogen content**

##### **3.14.2.1 Digestion of plant samples for N content**

Dried plant material was powdered and passed through 70- mesh screen and stored in polythene vials. This plant material was used for the estimation of N content.

100 mg of the oven dried powder from each replicate was transferred to a 50 mL Kjeldahl flask to which 2 mL of sulphuric acid was added. The content of the flask was heated on temperature-controlled assembly for about 2 hours to allow complete reduction of nitrates present in the plant material by organic matter itself. As a result, the contents of the flask were turned black. After cooling the flask for about 15 minutes, 0.5 mL of 30 %  $\text{H}_2\text{O}_2$  was added drop by drop and the solution was heated again until the colour turns from black to light yellow. Again after cooling for 30 minutes an additional 3-4 drops of 30 %  $\text{H}_2\text{O}_2$  were added, followed by heating for another 15 minutes.



The process was repeated until the contents of the flask turned colourless. The peroxide digested material was transferred from Kjeldahl flask to 100 mL volumetric flask with three washings of double distilled water. The volume of the flask was made up to the mark with double distilled water. This peroxide digested material was used for the estimation of N content.

#### **3.14.2.2 Estimation of nitrogen**

Nitrogen was estimated according to Lindner (1944). A 10 mL aliquot of the digested material was taken in 50 mL volumetric flask. To this, 2 mL of 2.5 N NaOH and 1 mL of 10 % sodium silicate solution were added which neutralizes excess of acid and prevents turbidity. The volume of the solution was made up to the mark with double distilled water. In a 10 mL graduated test tube, 5 mL of this solution was taken and 0.5 mL of Nessler's reagent was added. The final volume was made up with double distilled water. The contents of the tube were allowed to stand for 5 minutes for maximum colour development and absorbance was read at 525 nm on spectrophotometer (SL171 Elico, Hyderabad, India).

##### **3.14.2.2.1 Standard curve for nitrogen**

50 mg ammonium sulphate was dissolved in 1 litre double distilled water. From this solution, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 and 1.0 ml were pipetted to ten different test tubes. The solution in each test tube was diluted to 5 mL with double distilled water. In each test tube, 0.5 mL Nessler's reagent was added and after 5 minutes, the absorbance was read at 525 nm on spectrophotometer (SL171 Elico, Hyderabad, India). A blank was run with each set of determination. Standard curve was plotted using different concentrations of ammonium sulphate solution versus absorbance and with the help of this standard curve the amount of nitrogen present in the sample was determined.

### **3.14.3 N accumulation**

Nitrogen content of the plant at different sampling stages and their respective dry matter at these stages were used (as a product) to calculate the N accumulation

### **3.15 Yield Characteristics**

Following parameters were recorded at harvest.

1. Number of pods per plant
2. Number of seeds per pod
3. 1000 seed weight
4. Seed yield
5. Biological yield
6. Harvest index
7. Oil yield

In addition to the above parameters, following parameters were also included in the study in Experiments 3-5 because these experiments included nitrogen as a treatment.

1. Seed N
2. Nitrogen harvest index
3. Nitrogen yield potential

#### **3.15.1 Number of pods per plant**

At harvest, 12 plants from each treatment equivalent to 1 m<sup>2</sup> land were removed. The pods were collected and counted.

#### **3.15.2 Number of seeds per pod**

Seeds of 12 pods from each plant in a treatment were collected and counted.

#### **3.15.3 1000 seed weight**

From the produce of the plot, 1000 seeds were randomly drawn and the weight was recorded.

#### 3.15.4 Seed yield

Total seeds from one-meter square area of the plot were cleaned and weighed to compute the seed yield.

#### 3.15.5 Biological yield

Total biological yield from one-meter square area was recorded from sun-dried samples before threshing.

#### 3.15.6 Harvest index

Harvest index was computed by dividing the seed yield with biological yield and expressed in percentage.

$$\text{Harvest index (\%)} = \frac{\text{Seed yield}}{\text{Biological yield}} \times 100$$

#### 3.15.7 Seed N content per plant

Seed N content was determined as a product of N concentration and the dry weight of the seed.

#### 3.15.8 Nitrogen harvest index

Nitrogen harvest index is a measure of distribution of nitrogen from vegetative part to seed. This was calculated by dividing the seed N with plant N.

$$\text{Nitrogen harvest index} = \frac{\text{Seed N}}{\text{Plant N}}$$

#### 3.15.9 Nitrogen yield potential

Nitrogen yield potential denotes the efficiency of plants to mobilize total N contained in plants to seed during pod fill. This was calculated as a product of nitrogen harvest index and seed N i.e.

$$\text{Nitrogen yield merit} = \text{Nitrogen harvest index} \times \text{Seed N}$$

### 3.15.10 Oil yield

The per cent oil content in seeds when multiplied with seed yield gave the oil yield.

### 3.16 Quality Characteristics

The seed samples were crushed to get a fine meal for extracting the oil after separating them from extraneous material. The oil was analysed for following quality parameters.

1. Oil content
2. Acid value
3. Iodine value
4. Saponification value

#### 3.16.1 Determination of oil content

25 g of ground seeds meal was transferred to a Soxhlet apparatus and sufficient quantity of petroleum ether was added. The apparatus was kept on a hot water bath running at 60 °C for about 6 h, for extraction of oil. Petroleum ether from the extracted oil was evaporated after some time. The extracted oil was expressed as a percentage by mass of the seeds and was calculated by the following formula:

$$\text{Oil content (\%)} = \frac{m_0}{m_s} \times 100$$

Here,  $m_0$  = Sum of the mass of oil

$m_s$  = Seed sample mass

#### 3.16.2 Determination of acid value

Acid value of oil is the amounts of potassium hydroxide spend to neutralize free acid in one gram of oil. It was determined by the following method (Anonymous, 1970).

2 g of oil was dissolved in 50 mL solvent mixture of 95 % alcohol and diethyl ether (1:1) in a 250 mL conical flask. Titration was carried out with 0.1

N potassium hydroxide. Phenolphthalien was used as an indicator and the amount of mL 'a' of 0.1 N NaOH required was noted. The acid value was calculated by the following formula.

$$\text{Acid value} = \frac{\text{'a'} \times 0.05661 \times 1000}{W}$$

Here, 'a' = ml of 0.1N KOH used in titration

$W = \overset{w}{\text{weight of oil}}$

### 3.16.3 Determination of iodine value

Iodine value of oil is the number of gm of iodine absorbed by 100 gm of oil and expressed as the weight of iodine. It was determined by using iodine monochloride method described below (Anonymous, 1970).

Oil (2 gram) was taken in a dry ground neck flask to which 10 mL carbon tetrachloride and 20 mL iodine monochloride solution were added. The flask was stopper and allowed to stand in a dark place for about 30 minutes. After 30 minutes, 15 mL potassium iodide and 100 mL double distilled water was poured into it with proper shaking. Titration was carried out with 0.1 N sodium thiosulphate ( $\text{Na}_2\text{S}_2\text{O}_3$ ) solution using starch solution as an indicator. Number of mL 'a' of sodium thiosulphate used was noted. For blank, similar operation was put in practice without the oil and the number of ml 'b' of 0.1 N sodium thiosulphate solution used was noted. Iodine value was calculated by the following formula (Anonymous, 1970).

$$\text{Iodine value} = \frac{(b - a) \times 0.01269 \times 100}{W}$$

Here,  $a = \overset{\checkmark}{\text{number of mL of 0.1 N Na}_2\text{S}_2\text{O}_3 \text{ solution used in the sample}}$

$b = \overset{\checkmark}{\text{number of mL of 0.1 N Na}_2\text{S}_2\text{O}_3 \text{ solution used in blank}}$

$W = \overset{w}{\text{weight of oil}}$

#### 3.16.4 Determination of saponification value

Saponification value of oil is the amount of mg of KOH consumed by 1gm of the oil to neutralize the fatty acid resulting from complete hydrolysis.

2 gm oil was taken in a 250 mL conical flask to which 25 mL of 0.5 N KOH was added. The flask was attached with reflux condenser and heated on water bath for about 1h with frequent rotation of the contents of the flask. The excess of alkali was titrated with 0.5 N HCl. The number of mL (a) of 0.5 N HCl was noted. A similar practice was repeated without oil and the number of mL (b) of 0.5 N HCl required was noted (Anonymous, 1970). Saponification value was calculated by the following formula:

$$\text{Saponification value} = \frac{(b - a) \times 0.02805 \times 1000}{W}$$

Where a and b are number of mL of 0.05 N HCl used in the sample and blank titration respectively, and W is weight of oil

#### 3.17 Statistical Analysis

All the experimental data were subjected to statistical analysis by adopting analysis of variance techniques according to the design of the experiments (Gomez and Gomez, 1984) and the significance of the results were determined at 5% levels of probability. Pooled analysis of Experiments 1 and 2 and Experiments 3 and 4 was performed from a split plot design to evaluate the effect of combinations of factors. If the data were found significant, Least Significant Difference (LSD) was calculated. Correlation values between various traits were also worked out.

***CHAPTER-4***  
***EXPERIMENTAL***  
***RESULTS***

## CONTENTS

### EXPERIMENTAL RESULTS

4.1	Experiment 1 and 2	68
4.1.1	Growth characteristics	68
4.1.1.1	Plant height	68
4.1.1.2	Plant leaf area	69
4.1.1.3	Leaf area index	69
4.1.1.4	Specific leaf area	70
4.1.1.5	Specific leaf weight	70
4.1.1.6	Plant dry weight	70
4.1.1.7	Dry weight of different plant parts	71
4.1.1.7.1	Leaf dry weight	71
4.1.1.7.2	Stem dry weight	71
4.1.1.7.3	Pod dry weight	71
4.1.1.8	Per cent distribution of dry weight in plant parts	72
4.1.1.8.1	Leaf	72
4.1.1.8.2	Stem	72
4.1.1.8.3	Pod	73
4.1.1.9	Leaf fresh weight	73
4.1.1.10	Leaf turgid weight	73
4.1.1.11	Leaf relative water content	74
4.1.2	Physiological characteristics	74
4.1.2.1	Rate of photosynthesis	74
4.1.2.2	Stomatal conductance	75
4.1.2.3	Internal CO <sub>2</sub> concentration	75
4.1.2.4	Transpiration rate	75
4.1.2.5	Carboxylation efficiency	76
4.1.2.6	Photosynthetic water use efficiency	76
4.1.2.7	Plant water use efficiency	76
4.1.3	Biochemical characteristics	77
4.1.3.1	N content	77
4.1.3.2	N accumulation	77
4.1.4	Yield characteristics	78
4.1.4.1	Number of pods per plant	78
4.1.4.2	Number of seeds per pod	78
4.1.4.3	1000 seed weight	78
4.1.4.4	Seed yield	79
4.1.4.5	Biological yield	79
4.1.4.6	Harvest index	79
4.1.4.7	Oil yield	79
4.1.5	Quality characteristics	80



4.1.5.1	Oil content	80
4.1.5.2	Acid value	80
4.1.5.3	Iodine value	80
4.1.5.4	Saponification value	80
4.1.6	Pooled analysis of Experiment 1 and 2	81
4.2	Experiment 3 and 4	81
4.2.1	Growth characteristics	82
4.2.1.1	Plant height	82
4.2.1.2	Plant leaf area	82
4.2.1.3	Leaf area index	83
4.2.1.4	Specific leaf area	83
4.2.1.5	Specific leaf weight	83
4.2.1.6	Plant dry weight	84
4.2.1.7	Dry weight of different plant parts	84
4.2.1.7.1	Leaf dry weight	84
4.2.1.7.2	Stem dry weight	85
4.2.1.7.3	Pod dry weight	85
4.2.1.8	Per cent distribution of dry weight in plant parts	86
4.2.1.8.1	Leaf	86
4.2.1.8.2	Stem	86
4.2.1.8.3	Pod	87
4.2.1.9	Leaf fresh weight	87
4.2.1.10	Leaf turgid weight	88
4.2.1.11	Leaf relative water content	88
4.2.2	Physiological characteristics	89
4.2.2.1	Rate of photosynthesis	89
4.2.2.2	Stomatal conductance	89
4.2.2.3	Internal CO <sub>2</sub> concentration	89
4.2.2.4	Transpiration rate	90
4.2.2.5	Carboxylation efficiency	90
4.2.2.6	Photosynthetic water use efficiency	91
4.2.2.7	Plant water use efficiency	91
4.2.2.8	ACC content	92
4.2.2.9	ACC oxidase	92
4.2.4.10	Ethylene evolution	93
4.2.3	Biochemical characteristics	93
4.2.3.1	Nitrate reductase activity	93
4.2.3.2	N content	93
4.2.3.3	N accumulation	94
4.2.4	Yield characteristics	94
4.2.4.1	Number of pods per plant	94
4.2.4.2	Number of seeds per pod	94
4.2.4.3	1000 seed weight	95

4.2.4.4	Seed yield	95
4.2.4.5	Biological yield	95
4.2.4.6	Harvest index	95
4.2.4.7	Oil yield	95
4.2.4.8	Seed nitrogen per plant	96
4.2.4.9	Nitrogen harvest index	96
4.2.4.10	Nitrogen yield potential	96
4.2.5	Quality parameters	97
4.2.5.1	Oil content	97
4.2.5.2	Acid value	97
4.2.5.3	Iodine value	97
4.2.5.4	Saponification value	97
4.2.6	Pooled analysis of Experiment 3 and 4	97
4.3	Experiment 5	98
4.3.1	Growth characteristics	98
4.3.2	Physiological characteristics	98
4.3.3	Biochemical characteristics	99
4.3.4	Yield characteristics	99
4.3.5	Quality characteristics	99

## **EXPERIMENTAL RESULTS**

It has been mentioned in Materials and Methods section that the two experiments (1 and 2) and (3 and 4) were conducted on the same lines with difference of irrigated and non-irrigated conditions. The results of Experiment 1 and 2, and Experiment 3 and 4 are described together in the following pages. The data of Experiment 1 (Tables 7–34), Experiment 2 (Tables 35–62), pooled analysis of Experiment 1 and 2 (Tables 63–71), Experiment 3 (Tables 72–104), Experiment 4 (105–137), pooled analysis of Experiment 3 and 4 (Tables 138–146) and Experiment 5 (Tables 147–180) are given at the end of the thesis.

### **4.1 Experiment 1 and 2**

In these experiments concentrations of ethrel applied at 60d after sowing were 0, 100, 200, 400 and 600 $\mu$ L/L on Alankar and PBM16 cultivars of mustard. Growth characteristics, plant height, plant leaf area, leaf area index, specific leaf area, specific leaf weight, plant dry weight, dry weight and its per cent distribution in leaf, stem and pod, leaf fresh weight, leaf turgid weight and leaf relative water content; biochemical characteristics, N concentrations and its accumulation were recorded at 80, 100 and 120 DAS. Physiological characteristics, net photosynthetic rate, stomatal conductance, internal CO<sub>2</sub> concentration, transpiration rate, carboxylation efficiency, photosynthetic water use efficiency and plant water use efficiency were recorded at 80 and 100 DAS. At harvest pod number, seed number, 1000 seed weight, seed yield, biological yield, harvest index, oil yield, oil content, acid value, iodine value and saponification value of oil were determined.

#### **4.1.1 Growth characteristics**

##### **4.1.1.1 Plant height**

Plant height was significantly affected by ethrel spray and both cultivars differed each other at all growth stages. The interaction effect between ethrel

spray and cultivar was non-significant (Tables 7, 35). Similar response was noted in both the experiments.

Spraying of 200 $\mu$ L/L ethrel registered maximum effect on plant height and water spray recorded minimum value for plant height at all growth stages. Among the cultivars, Alankar was taller than PBM16 at all sampling times (Tables 7, 35) in Experiment 1 as well as in Experiment 2.

#### **4.1.1.2 Plant leaf area**

Crop canopy, which is represented by leaf area, was significantly affected by ethrel spray and cultivars also differed significantly at all sampling times in the two experiments. The interaction effect between these two was found significant at 80 DAS in Experiment 1 and 80 and 100 DAS in Experiment 2 (Tables 8, 36).

A concentration of 200 $\mu$ L/L ethrel showed maximum value at all three growth stages and significantly minimum value was recorded for water sprayed under irrigated and non-irrigated conditions.

Alankar showed maximum leaf area as compared to PBM16 at all sampling stages in the two experiments. It was observed that 200 $\mu$ L/L ethrel spray on Alankar gave maximum value and water spray on PBM16 gave minimum value at 80 DAS in Experiment 1. Similar effect was noted at 80 and 100 DAS in Experiment 2 (Tables 8, 36).

#### **4.1.1.3 Leaf area index**

Leaf area index was significantly affected by ethrel spray and cultivars differed significantly at all sampling times. Interaction effect between ethrel spray and cultivar was significant only for 80 DAS. A similar response was noted in both the experiments (Tables 9, 37).

Significantly maximum value was recorded with 200 $\mu$ L/L ethrel spray at every growth stage and minimum value was found in water spray. Alankar showed higher leaf area index than PBM16.

Among different interactions, 200 $\mu$ L/L ethrel spray on Alankar registered maximum value and minimum value was recorded in water spray on PBM16 (Tables 9, 37).

#### **4.1.1.4 Specific leaf area**

Ethrel spray significantly affected specific leaf area and cultivars differed significantly for this characteristic at all sampling times in both the experiments. The interaction effect of these two remained non-significant in both the experiments (Tables 10, 38).

Spray of 200 $\mu$ L/L ethrel and water gave maximum and minimum values respectively in both the experiments. Alankar showed maximum and PBM16 the minimum values.

#### **4.1.1.5 Specific leaf weight**

Values recorded in ethrel spray and cultivar differed significantly at all the sampling times in both the experiments. Interaction effect was noted to be non-significant in both the experiments (Tables 11, 39).

Significantly maximum and minimum values were recorded with water spray and 200 $\mu$ L/L ethrel spray respectively. The trend was similar in the two experiments, and Alankar registered higher value than PBM16.

#### **4.1.1.6 Plant dry weight**

Plant dry weight was affected by ethrel spray at every growth stage. The effect of ethrel spray on cultivars differed significantly at 80 DAS only, however, at other sampling times, the two cultivars responded equally to every spray treatment (Tables 12, 40) in Experiment 1. Highest plant dry weight was recorded with 200 $\mu$ L/L ethrel spray at all sampling times, while the lowest value was found in water sprayed control.

Alankar showed maximum plant dry weight as compared to PBM16. Among interactions, 200 $\mu$ L/L ethrel spray on Alankar proved best and gave significantly maximum value as compared to other interactions. Minimum value was noted in water sprayed on PBM16.

#### **4.1.1.7 Dry weight of different plant parts**

##### **4.1.1.7.1 Leaf dry weight**

Ethrel spray significantly affected leaf dry weight at all sampling times. Ethrel spray on the two cultivars significantly differed at 80DAS in Experiment 1 and at 100 DAS in Experiment 2 (Tables 13, 41).

Maximum leaf dry weight was recorded with 200 $\mu$ L/L ethrel spray at all sampling times in both the experiments. Significantly minimum value was found in water sprayed control treatments.

Among the cultivars, Alankar showed maximum value as compared to PBM16 in the two experiments. Among interactions, 200 $\mu$ L/L ethrel spray on Alankar proved best and gave significantly higher value while the lowest value was recorded in water sprayed on PBM16.

##### **4.1.1.7.2 Stem dry weight**

Stem dry weight was significantly affected by ethrel spray at all sampling times in both the experiments. Effect of ethrel spray on the two cultivars significantly differed at 80 and 100 DAS in Experiment 1, whereas the cultivars responded similarly to the concentrations of ethrel in Experiment 2 at all sampling times (Tables 14, 42).

In both the experiments, 200 $\mu$ L/L ethrel spray resulted in maximum value for stem dry weight, while minimum value was found in water sprayed control.

Alankar registered higher dry weight than PBM16 at all sampling times under irrigated and non-irrigated conditions. Regarding the interaction between ethrel spray and cultivar, it was observed that 200 $\mu$ L/L ethrel spray on Alankar gave maximum value at 80 and 100 DAS.

##### **4.1.1.7.3 Pod dry weight**

Effect of ethrel spray was significant at all sampling times and the interaction effect between spray and cultivar was found significant at 100 DAS in Experiment 1 and 2 (Tables 15, 43).

A 200 $\mu$ L/L ethrel spray gave highest value at all sampling times. Significantly lowest value was recorded for water spray control. At all growth stages, Alankar showed higher value than PBM16.

Among the interactions, 200 $\mu$ L/L ethrel spray on Alankar gave maximum value. However, minimum value was registered with water spray on PBM16 (Tables 15, 43).

#### **4.1.1.8 Per cent distribution of dry weight in plant parts**

##### **4.1.1.8.1 Leaf**

Effect of ethrel spray was found significant at all sampling times in both the experiments. The two cultivars differed significantly at 100 and 120 DAS in Experiment 2. The interaction effect between ethrel spray and cultivar was non-significant in both the experiments (Tables 16, 44).

Significantly maximum value was recorded with 200 $\mu$ L/L ethrel spray at every growth stages, while minimum value was found with water spray.

Value recorded for Alankar was higher than PBM16, when they differed significantly (Tables 16, 44).

##### **4.1.1.8.2 Stem**

Effect of ethrel spray was significant at all the three growth stages in both the experiments. Similarly, cultivars differed each other significantly. The interaction effect was significant at 100 DAS in Experiment 1, but it was non-significant at all sampling times in Experiment 2.

Per cent stem dry weight decreased with the increase in ethrel concentration at all growth stages and water-sprayed control registered maximum value. Significantly minimum value was recorded with 200 $\mu$ L/L ethrel spray. The trend was similar in both the experiments.

At all growth stages, PBM16 gave more value than Alankar. The interaction effect of water spray and PBM16 gave maximum value. Minimum value was given by 200 $\mu$ L/L ethrel sprayed on Alankar (Tables 17, 45).

#### 4.1.1.8.3 Pod

Distribution of dry weight towards pods was significantly influenced by ethrel spray and cultivar also differed significantly. Similar response was noted in the Experiment 1 and 2. However, interaction effect of ethrel spray and cultivar was significant at 100 DAS in Experiment 1 and non-significant at all sampling times in Experiment 2 (Tables 18, 46).

A 200 $\mu$ L/L ethrel spray showed maximum value and significantly minimum value was registered for water sprayed control in both the experiments.

Alankar showed higher value than PBM16 at all sampling times in both the experiments. Spray of 200 $\mu$ L/L ethrel on Alankar proved best and gave maximum value, while minimum value was found in water spray on PBM16 (Tables 18, 46).

#### 4.1.1.9 Leaf fresh weight

Ethrel spray significantly affected plant fresh weight. The ethrel spray on the two cultivars was significant only at 80 DAS in Experiment 1 and non-significant in Experiment 2 (Tables 19, 47). Spray of 200 $\mu$ L/L ethrel gave highest value at all sampling times, while the minimum value was found with water-sprayed control.

Cultivar, Alankar, elicited maximum value as compared to PBM16. Interaction of ethrel spray and cultivar produced higher leaf fresh weight when spray of 200 $\mu$ L/L ethrel was done on Alankar, while the minimum value was found in water sprayed on PBM16.

#### 4.1.1.10 Leaf turgid weight

Turgid weight of leaves was significantly affected by ethrel spray at all sampling times. Spray of ethrel on the two cultivars significantly differed at 80 DAS, whereas the two cultivars responded similarly to concentrations of ethrel in Experiment 2 (Tables 20, 48).



Spray concentration of 200 $\mu$ L/L ethrel showed maximum value at all sampling times in both the experiments. At all sampling times, Alankar showed maximum value as compared to PBM16 in the two experiments.

Regarding interaction effect, it was observed that 200 $\mu$ L/L ethrel spray on Alankar expressed maximum effect. Minimum value was observed in water sprayed on PBM16.

#### **4.1.1.11 Leaf relative water content**

Relative water content in leaf significantly responded towards ethrel spray and for cultivars at all sampling stages in both the experiments. The interaction effect of these two was found significant at 80 DAS in Experiment 1 and at 80 and 100 DAS in Experiment 2 (Tables 21, 49).

Maximum value was recorded for 200 $\mu$ L/L ethrel at all growth stages and significantly minimum value was found in water sprayed control. Alankar showed higher values than PBM16 in both the experiments.

Regarding the interaction between ethrel spray and cultivar, it was noted that 200 $\mu$ L/L ethrel spray on Alankar expressed the maximum value at 80 DAS in Experiment 1. Similarly, 200 $\mu$ L/L ethrel on Alankar proved best at 80 and 100 DAS in Experiment 2 (Tables 21, 49).

### **4.1.2 Physiological characteristics**

#### **4.1.2.1 Rate of photosynthesis**

Rate of photosynthesis significantly reciprocated towards ethrel spray and for cultivar at all sampling stages. The interaction effect between ethrel spray and cultivar remained non-significant. Similar response was noted in both the experiments (Tables 22, 50).

Spraying concentration of 200 $\mu$ L/L ethrel gave maximum value at all sampling stages and significantly minimum value was found in water-sprayed control in both the experiments. Alankar showed higher value than PBM16 at all sampling times (Tables 22, 50).

#### 4.1.2.2 Stomatal conductance

Stomatal conductance was significantly influenced by ethrel spray and cultivars differed significantly at all growth stages, but the interaction effect between ethrel spray and cultivar was non-significant (Tables 23, 51) in both the experiments.

Highest value for stomatal conductance was recorded with 200 $\mu$ L/L ethrel spray at all growth stages and significantly lowest value was found with water- sprayed control in Experiment 1 and Experiment 2. Alankar showed higher value than PBM16.

#### 4.1.2.3 Internal CO<sub>2</sub> concentration

Effect of ethrel spray was significant on internal carbon dioxide concentration and cultivars also differed significantly at all sampling times. The interaction effect was non-significant (Tables 24, 52). A similar response was observed in both the experiments.

Maximum value for internal carbon dioxide concentration was noted with 200 $\mu$ L/L ethrel spray and Alankar exhibited higher values at all sampling times (Tables 24, 52).

#### 4.1.2.4 Transpiration rate

In both the experiments, effect of ethrel spray and cultivar difference was significant at all growth stages, while the interaction effect between these two was found significant at 80 DAS (Tables 25, 53).

At all growth stages, 200 $\mu$ L/L ethrel spray gave maximum value. Significantly minimum value was recorded in water sprayed control. Alankar showed higher value as compared to PBM16 at all sampling stages.

Among various interactions, 200 $\mu$ L/L ethrel spray on Alankar expressed the maximum value. Significantly minimum value was given by water-sprayed control x PBM16 (Tables 25, 53).

#### 4.1.2.5 Carboxylation efficiency

Effect of ethrel spray was found significant at all sampling times and cultivars differed significantly at 100 DAS in Experiment 1 and at all stages in Experiment 2. The interaction effect in both the experiments remained non-significant (Tables 26, 54).

Value recorded with 200 $\mu$ L/L ethrel spray was significantly maximum at all growth stages. Minimum value was recorded with water-sprayed control. Alankar showed maximum value as compared to PBM16 at all sampling stages.

#### 4.1.2.6 Photosynthetic water use efficiency

Photosynthetic water use efficiency was significantly affected by ethrel spray and cultivars differed significantly at all growth stages. The interaction effect was significant at 80 DAS (Tables 27, 55) in both the experiments.

Spraying concentration of 200 $\mu$ L/L ethrel gave highest value at all growth stages. Significantly lowest value was found with water-sprayed control treatment. Value recorded for Alankar was significantly higher as compared to PBM16 at all growth stages.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel spray on Alankar gave maximum value and was at par with that for 200 $\mu$ L/L ethrel spray on PBM16 at 80 DAS. The effect was similar in both the experiments (Tables 27, 55).

#### 4.1.2.7 Plant water use efficiency

Plant water use efficiency significantly reciprocated towards both ethrel spray and for cultivars difference at all sampling stages in the Experiment 1 and 2. The interaction effect was significant only at 80 DAS in Experiment 1 and non-significant in Experiment 2 (Tables 28, 56).

Values recorded for 200 $\mu$ L/L ethrel spray in Experiment 1 and 2 were significantly maximum at all growth stages and minimum values were found in water-sprayed control. Alankar proved better than PBM16.

Interaction effect of 200 $\mu$ L/L ethrel spray on Alankar registered higher water use efficiency, which differed critically from all other combinations.

#### 4.1.3 Biochemical characteristics

##### 4.1.3.1 N content

Nitrogen content in plant tissues significantly responded towards ethrel spray and cultivars also differed significantly at every sampling time in both the experiments. The interaction effect was found to be significant only at 80 DAS in Experiment 1 and at all sampling times in Experiment 2 (Tables 29, 57).

Spray of 200 $\mu$ L/L ethrel and Alankar registered maximum values at all sampling times in the two experiments. Water-sprayed control gave significantly minimum values.


Among different interactions, 200 $\mu$ L/L ethrel spray on Alankar gave maximum value at 80 DAS in Experiment 1 but in Experiment 2, this interaction gave at par value with that for 200 $\mu$ L/L ethrel on PBM16 at 100 DAS.

##### 4.1.3.2 N accumulation

Nitrogen accumulation was significantly influenced by both ethrel spray and for cultivars at all sampling times and the interaction effect was non-significant in Experiment 1. The interaction effect in Experiment 2 was significant at 80 and 100 DAS (Tables 30, 58).

Maximum value was recorded for 200 $\mu$ L/L ethrel spray at all sampling times. Significantly minimum value was found in water sprayed control. Alankar registered maximum value.

Interaction 200 $\mu$ L/L ethrel on Alankar proved best at 80 and 100 DAS in Experiment 2. The values differed significantly from other combinations (Tables 30, 58).



#### **4.1.4 Yield characteristics**

##### **4.1.4.1 Number of pods per plant**

Number of pods per plant was significantly affected by ethrel spray in Experiment 1 and 2. Similarly, in both the experiments, cultivars differed significantly. The interaction effect was non-significant in both the experiments (Tables 31, 59).

Significantly maximum value was recorded for 200 $\mu$ L/L ethrel spray, while the minimum value was found with water-sprayed control in both the experiments.

Among the cultivars, Alankar showed higher value as compared to PBM16 in both the experiments (Tables 31, 59).

##### **4.1.4.2 Number of seeds per pod**

Number of seeds per pod was significantly affected by both ethrel spray and cultivars. The interaction effect was non-significant (Tables 31, 59) in both the experiments.

Highest value for number of seeds per pod was recorded for 200 $\mu$ L/L ethrel spray. Significantly lowest value was found in water sprayed control in Experiment 1 and 2. Alankar showed higher value than PBM16 (Tables 31, 59).

##### **4.1.4.3 1000 seed weight**

Effect of ethrel spray and cultivar difference was found significant, while their interaction was found to be non-significant in both the experiments (Tables 31, 59).

Maximum value was recorded for 200 $\mu$ L/L ethrel spray. Significantly minimum value was noted in water sprayed control. Among the cultivars, Alankar showed higher value than PBM16 (Tables 31, 59). The response was similar in both the experiments.

#### 4.1.4.4 Seed yield

Like yield attributes, seed yield was also significantly influenced by ethrel spray and cultivar difference was also significant. But interaction between these two remained non-significant. Similar response was noted in Experiment 1 and 2 (Tables 32, 60).

Spray concentration of 200 $\mu$ L/L ethrel gave significantly maximum value. Significantly lowest value was recorded for water sprayed control in both the experiments. Alankar gave higher value than PBM16 (Tables 32, 60).

#### 4.1.4.5 Biological yield

Biological yield significantly responded towards both ethrel spray and the cultivar. The interaction effect between ethrel spray and cultivar was found non-significant in both the experiments (Tables 32, 60).

Significantly highest value was registered for 200 $\mu$ L/L ethrel spray, while the lowest value was found in water sprayed control in Experiment 1 and 2.

Value recorded for Alankar was significantly higher as compared to PBM16 in both the experiments (Tables 32, 60).

#### 4.1.4.6 Harvest index

Harvest index was significantly influenced by both ethrel spray and cultivars, but no significant impact was observed when these two factors interacted (Tables 32, 60). Similar response was noted in Experiment 1 and 2.

Effect of ethrel spray at 200 $\mu$ L/L gave significantly maximum value. Significantly minimum value was recorded for water sprayed control treatment. Alankar showed more value for harvest index than PBM16 in Experiment 1 and 2 (Tables 32, 60).

#### 4.1.4.7 Oil yield

Ethrel spray significantly affected oil yield. Cultivars also differed significantly. However, interaction effect was not significant in both the experiments (Tables 33, 61).

Maximum value was recorded for 200 $\mu$ L/L ethrel, which differed from other values recorded in other spray treatments in both the experiments. Alankar out yielded PBM16.

#### **4.1.5 Quality characteristics**

##### **4.1.5.1 Oil content**

Effect of ethrel spray was significant individually as well as on its effects on the cultivars. Cultivars also differed significantly for the oil content (Tables 33, 61) in Experiment 1 and 2.

Spray of 200 $\mu$ L/L ethrel registered significantly highest oil content in both the experiments. Cultivar Alankar proved better than PBM16. This resulted in maximum oil content in Alankar sprayed with 200 $\mu$ L/L ethrel than any other concentrations of ethrel on either cultivars in both the experiments (Tables 33, 61).

##### **4.1.5.2 Acid value**

Effect of ethrel spray and cultivar was significant in both the experiments. The interaction effect between these two was significant in Experiment 1 and non-significant in Experiment 2 (Tables 34, 62).

Ethrel at 200 $\mu$ L/L concentration gave significantly maximum value and Alankar registered highest acid value in both the experiments. Interaction of 200 $\mu$ L/L ethrel spray x Alankar registered higher acid value, which differed critically from all other combinations in Experiment 1 (Tables 34, 62).

##### **4.1.5.3 Iodine value**

Effect of ethrel spray, cultivar and their interaction effect were non-significant in the two experiments (Tables 34, 62).

##### **4.1.5.4 Saponification value**

Saponification value was significantly reciprocated to both ethrel spray and cultivar while the interaction effect between these two remained non-significant (Tables 34, 62). Similar response was noted in Experiment 1 and 2.

Spraying concentration of 200 $\mu$ L/L ethrel gave maximum value. Significantly minimum value was noted for water sprayed control. Alankar registered significantly higher value as compared to PBM16 (Tables 34, 62).

#### 4.1.6 Pooled analysis of Experiment 1 and 2

Pooled analysis of the data of Experiments 1 and 2 was done to evaluate the performance of the cultivars in irrigated and non-irrigated conditions. Also the impact of ethrel spray on the two cultivars under irrigated and non-irrigated experiments was evaluated (Tables 63, 71). The data showed that there was no significant effect of irrigation and cultivars also did not differ significantly under irrigated and non-irrigated conditions. Similarly, ethrel spray was equally effective under the conditions of irrigation. The three-way interaction was found non-significant.

#### 4.2 Experiment 3 and 4

Experiment 3 was conducted under irrigated conditions and Experiment 4 under non-irrigated conditions. These experiments were conducted to investigate the effect of leaf-applied 0, 100 and 200 $\mu$ L/L ethrel (selected on the basis of Experiment 1 and Experiment 2) at 60d after sowing (DAS, flowering stage) on mustard (*Brassica juncea* L.) cultivar Alankar grown with basally applied 0, 40, 60 or 80kg N/ha, on growth, physiological, biochemical, yield and quality characteristics determined at various sampling times as described in Experiment 1 and Experiment 2. Among physiological characteristics, 1-aminocyclopropane-1-carboxylic acid, ACC oxidase and ethylene evolution were also noted at 80 and 100 DAS samplings. Among biochemical characteristics nitrate reductase activity was also recorded at 80 and 100 DAS samplings. Among yield characteristics, seed nitrogen content per plant, nitrogen harvest index and nitrogen yield potential were also studied at harvest. The details of the results are given below and summarized in Tables 72–137.



#### **4.2.1 Growth characteristics**

##### **4.2.1.1 Plant height**

Plant height was significantly affected by ethrel spray and nitrogen at all sampling times in both the experiments. However, the interaction effect between ethrel spray and nitrogen was found significant at 80 and 100 DAS in Experiment 3, and at 80 DAS in Experiment 4 (Tables 72, 105).

At all sampling times, 200 $\mu$ L/L ethrel spray showed significantly maximum value, while the minimum value was recorded in water-sprayed control in the two experiments.

Maximum height was recorded with 80kg N/ha and minimum with 0kg N/ha at all sampling times in both the experiments.

Among interactions, 200 $\mu$ L/L ethrel spray x 80kg N/ha proved best and gave significantly maximum value. Minimum value was recorded with water sprayed control x 0kg N/ha. This was the effect in both the experiments when the effect was significant (Tables 72, 105).

##### **4.2.1.2 Plant leaf area**

Leaf area was significantly affected by ethrel spray and nitrogen application at all sampling times in both the experiments, while their interaction effect was significant at all sampling times only in Experiment 3 (Tables 73, 106).

Maximum value was recorded with 200 $\mu$ L/L ethrel whereas minimum value was found with water-sprayed plants at all sampling times.

At all sampling times, treatment 80kg N/ha gave significantly maximum value, while the minimum value was registered with 0kg N/ha in both the experiments.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel spray x 80kg N/ha gave maximum value. Minimum value was recorded with water-sprayed control x 0kg N/ha (Tables 73, 108).

#### 4.2.1.3 Leaf area index

Application of ethrel spray and nitrogen resulted into significant impact on leaf area index at all sampling times in both the experiments. But the interaction effect between ethrel spray x nitrogen was significant at all sampling days in Experiment 3 only (Tables 74, 107)

Maximum and minimum values were recorded with 200 $\mu$ L/L ethrel spray and water-sprayed control respectively at all sampling times

At all sampling times, treatment 80kg N/ha gave significantly maximum value, whereas minimum value was recorded with 0kg N/ha in both the experiments

Interaction effect of ethrel spray and nitrogen produced higher leaf area index in combination of 200 $\mu$ L/L ethrel spray x 80kg N/ha. Minimum value was recorded with water-sprayed control x 0kg N/ha (Tables 74, 107)

#### 4.2.1.4 Specific leaf area

The effect of ethrel spray and nitrogen proved significant at all sampling times, while their interaction effect was found non-significant in both the experiments (Tables 75, 108)

At all sampling times, 200 $\mu$ L/L ethrel spray gave maximum value and minimum value was recorded with water-sprayed control in both the experiments

Application of nitrogen at 80kg N/ha exhibited maximum value and the minimum value was registered in 0kg N/ha at all sampling times in both the experiments (Tables 75, 108)

#### 4.2.1.5 Specific leaf weight

Effect of ethrel spray and nitrogen was significant at all sampling times, while the interaction effect between these two was non-significant in both the experiments (Tables 76, 109)

At all sampling times, water-sprayed plants gave significantly maximum value, while the minimum value was recorded with 200 $\mu$ L/L ethrel spray

Treatment 0kg N/ha expressed maximum value, while the minimum value was recorded with 80kg N/ha at all sampling times in both the experiments (Tables 76, 109).

#### **4.2.1.6 Plant dry weight**

Ethrel spray and nitrogen significantly affected plant dry weight at all sampling times in both the experiments. The interaction effect between these two was significant at all sampling times in Experiment 3 only.

Maximum value was recorded with 200 $\mu$ L/L ethrel spray and minimum value with water-sprayed control at all sampling times in both the experiments.

At all sampling times, 80kg N/ha gave significantly maximum value, while the minimum value was recorded with 0kg N/ha.

Interaction effect of ethrel spray and nitrogen produced higher plant dry weight in the combination of 200 $\mu$ L/L ethrel x 80kg N/ha. Minimum value was recorded with water-sprayed control x 0kg N/ha (Tables 77, 110).

#### **4.2.1.7 Dry weight of different plant parts**

##### **4.2.1.7.1 Leaf dry weight**

Application of ethrel spray and nitrogen resulted in significant impact on leaf dry weight at all sampling times in both the experiments. The interaction between ethrel and nitrogen was significant at 80 and 100 DAS in Experiment 3, whereas in Experiment 4, interaction effect was significant only at 120 DAS.

Maximum value was recorded with 200 $\mu$ L/L ethrel spray and minimum value with water-sprayed control at all sampling times in Experiment 3 and 4.

Application of nitrogen at 80kg N/ha gave significantly maximum value and differed from other treatments at all sampling times in Experiment 3 and 4.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel spray along with 80kg N/ha proved best compared to any other combinations in both the experiments (Tables 78, 111).

#### 4.2.1.7.2 Stem dry weight

Effect of ethrel spray and basal nitrogen application was significant at all sampling times in Experiments 3 and 4. The interaction effect was significant at 120 DAS in Experiment 3, while in Experiment 4, interaction effect was significant only at 80 DAS (Tables 79, 112).

At all sampling times, 200 $\mu$ L/L ethrel showed maximum value, while minimum value was recorded with water-sprayed control in both the experiments.

Treatment 80kg N/ha exhibited maximum value and the minimum value was registered with 0kg N/ha at all sampling times in both the experiments.

Among interactions, 200 $\mu$ L/L ethrel x 80kg N/ha gave maximum value, while the minimum value was recorded with water-sprayed control x 0kg N/ha in both the experiments (Tables 79, 112).

#### 4.2.1.7.3 Pod dry weight

Pod dry weight was significantly affected by ethrel spray and nitrogen at all sampling times in both the experiments, while their interaction effect was significant at all sampling times in Experiment 3, and at 120DAS in Experiment 4 (Tables 80, 113).

Significantly maximum value was recorded with 200 $\mu$ L/L ethrel spray and minimum value was found with water-sprayed control at all sampling times in Experiments 3 and 4.

At all sampling times, treatment 80kg N/ha gave significantly maximum value, while the minimum value was recorded with 0kg N/ha in both the experiments.

Interaction between ethrel spray and nitrogen produced higher pod dry weight for the combination of 200 $\mu$ L/L ethrel spray x 80kg N/ha. Minimum value was recorded with water-sprayed control x 0kg N/ha in both the experiments (Tables 80, 113).

#### **4.2.1.8 Per cent distribution of dry weight in plant parts**

##### **4.2.1.8.1 Leaf**

Effect of ethrel spray and nitrogen was significant at all sampling times in both the experiments. But the interaction effect between these two was significant at 80 and 100 DAS in Experiment 3 and at 120 DAS in Experiment 4 (Tables 81, 114)

At all sampling times, 200 $\mu$ L/L ethrel spray gave significantly maximum value, while the minimum value was recorded with water-sprayed control in both the experiments.

Application of nitrogen at 80kg N/ha gave significantly higher value, whereas the minimum value was registered with 0kg N/ha at all sampling times in Experiments 3 and 4.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel x 80kg N/ha gave maximum value, which differed critically from other combinations in both the experiments (Tables 81, 114).

##### **4.2.1.8.2 Stem**

Effect of ethrel spray and application of nitrogen was significant at all sampling times in both the experiments. The interaction effect was significant at 80 and 100 DAS in Experiment 3, whereas in Experiment 4, it was significant only at 120DAS (Tables 82, 115).

Water-sprayed plants and 200 $\mu$ L/L ethrel spray exhibited maximum and minimum values respectively at all sampling times in both the experiments.

At all sampling times, 0kg N/ha gave maximum value and minimum value was recorded with 80kg N/ha in Experiment 3 and 4.

Among interactions, water-sprayed control x 0kg N/ha showed maximum value, while minimum value was registered with 200 $\mu$ L/L ethrel x 80kg N/ha (Tables 82, 115).

#### 4.2.1.8.3 Pod

Effect of ethrel spray and nitrogen was significant at all sampling times in Experiments 3 and 4. However, the interaction effect was significant at all sampling times in Experiment 3, whereas in Experiment 4, it was significant at 100 and 120 DAS (Tables 83, 116).

Highest per cent pod dry weight was recorded in 200 $\mu$ L/L ethrel spray, while minimum value was recorded in water sprayed control at all sampling times with both the experiments.

Application of 80kg N/ha resulted in significantly higher pod dry weight in both the experiments.

Regarding interaction effect, it was noted that 200 $\mu$ L/L ethrel x 80kg N/ha proved best and differed significantly from other combinations (Tables 83, 116) in both the experiments.

#### 4.2.1.9 Leaf fresh weight

Leaf fresh weight was significantly affected by ethrel spray and nitrogen at all sampling times in both the experiments. But the interaction between these two was found significant at 80 and 100 DAS in Experiment 3, whereas in Experiment 4, it was significant at 120 DAS (Tables 84, 117).

Highest value of leaf fresh weight was recorded with 200 $\mu$ L/L ethrel spray, while the minimum value was found with water-sprayed control at all sampling times in both the experiments.

Application of nitrogen at 80kg N/ha showed significantly maximum value. Minimum value was recorded with 0kg N/ha at all sampling times in both the experiments.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel spray x 80kg N/ha proved best than any other combinations (Tables 84, 117) at sampling times when effect was significant.

#### 4.2.1.10 Leaf turgid weight

Effect of ethrel spray and application of nitrogen was significant at all sampling times in both the experiments. The interaction effect was significant at 80 and 100 DAS in Experiment 3, whereas in Experiment 4, it was significant only at 120 DAS.

At all sampling times, 200 $\mu$ L/L ethrel showed significantly maximum value. Minimum value was registered with water-sprayed control at both the experiments.

At all sampling times, 80kg N/ha recorded maximum value and the minimum value was noted with 0kg N/ha.

Among interactions, 200 $\mu$ L/L ethrel x 80kg N/ha gave maximum value, which differed significantly from other values. Minimum value was recorded with water-sprayed control x 0kg N/ha (Tables 85, 118).

#### 4.2.1.11 Leaf relative water content

Relative water content in leaf tissues was significantly affected by both ethrel spray and nitrogen at all sampling times in both the experiments. However, the interaction effect between these two was found significant at 80 and 120 DAS only in Experiment 4 (Tables 86, 119).

Spray of 200 $\mu$ L/L ethrel gave significantly maximum value, whereas minimum value was recorded with water-sprayed control at all sampling times in both the experiments.

At all sampling times, 80kg N/ha exhibited significantly maximum value, while minimum value was found with 0kg N/ha in both the experiments.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel spray x 80kg N/ha proved best as compared to any other combinations (Table 86, 119).

## **4.2.2 Physiological characteristics**

### **4.2.2.1 Rate of photosynthesis**

Effect of ethrel spray, nitrogen and their interactions were found significant at all sampling times in both the experiments (Tables 87, 120).

At all sampling times, 200 $\mu$ L/L ethrel spray gave significantly maximum value and the minimum value was recorded with water-sprayed control under

Application of nitrogen at 80kg N/ha gave maximum value, while minimum value was registered with 0kg N/ha at all sampling times in both the experiments

Among interactions, 200 $\mu$ L/L ethrel spray x 80kg N/ha gave maximum value and minimum value was given by water-sprayed control x 0kg N/ha (Tables 87, 120)

### **4.2.2.2 Stomatal conductance**

In Experiment 3 and 4, the effects of ethrel spray and nitrogen were significant, while their interaction effect was non-significant at all sampling times (Tables 88, 121).

At all sampling times, 200 $\mu$ L/L ethrel showed significantly maximum value, while the minimum value was recorded with water-sprayed control in Experiment 3 and Experiment 4

Treatment 80kg N/ha showed maximum value and the minimum value was registered with 0kg N/ha at all sampling times in both the experiments (Tables 88, 121).

### **4.2.2.3 Internal CO<sub>2</sub> concentration**

Effects of ethrel spray and nitrogen were significant at all sampling times in both the experiments. The interaction effect was significant at all sampling times in Experiment 3, whereas in Experiment 4, it was significant only at 80 DAS (Tables 89, 122).



Maximum value was recorded with 200 $\mu$ L/L ethrel, which differed critically from other values and the minimum value was found with water-sprayed control at all sampling times in both the experiments

At all sampling times, application of nitrogen at 80kg N/ha gave maximum value, while the minimum value was recorded with 0kg N/ha

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel x 80kg N/ha proved best, when compared to any other combinations (Tables 89, 122)

#### **4.2.2.4 Transpiration rate**

Rate of transpiration was significantly responded towards both ethrel spray and nitrogen at all sampling times in both the experiments. However, the interaction effect was significant at all sampling times in Experiment 3, whereas in Experiment 4, it was significant only at 100 DAS (Tables 90, 123)

Maximum transpiration rate was recorded with 200 $\mu$ L/L ethrel spray, while minimum value was recorded in water-sprayed control at all sampling times

At all sampling times, treatment 80kg N/ha gave significantly maximum value and minimum value was registered with 0kg N/ha in both the experiments.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel spray x 80kg N/ha gave maximum value. Minimum value was recorded with water-sprayed control x 0kg N/ha in both the experiments (Tables 90, 123).

#### **4.2.2.5 Carboxylation efficiency**

Application of ethrel spray and nitrogen resulted in significant impact on carboxylation efficiency at all sampling times in both the experiments. But the interaction effect between ethrel spray and nitrogen was significant at 100 DAS in Experiment 4, whereas in Experiment 3, the interaction effect was non-significant (Tables 91, 124).

At all sampling times, 200 $\mu$ L/L ethrel spray gave significantly maximum value. Minimum value found in water-sprayed control at all sampling times in both the experiments.

At all sampling times, treatment 80kg N/ha showed significantly maximum value, while the minimum value was noted with 0kg N/ha in both the experiments.

Among interactions, 200 $\mu$ L/L ethrel x 80kg N/ha gave significantly maximum value and minimum value was recorded with water-sprayed control x 0kg N/ha in Experiment 4 at 100 DAS (Tables 91, 124).

#### **4.2.2.6 Photosynthetic water use efficiency**

Effect of ethrel spray and application of nitrogen were significant at all sampling times in both the experiments. The interaction effect was significant only at 80 DAS in Experiment 3, and at both stages (80 and 100 DAS) in Experiment 4 (Tables 92, 125).

In Experiment 3 and 4, greatest values for photosynthetic water use efficiency were recorded with 200 $\mu$ L/L ethrel spray treatment at both sampling time and minimum value was noted in water-sprayed control.

Treatment 80kg N/ha showed maximum values at both sampling times in the two experiments. Significantly maximum value was given by 200 $\mu$ L/L ethrel x 80kg N/ha, while minimum value was recorded with water-sprayed control x 0kg N/ha in both the experiments (Tables 92, 125).

#### **4.2.2.7 Plant water use efficiency**

Plant water use efficiency was significantly affected by ethrel spray and nitrogen at all sampling times in both the experiments. The interaction effect between ethrel spray and nitrogen was significant at 80 DAS in Experiment 4 (Tables 93, 126).

At all sampling times, 200 $\mu$ L/L ethrel spray resulted in significantly maximum value and the minimum value was recorded with water-sprayed control.

Treatment 80kg N/ha gave significantly maximum value, while the minimum value was recorded with 0kg N/ha at all sampling times in both the experiments.

Among interactions, plant water use efficiency was affected by ethrel spray at 200 $\mu$ L/L concentration along with 80kg N/ha (Tables 93, 126).

#### **4.2.2.8 ACC content**

l-aminocyclopropane-1-carboxylic acid content was significantly affected by ethrel spray and nitrogen at all sampling times in both the experiments. The interaction effect between ethrel spray and nitrogen was significant at 100 DAS in Experiment 3, and at all sampling times in Experiment 4 (Tables 94, 127). A concentration of 200 $\mu$ L/L ethrel showed maximum value at all sampling times. Among applied nitrogen levels, 80kg N/ha gave significantly maximum value and the minimum value was recorded with 0kg N/ha at all sampling times in both the experiments.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel sprayed on plants grown with basal 80kg N/ha registered maximum value in both the experiments (Tables 94, 127).

#### **4.2.2.9 ACC oxidase**

Application of ethrel spray and nitrogen resulted in significant impact on ACC oxidase at all sampling times. The interaction effect was significant at 80 DAS in Experiment 3 and at 100 DAS in Experiment 4 (Table 95, 128).

Spray of 200 $\mu$ L/L ethrel and water spray gave significantly maximum and minimum values respectively at all sampling times in both the experiments. Application of nitrogen at 80kg N/ha resulted in maximum value and minimum value was registered with 0kg N/ha at all sampling times in both the experiments (Tables 95, 128).

Among interactions, 200 $\mu$ L/L ethrel x 80kg N/ha proved best, when compared to any other combinations (Tables 95, 128).

#### **4.2.4.10 Ethylene evolution**

Ethrel spray and nitrogen significantly affected the ethylene evolution at all sampling times in both the experiments. The interaction effect between these two was significant at 80 DAS in Experiment 4 (Tables 96, 129).

Ethylene evolution increased with increase in ethrel spray at both sampling times in the two experiments. Similarly, increase in nitrogen levels increased ethylene evolution. Such effect was seen in spray of 200 $\mu$ L/L ethrel on plants grown with 80kg N/ha, which showed significantly maximum value (Tables 96, 129).

### **4.2.3 Biochemical characteristics**

#### **4.2.3.1 Nitrate reductase activity**

Nitrate reductase activity in leaf tissues was significantly affected by ethrel spray and nitrogen at all sampling times in both the experiments. But the interaction effect between ethrel spray and nitrogen was non-significant in both the experiments (Tables 97, 130).

Spray of 200 $\mu$ L/L ethrel and water spray gave significantly maximum and minimum values respectively at all sampling times.

Application of nitrogen at 80kg N/ha resulted in maximum value and minimum value was registered with 0kg N/ha at all sampling times in both the experiments (Tables 97, 130).

#### **4.2.3.2 N content**

Nitrogen content in plant tissues significantly responded towards both ethrel spray and levels of nitrogen at all sampling stages in both the experiments. The interaction effect was significant at 80 DAS and 120 DAS in Experiment 3 and at 80 DAS in Experiment 4 (Tables 98, 131).

Spray of 200 $\mu$ L/L ethrel showed significantly maximum value at all sampling times. Among applied nitrogen levels, 80kg N/ha showed significantly maximum values at all sampling times in both the experiments.

Among interactions, 200 $\mu$ L/L ethrel x 80kg N/ha gave significantly maximum value and minimum value was recorded with water-sprayed control x 0kg N/ha in both the experiments (Tables 98, 131).

#### **4.2.3.3 N accumulation**

Nitrogen accumulation was significantly affected by ethrel spray and nitrogen at all sampling times. The interaction effect was significant at 80 and 100 DAS in Experiment 3 and at 80 and 120 DAS in Experiment 4 (Tables 99, 132).

At all sampling times, 200 $\mu$ L/L ethrel spray gave significantly maximum value, while the minimum value was recorded with water-sprayed control.

Application of nitrogen at 80 and 0kg N/ha showed significantly maximum and minimum values respectively at all sampling times in both the experiments.

Regarding the interaction effect, it was observed that accumulation of nitrogen was significantly affected by 200 $\mu$ L/L ethrel x 80kg N/ha (Tables 99, 132).

#### **4.2.4 Yield characteristics**

##### **4.2.4.1 Number of pods per plant**

Ethrel spray and nitrogen significantly affected pod number in both the experiments. The interaction effect between ethrel spray and nitrogen was non-significant in both the experiments (Tables 100, 133).

Ethrel spray at 200 $\mu$ L/L concentration and water spray resulted in significantly maximum and minimum values respectively. Basal 80kg N/ha proved best for pod number in both the experiments (Tables 100, 133).

##### **4.2.4.2 Number of seeds per pod**

Number of seeds per pod was significantly influenced by ethrel spray and nitrogen in both the experiments and the interaction effect was significant in Experiment 4 (Tables 100, 133).

Individual as well as combined effect of 200 $\mu$ L/L ethrel spray and 80kg N/ha proved superior than any other treatments (Tables 100, 133).

#### **4.2.4.3 1000 seed weight**

The seed weight was also significantly affected by ethrel spray and nitrogen in both the experiments, proving 200 $\mu$ L/L ethrel and 80kg N/ha most effective (Tables 100, 133).

#### **4.2.4.4 Seed yield**

Seed yield of the crop was significantly affected by ethrel spray and nitrogen, but their interaction effect was non-significant in both the experiments (Tables 101, 134).

Seed yield was significantly maximum in 200 $\mu$ L/L ethrel concentration and 80kg N/ha, while the minimum value was noted with water-sprayed control in both the experiments (Tables 101, 134).

#### **4.2.4.5 Biological yield**

Biological yield was significantly influenced by ethrel spray and nitrogen in both the experiments. But the interaction effect between these two was found significant only in Experiment 3 (Tables 101, 134).

Concentration of 200 $\mu$ L/L ethrel spray and 80kg N/ha individually and interactively gave maximum value and the minimum value was registered with water-sprayed control (Tables 101, 134).

#### **4.2.4.6 Harvest index**

Individual and combined effect of ethrel spray and nitrogen were found significant in both the experiments (Tables 101, 134).

Significantly maximum value was recorded with 200 $\mu$ L/L ethrel spray and 80kg N/ha and their interaction in both the experiments (Tables 101, 134).

#### **4.2.4.7 Oil yield**

Application of ethrel spray and nitrogen resulted in significant impact on oil yield. But the interaction effect was non-significant (Tables 103, 136).

Maximum value was recorded with 200 $\mu$ L/L ethrel and the minimum value was noted with water-sprayed control in both the experiments.

Application of nitrogen at 80kg N/ha showed maximum value, while the minimum value was recorded with 0kg N/ha (Tables 103, 136).

#### **4.2.4.8 Seed nitrogen per plant**

Effect of ethrel spray, nitrogen and their interaction were found to be significant in both the experiments (Tables 102, 135).

Among ethrel treatments, 200 $\mu$ L/L ethrel spray gave maximum value and the minimum was found with water-sprayed control in both the experiments.

Among applied nitrogen levels, 80kg N/ha registered highest value while control recorded significantly lowest value.

Regarding the interaction effect, it was observed that 200 $\mu$ L/L ethrel spray x 80kg N/ha proved best in comparison to any other combinations (Tables 102, 135).

#### **4.2.4.9 Nitrogen harvest index**

Nitrogen harvest index was non-significant for ethrel spray, applied nitrogen and their interaction (Tables 102, 135).

#### **4.2.4.10 Nitrogen yield potential**

Effect of ethrel spray application, levels of nitrogen and their interaction effect were found significant in both the experiments (Tables 102, 135).

Highest nitrogen yield potential was recorded with 200 $\mu$ L/L ethrel spray and the minimum value was found with water-sprayed control.

Application of nitrogen at 80kg N/ha gave maximum value and the minimum value was recorded with 0kg N/ha in both the experiments.

Among interactions, 200 $\mu$ L/L ethrel spray x 80kg N/ha gave significantly maximum value, while the minimum value was registered with water-sprayed control x 0kg N/ha (Tables 102, 135).

#### **4.2.5 Quality parameters**

##### **4.2.5.1 Oil content**

Effect of ethrel spray was significant, while the effect of nitrogen and interaction effect was non-significant in both the experiments (Tables 103, 136)

Ethrel spray increased oil content and 200 $\mu$ L/L proved most effective in both the experiments (Tables 103 136)

##### **4.2.5.2 Acid value**

Effect of ethrel spray was significant, while nitrogen effect and their interaction effect was non-significant (Tables 104, 137)

Spray of 200 $\mu$ L/L ethrel showed maximum value, while the minimum was recorded with water-sprayed control (Tables 104, 137)

##### **4.2.5.3 Iodine value**

Effect of ethrel spray only was significant in both the experiments (Tables 104, 137) Maximum iodine value was recorded with 200 $\mu$ L/L ethrel

##### **4.2.5.4 Saponification value**

Saponification value was significantly affected by ethrel spray, while nitrogen and the interaction effect was found non-significant in both the experiments (Tables 104, 137)

Concentration of 200 $\mu$ L/L ethrel spray gave maximum value and the minimum value was recorded with water-sprayed control in both the experiments (Tables 104, 137)

#### **4.2.6 Pooled analysis of Experiment 3 and 4**

Pooled analysis of the data of Experiments 3 and 4 was done to evaluate the effect of ethrel spray and nitrogen application under irrigated and non-irrigated conditions (Tables 138–146) Irrigation did not significantly affect the characteristics studied The nitrogen application and ethrel spray under irrigated and non-irrigated conditions were equally effective The three-way interaction was found non-significant



### 4.3 Experiment 5

This experiment was conducted to confirm the effects of 200 $\mu$ L/L ethrel spray (reported in Experiments 1–4) on growth, physiological, biochemical, yield and quality characteristics of mustard grown under irrigated and non-irrigated conditions. For this, plants were grown with uniform basal dose of 80kg N/ha and were sprayed with either 0, 200 $\mu$ L/L ethrel or 1mM of silver thiosulphate (STS; a ethylene action inhibitor) at 60d after sowing (DAS; flowering stage). Growth of the plants was assessed for plant leaf area and plant dry weight, whereas other characteristics studied were similar as in Experiment 3 and Experiment 4. The details of the results are given below and summarised in Tables 147–166.

#### 4.3.1 Growth characteristics

Spray significantly affected the plant leaf area and plant dry weight at all sampling times. The effect of spray under irrigated and non-irrigated plants for these two characteristics was similar (Tables 147, 148). Highest plant dry weight was recorded with 200 $\mu$ L/L ethrel spray, but the plants sprayed with silver thiosulphate (STS) showed lowest plant dry weight.

#### 4.3.2 Physiological characteristics

Among the physiological characteristics studied, spray of 200 $\mu$ L/L ethrel enhanced the traits in comparison to no ethrel spray. However, spray of silver thiosulphate (1mM) reduced the effects of ethrel spray and value decreased to lower than the control.

Irrigation effect and interaction of spray with irrigation were non-significant. Similar results were observed for the photosynthetic rate, stomatal conductance, internal CO<sub>2</sub> concentration, carboxylation efficiency, photosynthetic water use efficiency, plant water use efficiency, 1-aminocyclopropane carboxylic acid content, ACC oxidase and ethylene evolution (Tables 149–158).

#### **4.3.3 Biochemical characteristics**

Nitrate reductase activity, nitrogen content and nitrogen accumulation were significantly affected by spray. Ethrel spray enhanced nitrogen content and accumulation in plants compared with no ethrel spray. Silver thiosulphate spray decreased these contents. Under irrigated and non-irrigated conditions, spray of ethrel was equally effective (Tables 159–161).

#### **4.3.4 Yield characteristics**

Number of pods and seeds, 1000 seed weight, seed yield, biological yield, harvest index, oil yield, seed nitrogen content per plant and nitrogen yield potential were significantly increased by 200 $\mu$ L/L ethrel spray. Silver thiosulphate spray confirmed the positive effect of ethylene as ethrel spray (Tables 162–164).

#### **4.3.5 Quality characteristics**

Quality assessed in terms of oil content, acid, iodine and saponification values were significantly affected by 200 $\mu$ L/L ethrel spray. Spray effects under irrigated conditions did not differ significantly (Tables 165–166).

# ***CHAPTER–5***

## ***DISCUSSION***

## **CONTENTS**

## **DISCUSSION**

5.1	Introduction	100
5.2	Comparison of cultivars	101
5.3	Effects of ethrel, a ethylene releasing compound	103
5.3.1	Growth characteristics	103
5.3.2	Physiological characteristics	106
5.3.3	Biochemical characteristics	110
5.3.3.1	Nitrogen use efficiency	112
5.3.4	Yield characteristics	113
5.3.5	Quality characteristics	115
5.4	Conclusion	116
5.5	Future prospects	116

## DISCUSSION

## 5.1 Introduction

The efforts to increase land productivity have increased due to expanding food demand because of population explosion and decreasing land-man ratio. In post-independent India a new surge began in the shape of well known 'Green Revolution', which became meaningless after five decades due to non-sustainable food production for the ever-increasing population. The task of Green Revolution was achieved through the use of chemical fertilizers, <sup>as</sup> one of the several factors. With this chemical fertilizer tool, although food grain production in the country was achieved but its use also resulted in problems of soil salination<sup>is</sup>, ground water pollution, nutrient imbalance, emergence of new pest diseases and environmental degradation. The problem further aggravates by the constant rise in population, which is estimated that by 2020, India's population is likely to be around 1.3 billion. In addition to this, All India Yield Indices of Major Crops is found to be stagnating or decreasing due to stagnating yield and decreasing land-man ratio. This has exerted a massive pressure for managing the available resources in such a way as to be benefited maximally through environmentally sustainable techniques. Nitrogen is one of the mineral nutrient elements of prime importance in increasing crop productivity. It has a well-established role in the crop plants and is taken up in large quantity by plants. It is estimated that terrestrial plants assimilate approximately 1.4 gigatons of nitrogen annually and about 90 to 95% of the total in form of mineral nitrogen (Marschner, 1986). Although more input of nitrogen increases growth and yield of the crops, but its excessive use causes environmental degradation phenomenon like eutrophication of the resources.

Therefore, an approach is to be explored which minimizes the use of nitrogen without decreasing the growth and yield of crops. In this context, use of plant hormones may prove its potential as it has been found to enhance

growth and productivity of the crop plants (Leopold and Kriedmann, 1979; Khan, 1996a; Singh, 1996; Khan *et al.*, 1997; Khan *et al.*, 2000; 2001). Of several naturally occurring phytohormones, ethylene influences about all aspects of plant growth and development (Mattoo and White, 1991; Abeles *et al.*, 1992) as well as the induction of some plant defence responses (Boller, 1991). Ethylene produced in trace amount elicits many physiological responses, acting at a concentration as low as 0.01 $\mu$ L/L (Reid, 1987). Ethylene releasing compounds are applied to cereal crops to prevent lodging, thereby reducing yield losses and deterioration of grain quality (Foster *et al.*, 1991; Moes and Stobbe, 1991a, b), and to mustard for increasing yield (Khan, 1996b; Khan *et al.*, 2000).

Keeping in view the above facts, five field trials were conducted on mustard (*Brassica juncea* L. Czern & Coss.) on the following lines:

1. To study the effect of ethrel spray on growth, physiological, biochemical, yield and quality characteristics of mustard cultivars under irrigated and non-irrigated conditions.
2. To study the effect of ethrel spray on mustard grown with nitrogen levels under irrigated and non-irrigated conditions for nitrogen use efficiency and yield of the crop.
3. To test the hypothesis that ethylene has a central role in mediating plant responses with the use of ethylene action inhibitor silver thiosulphate

The results obtained in Chapter 4 are discussed below:

## 5.2 Comparison of Cultivars

In Experiment 1 and 2, Alankar and PBM16 cultivars of mustard (*Brassica juncea* L.) were used as experimental material. The rationale for use of the two cultivars is given in Materials and Methods section. The two cultivars responded similarly to ethrel application, giving maximal response with 200 $\mu$ L/L ethrel concentration. Both the cultivars showed inhibitory response to spray of ethrel higher than 200 $\mu$ L/L concentration.

For almost all growth characteristics, Alankar registered higher values than PBM16. Increase in inter-nodal length in Alankar was reflected in taller plants than PBM16. Leaf growth (area) in Alankar was more than PBM16 because of massive and irreversible expansion of daughter cell by meristematic divisions (Fig. 4). Also, Alankar showed higher inherent efficiency of retaining more moisture content in leaf tissues than PBM16. This was reflected in higher values for leaf fresh weight, leaf turgid weight and leaf relative water content (Tables 19, 47, 20, 48, 21, 49). Since leaf expansion is associated with trapping of solar energy and production of dry matter, therefore, Alankar registered higher dry weight (Fig. 6), and per cent distribution of dry weight towards leaf and pod (on emergence) was more in Alankar than PBM16. Thus, the efficiency of Alankar for translocating photoassimilates from leaf (source) to pod (sink) was greater than PBM16.

Alankar with large canopy structure, helped in efficiently receiving photosynthetically active radiation resulting in higher rate of net photosynthesis (Fig. 5) than PBM16. Higher leaf surface area as mentioned earlier was due to enhanced number of cell division and cell expansion. Increase in internal CO<sub>2</sub> concentration and stomatal conductance were reflected in more carboxylation efficiency, photosynthetic water use efficiency and water use efficiency in Alankar (Tables 26, 54, 27, 55, 28, 56).

The high efficiency of Alankar for conversion of biological matter to seed resulted in higher number of pods and seed yield (Fig. 7). The main contributing factor for increase in seed yield was pod number and seed number and 1000 seed weight was found almost equal in the two cultivars. Together with higher seed yield, oil yield of Alankar was more because of higher oil content than PBM16 (Tables 32, 60, 33, 61). It may be mentioned here that Alankar is the most suited cultivar grown in the region and surpassed other cultivars so far released. The cultivar was found equally suitable for irrigated and non-irrigated conditions.




Fig. 4. Effect of ethrel spray on leaf area of Alankar and PBM16 cultivars of mustard (*Brassica juncea* L.)






Fig. 5. Effect of ethrel spray on net photosynthetic rate of Alankar and PBM16 cultivars of mustard (*Brassica juncea* L.)




Fig. 6. Effect of ethrel spray on dry weight of Alankar and PBM16 cultivars of mustard (*Brassica juncea* L.)






Fig. 7. Effect of ethrel spray on seed yield of Alankar and PBM16 cultivars of mustard (*Brassica juncea* L.)

### 5.3 Effects of ethrel, a ethylene releasing compound

It is mentioned in Material and Methods section that Experiment 1 and 3 were conducted under irrigated conditions and Experiment 2 and 4 under non-irrigated conditions. The response of the crop to ethrel and nitrogen application was similar under these conditions of availability of water. This was possibly due to fact that mustard is a harder crop as moisture availability is concerned and may be said as drought tolerant. Moreover, monsoon showers during the month of December, when there is a higher need of moisture for reproductive parts formation, meet the water requirement. Therefore, the crop behaved similarly under irrigated and non-irrigated conditions, although with a difference in the absolute values for the plant characteristics. In the following pages, attempt has been made to discuss the results obtained in experiments with an assumption that similar phenomenon occurred due to the effect of ethrel or nitrogen under irrigated and non-irrigated conditions.

#### 5.3.1 Growth characteristics

Spray of 200 $\mu$ L/L ethrel increased growth characteristics maximally in Experiments 1-4. Further increase in ethrel concentration (400 to 800 $\mu$ L/L) proved inhibitory. Plant heights in Experiment 1 and 2 were increased by 14.9 and 16.2% respectively due to 200 $\mu$ L/L ethrel application. Such increasing effects of ethrel are basically on internode growth of plants (Sisler and Yang, 1984). Contrasting effects of ethylene sources on plant height have been reported in different plant species. For example, increased plant height in cauliflower (Jana and Kabir, 1991) due to ethrel application has been found. Contrarily, higher concentrations of ethrel/ethephon reduced plant height in sunflower (Sauerbrey *et al*, 1988), winter wheat (Van Sanford *et al.*, 1989), radish (Vreugdenhill and Harro, 1989), lupin (Ortuno *et al*, 1993), arabidopsis seedlings (Smalle *et al*, 1997), barley (Sanvicente *et al.*, 1999), linseed (Leitch and Kuat, 1999) and barley, oat and wheat (Rajala and Peltonen-Sainio, 2001). Wherever reduction of plant height was observed, this was due to use of higher

concentrations of ethrel. In this study also, higher concentrations of ethrel (400–800  $\mu\text{L/L}$ ) reduced plant height. These conflicting observations involved several mechanisms. Firstly, ethylene is found to promote the reorientation of cortical microtubules, thereby possibly controlling elongation (Shibaoka, 1994). Secondly, a role for ethylene has been suggested in regulating the expression of cell wall peroxidase involved in the control of the wall extensibility and cell growth (Ridge and Osborne, 1971). Regulating the levels of peroxidase activity by suitable concentration of ethrel possibly influenced the direction of growth of active tissues and organs. This effect of ethrel led to the emergence and formation of leaves with enhanced total leaf area of plant. Lower concentration of ethrel (200  $\mu\text{L/L}$ ) produced maximum leaf area, giving 27.0 and 35.7% increases over control in Experiment 1 and 2 respectively and higher concentration reduced the leaf area (Tables 8, 36, Fig. 4). Similar observation has been reported by Lee and Reid (1997) in sunflower. Ivenish and Kreibergs (1992), in cereal seedlings, reported that leaf emergence is associated with a peak of ethylene evolution. A similar phenomenon has been observed in *Arabidopsis* and burst of ethylene was accompanied by an increased expression of the ACC synthase gene 1, a gene suggested to be involved in the control of cell expansion (Rodrigues-Pousida *et al*, 1993). Thus, the induction of ethylene biosynthesis as in this may be associated with leaf emergence or in the control of cell expansion. This is also supported from the studies on ethylene-insensitive mutants, which have a large rosette than the wild type (Ecker, 1995) resulting from cell enlargement (Hua *et al*, 1995). Leaf fresh weight, leaf turgid weight, leaf relative water content and plant dry weight were significantly increased by 200  $\mu\text{L/L}$  ethrel (Tables 19, 47, 20, 48, 21, 49, 12, 40) because of observed increase in water use efficiency and photosynthetic water use efficiency in this ethrel treatment (Tables 27, 55, 28, 56). These aspects of efficiency parameters have been discussed in the following pages under a separate heading Physiological characteristics (Section 5.3.2).

It was found that the distribution of dry weight (on per cent basis) towards pods was higher in 200 $\mu$ L/L ethrel-sprayed plants. The dry weight thus accumulated was efficiently translocated to pod causing increase in per cent pod dry weight (Tables 18, 46). Linear regression analysis for various growth parameters with seed yield for Experiments 1–4 (Tables 167–178) also confirms the contribution of growth characteristics in yield.

Experiments 3 and 4 were conducted to assess the efficacy of foliar spray of ethrel on plants grown with nitrogen levels under irrigated (Experiment 3) and non-irrigated conditions (Experiment 4). Nitrogen is a major limiting nutrient for most plant species (Greenwood, 1982). Acquisition and assimilation of nitrogen is second in importance only to photosynthetic carbon assimilation for plant growth and development (Heickel, 1980; Araus *et al.*, 1993; Anten *et al.*, 1995; Arthamawar *et al.*, 1996). Non-availability of nitrogen in Experiment 3 and 4 affected growth characteristics (Tables 72–86, 105–119). The best dose of nitrogen was 80kg N/ha and nitrogen level lower than this showed poor growth. At low (0 to 60kg N/ha) nitrogen level, the N absorbed by the roots was utilized for protein synthesis from reserve root carbohydrates and supply of N to the top of plants was limited affecting the growth of the shoot. The lesser N content present in the plant (Tables 98, 131, 99, 132) also show that at lower levels of basally applied N, the plant growth was poor and not benefited much.

In this study the growth response of the plant was maximum with application of 80kg N/ha together with spray of 200 $\mu$ L/L ethrel. The values obtained for this combination were maximum and higher than any other combinations of nitrogen and ethrel concentration (Fig. 8–15). The positive effect of 200 $\mu$ L/L ethrel spray together with 80kg N/ha is attributed to the changes in ethylene evolution (discussed in section 5.3.2). The ethylene evolution is reported to be increased by excessive ammonia accumulation (Corey *et al.*, 1987; Arshad and Frenkenberger, 1991) and can be induced by




Fig. 8. Effect of ethrel spray on leaf area of cultivar Alankar of mustard (*Brassica juncea* L.) grown with different levels of nitrogen






Fig. 9. Effect of ethrel spray on net photosynthetic rate of cultivar Alankar of mustard (*Brassica juncea* L.) grown with different levels of nitrogen






Fig. 10. Effect of ethrel spray on dry weight of cultivar Alankar of mustard (*Brassica juncea* L.) grown with different levels of nitrogen




Fig. 11. Effect of ethrel spray on 1-aminocyclopropane-1-carboxylic acid (ACC) of cultivar Alankar of mustard (*Brassica juncea* L.) grown with different levels of nitrogen






Fig. 13. Effect of ethrel spray on ethylene evolution of cultivar Alankar of mustard (*Brassica juncea* L.) grown with different levels of nitrogen

urea fertilization (Barker and Cory, 1990). Thus, the suitable N availability led to tissue ammonia accumulation and increased ethylene evolution, which triggered many physiological effects and enhanced growth of plants. The obtained dry mass increase in plant (Fig. 10) and its higher distribution in leaf than in stem and pod were because of better management of available photosynthates and source-sink relation. These results were observed in the form of per cent leaf, stem and pod dry weights (Tables 81, 114, 82, 115, 83, 116), specific leaf weight (Tables 76, 109) and enhanced leaf area (Tables 73, 106).

The contribution of leaf dry mass to total dry mass was found to be relatively higher for nitrogen dose 80kg N/ha and sprayed with 200 $\mu$ L/L ethrel than any other treatments. As the growth progressed pod dry mass increased. This was due to remobilization of dry matter from source (leaves) to sink (pods) under the influence of balanced nutrient regime. The balanced nutrient profile helped in maintaining maximum growth characteristics. Together with this, spraying of ethrel also helped the plant to produce signals for better utilization of nutrients (Graan and Boyer, 1990; Zhang and Davies, 1990; Davies and Zhang, 1991; Nilsen and Orcutt, 1996).

The altered structure of canopies in relation to leaf size and leaf area index were helpful in improving solar energy harvesting ability of the leaves as evident from increased plant dry matter production (Tables 77, 110; Fig. 10). Earlier reports from the author's laboratory have shown that spray of 200 $\mu$ L/L of ethrel on mustard plants enhanced dry matter through increase in leaf area index (Khan, 1998; Khan *et al.*, 2000) of mustard grown under non-irrigated conditions. These studies were, however, limited to the use of only a single ethrel concentration and few plant characteristics were studied.

### 5.3.2 Physiological characteristics

A variety of plant physiological processes at the biochemical and whole plant level are the driving force behind biomass production. Canopy

photosynthesis, integrated over a whole growing season, and whole plant pattern of photoassimilates partitioning is more important physiological determinant of biomass production and crop yield. Rate of photosynthesis is responsive to number of factors, like canopy structure, interception of solar radiation, stomatal conductance, carbon dioxide concentration and levels of ethylene. The degree to which photosynthesis responds to exogenous levels of ethylene is not well understood.

Rates of photosynthesis were increased by 23.7, 27.2, 27.7 and 31.3% with 200 $\mu$ L/L ethrel over control in Experiments 1, 2, 3 and 4 respectively. This ethrel concentration has shown to increase leaf area and leaf area index (discussed in Growth characteristics section). Further, ethrel induced leaf area showed a strong correlation with ethrel-induced photosynthesis (Fig. 21A). Ethrel-induced photosynthesis exhibited strong relationship with ethrel-induced carboxylation efficiency (Fig. 21B). Observations of Buehler *et al.* (1978); Grewal and Kolar (1990) and Grewal *et al.* (1993) have shown the increase in photosynthesis with ethrel due to increase in chlorophyll per unit of leaf area. The increase in photosynthesis with ethrel has also been reported by Subrahmanyam and Rathore (1992a), Pua and Chi (1993) and Khan *et al.* (2000). Retaining higher leaf area index in ethrel-treated plants helped in an increase in photosynthesis. However, at 100d, inspite of increase in leaf area, the photosynthesis rate declined because of mutual shading of leaves. Ethrel concentrations higher than 200 $\mu$ L/L in Experiments 1 and 2 decreased the photosynthesis. Inhibition of photosynthesis in these experiments was because of higher amount of ethylene evolution due to 400–600 $\mu$ L/L ethrel application. Such conditions of inhibition of photosynthesis by ethylene have also been suggested by Kays and Pallas Jr (1980) and Rajala and Peltonen-Sainio (2001). This is possible that threshold value for ethylene with 200 $\mu$ L/L ethrel was comparable to that which elicits the ethylene induced hormonal responses.

In the reported observation, a correlation between ethrel-induced photosynthesis and stomatal conductance (Fig. 22B) suggests that differences in stomatal conductance contributed significantly to the variation in photosynthesis. Analysis of  $A/C_i$  values (carboxylation efficiency) also suggests non-stomatal limitation to photosynthesis. The correlation studies showed that variation in photosynthetic capacity (carboxylation efficiency) accounted for the differences in photosynthetic rate (Fig. 21B) more so than differences in stomatal limitation. This view is again strengthened by the observed relationship of carboxylation efficiency with photosynthetic water use efficiency (Fig. 22A). Vanden Boogard *et al.* (1995) reported a correlation between photosynthetic water use efficiency with rubisco activity and photosynthetic water use efficiency as a measure of rubisco activity. Water use efficiency measured from the data on biomass and transpiration rate showed that it was increased maximally with 200  $\mu\text{L/L}$  ethrel. The effect of ethrel on water use efficiency was through maintenance of turgor potential and stomatal movement. At maturity, high water use efficiency was caused by lower transpiration rate associated with a high leaf area per unit plant weight, also observed by Vanden Boogard (1996a, b). Similarly, nutrients have also been shown to have a role in water use efficiency of plants (Payne *et al.*, 1992; Bruck *et al.*, 2000). Ethylene-enhanced net photosynthetic rate helped in an increase in plant dry weight, therefore photosynthetic water use efficiency showed relationship with plant dry weight (Fig. 23A–B).

Contrasting reports on exogenously applied ethylene sources on transpiration rate appear in the literature. Transpiration is reportedly non responsive to ethylene in both herbaceous and woody species (Aharnoi, 1978; Pallaghy and Raichke, 1972; Johnson, 1984). In contrast, significant responses in transpiration are shown in several herbaceous species (Govindarajan and Pooviah, 1982; Kays and Pallas Jr, 1980; Pallas Jr and Kays, 1982). Increase in carboxylation efficiency and photosynthesis was related to the increase in

intercellular carbon dioxide concentration by ethrel. At later maturity stages decrease in photosynthesis was related to resistance to CO<sub>2</sub> diffusion and was stomatal limitation. Moreover, Mattoo and White (1991) reported that CO<sub>2</sub> could promote or inhibit ethylene evolution, depending on the tissue concentration. On the same lines Dharan *et al.* (1981), Kao and Yang (1982) and Grodzinski *et al.* (1982) reasoned that inhibition of ethylene evolution resulted from a decrease in internal CO<sub>2</sub> concentration and regulated photosynthesis. In this study, 200µL/L ethrel significantly favoured ethylene evolution that fundamentally influenced the central regulatory system. Further, increase in ethrel concentration (400 or 600µL/L) possibly promoted ethylene evolution to inhibit the physiological and biochemical processes. It seems probable that there is some requirement of ethylene for optimum response. Low and high concentrations represent the two ends of an optimum curve, like that for many hormones, promoting at low concentration and inhibiting at high. There is, thus, an interrelation between CO<sub>2</sub> metabolism and ethylene evolution, which controls other biochemical and physiological changes. Ethylene evolution was linked with activity of ACC oxidase. It was possible that stimulatory effect of ethrel on ACC oxidase might involved not only on enhancement of the activity but also its synthesis and degradation, thereby, resulting in higher ACC dependent ethylene evolution. It may be emphasized that ethrel application promoted ACC dependent ethylene evolution (Tables 96, 129).

Mattoo *et al.* (1997) observed that autocatalytic effect of ethylene was physiologically achieved by an enhancement in the activities of both ACC synthase and ACC oxidase. It may be reiterated that the high conversion rate of ACC to ethylene permitted to influence the fundamental control system. The concentration of ethylene with spray of 200µL/L ethrel showed most suitable concentration for plant growth and development. The concentration of ethylene

with 100 $\mu$ L/L ethrel remained below a critical threshold and higher concentration than 200 $\mu$ L/L possibly exceeded threshold values of ethylene.

In Experiment 5, application of 200 $\mu$ L/L ethrel was used to promote ethylene evolution together with 1mM of silver thiosulphate (STS) to block the action of ethylene. The objective was to test the hypothesis that ethylene has a central role in mediating plant responses, reported in Experiment 1–4.

The hypothesis that ethylene was responsible for the observed enhancement in plant characteristics with 200 $\mu$ L/L ethrel was substantiated by this experiment (Experiment 5) where ethrel and STS were supplied.

It was discussed in preceding pages that 200 $\mu$ L/L ethrel increased ethylene evolution to optimum concentration causing maximum increase in growth and physiological characteristics. Limiting the action of ethylene with STS reduced the plant characteristics. Thus, a reduction of growth and physiological rates was achieved by blocking the ethylene action (Tables 147–166). As STS-treated plants exhibited values lower than the plants treated with water (control), it is possible that the intrinsic ethylene biosynthesis was affected (Fig. 16–18).

### 5.3.3 Biochemical characteristics

Nitrogen related characteristics were studied in Experiment 3 and 4. Nitrogen status of a plant in a particular spatial or temporal zone exhibits the nutritional requirement of the crop during that period in that part of plant phase. In different water availability regime, the nutrient availability differed, and plant respond to the changing environment by proliferating the roots, which develop deep into the soil layers in search of nutrients and water. Thus, the amounts of nitrogen that can be transported to the shoots depend on the capacity of the roots to absorb nitrogen from the soil and transport them to the transpirational stream. The concentration of nitrogen in growth media exerts a considerable influence on growth and mineral composition of crop plants (Kurvitis and Kirkby, 1980; Gashew and Mugwira, 1981; Ansari, 1990;






Fig. 16. Effect of ethrel spray (E: 0 or 200  $\mu\text{L/L}$ ) or silver thiosulphate (S: 1mM) on leaf area of Alankar cultivar of mustard (*Brassica juncea* L.)




Fig. 17. Effect of ethrel spray (E: 0 or 200 μL/L) or silver thiosulphate (S: 1mM) on net photosynthetic rate of Alankar cultivar of mustard (*Brassica juncea* L.)




Fig. 18. Effect of ethrel spray (E: 0 or 200  $\mu\text{L/L}$ ) or silver thiosulphate (S: 1mM) on dry weight of Alankar cultivar of mustard (*Brassica juncea* L.)




Fig. 19. Effect of ethrel spray (E: 0 or 200  $\mu\text{L/L}$ ) or silver thiosulphate (S: 1mM) on seed yield of Alankar cultivar of mustard (*Brassica juncea* L.)

Jeschke *et al.*, 1992) and affects the relation uptake of cations and anions by plants (Kirkby, 1981; Lovatt, 1986).

In the present study, nitrogen content was found to increase with increasing doses of nitrogen. At initial growth stage, nitrogen content increased by 8.6 and 11.5% in Experiment 3 and Experiment 4 respectively with 80kg N/ha applied as compared to 0kg N/ha. The increase in the nitrogen content at early growth stage coincides with the increasing rate of nitrate reductase activity (Tables 97, 130). Nitrogen accumulation was found to be well coordinated with supply of nitrogen. It is believed that availability of a given nutrient may interact with the uptake of other nutrients and thus making the uptake more complex (Amoruwa *et al.*, 1987; Marschner, 1986). The response of accumulation to nitrogen deprivation (0kg N/ha) was negligible for nitrogen content at 80d sampling. The possible explanations are that there was a tendency for the root system to have more proliferation in the soil to meet the plant nutrient demand or it became larger relative to the shoot during nitrogen deprivation. Ethylene regulated differentiation in trachied (Zobel and Roberts, 1978), root elongation (Koning and Jackson, 1979), adventitious root formation (Liu *et al.*, 1990) and root/shoot ratio (Rajala and Peltonen-Sainio, 2001) have been observed. In a study on mustard, Khan *et al.* (1996a) reported increase in nitrate reductase activity with GA<sub>3</sub> application. Enhancement of nitrate reductase activity by GA<sub>3</sub> or GA<sub>3</sub> + cytokinin in tobacco leaves has been observed by Roth-Bejerano and Lips (1970). The accumulation of nitrogen was associated with increased photosynthetic rate, which resulted in higher dry matter accumulation. Together with this other possibility is of accumulation of other cations (preferably potassium), which helped in maintaining the rate of photosynthesis by improving the relative water content of the leaf through osmotic adjustment. In a study on mustard, Khan *et al.* (2000) reported that potassium accumulation increased with nitrogen supply, which caused increase in stomatal conductance, photosynthetic rate and dry matter accumulation.



Regarding this trial, it has been reported that this was a preliminary trial. Accumulation of potassium in guard cells provides the necessary amount of solute for developing the water potential gradient required for water movement into the guard cells for stomatal opening necessary for photosynthesis (Jensen and Tophoj, 1985; Tanguilag *et al.*, 1987 and Thakral *et al.*, 1997). The findings also encourage the view that there is some form of co-regulating of the nutrients accumulation in mustard, which may akin to that described for their accumulation (Vyas *et al.*, 1995; Khan *et al.*, 1997; Zaman and Choudhri, 1998).

#### 5.3.3.1 Nitrogen use efficiency

Nitrogen use efficiency has been defined as seed yield with per unit of available N (soil N + fertilizer N). It has two components, nitrogen uptake efficiency (plant N per unit of soil + fertilizer N) and nitrogen utilization efficiency (seed yield per unit of N in plant). A product of these two components results in nitrogen use efficiency (Moll *et al.*, 1982; Prasad *et al.*, 2000). In the present research, maximal use efficiency of N was found with 200 $\mu$ L/L ethrel and 80kg N/ha (Tables 179–184; Fig. 20). Nitrogen applied in sub-optimal level (40 or 60kg N/ha), the use efficiency was less. At optimal N (80kg N/ha) use efficiency was higher and application of 200 $\mu$ L/L ethrel on plants receiving 80kg N/ha enhanced use efficiency further. It may be mentioned here that treatment 80kg N/ha and 200 $\mu$ L/L ethrel enhanced top growth of plants maximally, which puts demand on soil N to meet the growing need of the shoot. Increased nitrate reductase activity in leaves with 80kg N/ha and ethrel (200 $\mu$ L/L) also supports the incorporation of nitrogen in the shoot. Moreover, plant N was translocated to seed N reflecting more of its translocation on application of ethrel finally increasing the seed yield. Moreover, nitrogen application results in enhanced ethylene evolution, which is reported to increase by excessive ammonia accumulation (Corey *et al.*, 1987; Arshad and Frenkenberger, 1991) and can be induced by urea fertilization




Fig. 20. Effect of ethrel spray on N uptake, N use and N utilization efficiency of cultivar Alankar of mustard (*Brassica juncea* L.) grown with different levels of nitrogen




Fig. 21. The relationship between (A) ethrel induced changes in leaf area and ethrel induced changes in net photosynthetic rate ( $P_N$ ) and (B) ethrel induced changes in carboxylation efficiency (CE) and ethrel induced changes in net photosynthetic rate ( $P_N$ ). The per cent increase in values from 200  $\mu\text{L/L}$  ethrel over control of the cultivars PBM16 and Alankar in Experiments 1 and 2 were used for these correlation studies.  $^{**}$  significant at  $P = 0.01$ .






Fig. 22. The relationship between (A) ethrel induced changes in photosynthetic water use efficiency (PWUE) and ethrel induced changes in carboxylation efficiency (CE) and (B) ethrel induced changes in stomatal conductance ( $g_s$ ) and ethrel induced changes in net photosynthetic rate ( $P_N$ ). The per cent increase in values from 200  $\mu\text{L/L}$  ethrel over control of the cultivars PBM16 and Alankar in Experiments 1 and 2 were used for these correlation studies. \* significant at  $P = 0.05$ , \*\*  $P = 0.01$ .




Fig. 23. The relationship between (A) ethrel induced changes in dry mass (DM) and ethrel induced changes in net photosynthetic rate ( $P_N$ ) and (B) ethrel induced changes in dry mass (DM) and ethrel induced changes in photosynthetic water use efficiency (PWUE). The per cent increase in values from 200  $\mu\text{L/L}$  ethrel over control of the cultivars PBM16 and Alankar in Experiments 1 and 2 were used for these correlation studies.  $^{**}$  significant at  $P = 0.01$ .

(Barker and Corey, 1990). Availability of N leads to tissue ammonia accumulation and increased ethylene formation (Feng and Barker, 1992, 1993). Nitrogen use efficiency with fertilizer N has also been reported by Gajri *et al.* (1993), Feiz *et al.* (1994), Gardner *et al.* (1994) and Sowers *et al.* (1994). The effect of GA<sub>3</sub> on the nitrogen use efficiency has been reported from the author's laboratory but the effect of ethrel on nitrogen use efficiency has not been reported earlier

#### 5.3.4 Yield characteristics

Seed yield of a crop may depend on the vegetative growth of the crop because photosynthesizing sites have determinant role in producing the photosynthates. Yield is the final manifestation of several intricate morphological and physiological traits that initiate at germination and terminates at harvest. Yield is dependent on the maintenance of an array of metabolic processes including photosynthesis and hormonal status. There is increasing evidence for metabolic growth regulator effects on various crops.

In fact, leaf size constitutes the canopy structure and is actively involved in interception of solar radiation and in contribution of photoassimilates to the developing pods. Two sequential steps are necessary for a mustard plant to produce pods, a sink of pollinated pods capable of further development must be created and this must be supplied with photosynthates over the subsequent period of development. Thus, seed yield at harvest may be determined either by the seed capacity established at pollination or by the quantity of photosynthate made available between pollination and maturity. There is a positive feed back cycle between photosynthetic products, growth and leaf area. The effectiveness of this cycle is increased under favourable conditions. The CO<sub>2</sub> enrichment through increase in leaf canopy may increase plant growth by stimulating photosynthetic rate and thereby accelerating the cycle (Rogers *et al.*, 1996).

The aim of the Experiments 1 and 2 was find out suitable concentration of ethrel spray on mustard and thereby, if possible to increase seed yield. It was

found that 200 $\mu$ L/L ethrel spray contributed for enhancement of pod number by 11.9 and 14.6 % over control in Experiment 1 and 2 respectively (Tables 31, 59). Primarily, the pod number enhancement contributed to increase of 32.3 and 40.5% seed yield in these experiments. Moreover, increase in number of seeds per pod (14.5% in Experiment 1 and 11.8% in Experiment 2) and slight increase in 1000 seed weight also helped in increasing the seed yield. Increase in seed yield resulted in an increase of 38.8 and 48.3% oil yield in Experiment 1 and Experiment 2 respectively (Tables 32, 60, 33, 61).

In Experiment 3 and 4, it was found that there was differential response of 200 $\mu$ L/L ethrel spray for the yield when the plants were grown with nitrogen deficient (0kg N/ha) to sufficient nitrogen (80kg N/ha). This combination of treatments improved the seed yield by increasing the proportion of the reproductive tissues (inflorescence and/or pods) to total plant dry matter (biological yield). There was 8.9 and 10.5% increase in pod number and 21.0 and 26.4% increase in seed yield in Experiments 3 and 4 respectively (Tables 100, 133, 101, 134). The increase in yield was directly related to the increase in sink capacity. Ethrel application enhanced the incorporation of soil nitrogen (Section 5.3.3), which was manifested in the better vegetative growth (Section 5.3.1) and better partitioning into reproductive parts, evident from increased pod number and harvest index. Linear regression analysis for growth and yield attributing characteristics with seed yield in Experiment 1–4 also confirms the view (Tables 167–178). The beneficial effects of ethrel have also been studied by many research scientists including Dahnous *et al.* (1982), Leary and Oplinger (1983), Wiersma *et al.* (1986), Joshi *et al.* (1987), Singh *et al.* (1987), Ramos *et al.* (1989), Singh and Kumar (1991), Bulman and Smith (1993a, b), Grewal *et al.* (1993) and Kasele *et al.* (1995) in several crop species.

Nitrogen yield potential provides a measure of nitrogen partitioning potential from vegetative parts to seeds. In Experiment 3 and 4, nitrogen yield potential was enhanced by 32.3 and 35.4% respectively due to 200 $\mu$ L/L ethrel

spray (Tables 102, 135). Increase in seed yield has also resulted in increase in oil yield by 30.6 and 36.7% in Experiments 3 and 4 respectively.

### 5.3.5 Quality characteristics

For assessing oil quality, oil content, acid, iodine and saponification values of oil were determined. It may be added that low acid and iodine values are considered good for oil quality and denote good keeping and easy hydrogenation. High saponification value is good for digestibility and soap making quality.

Ethrel application increased the oil content. The ethrel (ethylene) might have increased the supply carbon to lipid synthesis by induction of a specific transporter and played a role in determining the amount of oil. There is possibility of ethrel (ethylene) affecting acetyl CoA carboxylase. Other workers have also shown that an additional carbon input to lipid synthesis contribute to the oil formation (Post-Beittenmiller *et al.*, 1992; Hunter and Ohlrogge, 1998; Kozaki and Sasaki, 1999; Sawage and Ohlrogge, 1999). Leitch and Kuat (1999) demonstrated role of ethylene applied as ethrel in fatty acid synthesis, increasing the oleic acid in linseed oil. In the present study increase in iodine and saponification values indirectly provide evidence of increase in short chain unsaturated fatty acid. However, the possibility of ethylene affecting acetyl CoA carboxylase and to the flux of carbon to oil in developing embryos of oilseed modifying fatty acid composition needs to be investigated.

Application of nitrogen had not any significant effect in altering oil content of the seed in the present study. However, reports are available for reduced oil content due to N fertilization (Smith *et al.*, 1988; Gendy and Marquard, 1989; Khan *et al.*, 1990; Pinkerton, 1991; Samiullah *et al.*, 1991; Asare and Scarisbrick, 1995). In spite of non-significant effect on oil content, oil yield was significantly enhanced by ethrel (200 $\mu$ L/L) and 80kg N/ha due to increase in seed yield.

## 5.4 Conclusion

This has been shown in the study, reported in the thesis, that manipulating the quantity of ethylene alters the leaf expansion and total plant leaf area. Specifically, low concentration of ethylene releasing compound ethrel (200 $\mu$ L/L) stimulated leaf area and higher concentration of ethrel reduced the leaf area. Other effects of ethylene were documented in stomatal and non-stomatal limitations to carboxylation and photosynthesis, which resulted in higher biomass production. In experiments conducted with doses of nitrogen and ethrel, the stimulatory effects of ethylene on nitrate reductase activity and plant nitrogen use efficiency were seen. Enhanced plant top growth due to ethrel (200 $\mu$ L/L) exerted demand on the root for higher utilization of available soil-N.

Nitrogen use efficiency became high and nitrogen accumulation enhanced in the plant above ground part. Nitrogen was incorporated into macromolecules and reflected in higher dry matter accumulation. Efficient partitioning of dry matter resulted in increased seed yield and seed yield potential of the plants.

## 5.5 Future prospects

The role of plant hormones in growth and development is well known. However, the action of hormones is brought about by producing its effect on other hormones and also controlling the biosynthesis of other hormones. For example, it has been shown that ethylene inhibits GA induced elongation and GA reverses the effects of ethylene. Similarly, the action of ABA inhibits endogenous production of ethylene, but not that formed from ACC and regulates stomatal movements. ABA inhibits the formation of ethylene that is induced by IAA. These aspects of hormonal control over the action of ethylene need to be investigated. It is proposed to work out the level of GA and ABA to have in depth understanding on the regulatory role of ethylene in photosynthesis and biomass accumulation.

# ***CHAPTER-6***

## ***SUMMARY***

### **SUMMARY**

The thesis “Physiological significance of ethrel (2-chloroethyl phosphonic acid) and nitrogen in relation to growth and metabolism of mustard under irrigated and non-irrigated conditions”, contains six chapters.

Chapter 1 deals with the importance of the problem. Lacunae in the understanding of the problem and justifications for undertaking the present study have been put forth.

Chapter 2 is review of literature. Relevant available literature pertaining to individual as well as combined effect of plant growth regulators especially ethylene sources with nitrogen on crop growth and development has been given in this chapter

Chapter 3 describes the details of the material used in the study and methods employed in determining observations carried out in the experiments. Relevant information on meteorological and edaphic data has been included.

In Chapter 4, the results obtained in the experiments which were found significant at  $P < 0.05$  have been recorded in detail.

In Chapter 5, significant results have been discussed in the light of earlier reported findings. Possible explanations of the data obtained have also been given to reach a conclusion. The results of the five field experiments are summarized below.

Experiment 1 (1998-99) was conducted under irrigated conditions to study the response of two cultivars (Alankar and PBM16) of mustard (*Brassica juncea* L.) to leaf-applied ethrel at 60d after sowing (flowering stage). Alankar is a well established cultivar and accepted by the mustard growers for a decade, whereas PBM16 is a newly released. This was a factorial experiment conducted according to randomized complete block design. The response of the cultivars to ethrel treatments was assessed by determining growth and biochemical characteristics at 80 (pod fill), 100 (pod maturity) and 120d (harvest) after



sowing. Physiological characteristics were studied at 80 and 100d after sowing. Yield and quality characteristics were determined at harvest. Growth characteristics were plant height, plant leaf area, leaf area index, specific leaf area, specific leaf weight, plant dry weight and dry weights and per cent dry weight distribution in leaf, stem and pod, leaf fresh weight, leaf turgid weight and leaf relative water content. Physiological characteristics were, rate of photosynthesis, stomatal conductance, internal CO<sub>2</sub> concentration, transpiration rate, carboxylation efficiency, photosynthetic water use efficiency, plant water use efficiency. Biochemical characteristics included N content and N accumulation in plant. At harvest, yield characteristics studied were, number of pods per plant, number of seeds per pod, 1000 seed weight, seed yield, biological yield, harvest index and oil yield. The oil was assessed for acid, iodine and saponification values.

Among five concentrations of ethrel (0, 100, 200, 400 and 600µL/L) applied, 200µL/L was found superior over others in increasing the plant characteristics studied. The concentration less than 200µL/L was found less effective, whereas concentration higher than 200µL/L proved inhibitory. Spray of 200µL/L ethrel affected growth of the plants and increased total leaf area per plant. Thus, the increase in photosynthesising surface area led to increase in photosynthesis and CO<sub>2</sub> accumulation, which resulted in increased plant dry weight. The other effects of ethrel spray was noted in improved source-sink relationship, as seen in increase in per cent pod dry mass in ethrel-treated plants. More of the flowers developed into pods, evident from higher pod number and seed yield in ethrel-treated plants compared with water-sprayed control. The oil quality was also improved with 200µL/L ethrel spray.

Comparison of the cultivars showed that Alankar established its superiority over PBM16. Growth, physiological, biochemical, yield and quality characteristics were found superior in Alankar than in PBM16. Interaction effect between cultivar and ethrel spray was non-significant for most of the

plant characteristics. This suggests that the two cultivars responded similarly to ethrel spray.

Experiment 2 (1998-99) was a factorial randomized complete block design conducted on the same lines as Experiment 1 but under non-irrigated conditions. The scheme of the treatments, design of the experiment, ethrel spray treatments and cultivars of mustard were also similar as described for Experiment 1. The observations recorded at different sampling stages were similar to Experiment 1. In this experiment, it was noted that ethrel spray at 200 $\mu$ L/L concentration was more effective than any other concentrations used. The effect of the spray was manifested through changes in various characteristics as described for Experiment 1. Alankar cultivar surpassed PBM16 in performance.

Combined analysis of the two experiments showed that the factors irrigated and non-irrigated were non-significant. Ethrel spray under irrigated and non-irrigated conditions was equally effective. The two cultivars also behaved similarly in the two conditions of irrigation.

Experiment 3 (1999-2000) was a factorial randomized complete block design conducted under irrigated conditions to study the effect of leaf-applied 0, 100 and 200 $\mu$ L/L ethrel on the performance of Alankar cultivar of mustard (the cultivar was selected on the basis of Experiment 1) grown with soil-applied 0, 40, 60 and 80kg N/ha. Ethrel spray application was done at 60d after sowing (flowering stage) and performance of the crop was assessed by determining various plant characteristics at 80 (pod fill), 100 (pod maturity) and 120d (harvest) after sowing. Growth characteristics were those studied in Experiment 1. Among physiological characteristics, 1-aminocyclopropane-1-carboxylic acid content, ACC oxidase and ethylene evolution were also studied in addition to the characteristics studied in Experiment 1. Among biochemical characteristics, leaf nitrate reductase activity was also studied in addition to the determination of N content and N accumulation in plant. Among yield

characteristics, seed N, nitrogen harvest index and nitrogen yield potential were also studied in addition to the yield characteristics studied in Experiment 1. Quality characteristics were similar as in Experiment 1.

Ethrel at 200 $\mu$ L/L concentration and nitrogen at 80kg N/ha registered significantly superior values as compared to other treatments. Ethrel (200 $\mu$ L/L) enhanced growth, physiological, biochemical, yield and quality characteristics. In this experiment, it was found that the effect of 200 $\mu$ L/L ethrel was maximal when plants received soil-applied 80kg N/ha. This combination (200 $\mu$ L/L ethrel and 80kg N/ha) enhanced plant leaf area, photosynthesis, CO<sub>2</sub> accumulation and plant dry weight. Water relations characteristics such as leaf fresh weight, leaf turgid weight and leaf relative water content were also enhanced by 200 $\mu$ L/L ethrel x 80kg N/ha. Pod dry weight was maximal with 200 $\mu$ L/L ethrel x 80kg N/ha, which showed higher translocation of dry matter towards sink (pods). Among physiological characteristics, rate of photosynthesis, internal CO<sub>2</sub> concentration, transpiration rate, photosynthetic water use efficiency, water use efficiency, 1-aminocyclopropane-1-carboxylic acid content and ACC oxidase were maximal in 200 $\mu$ L/L ethrel x 80kg N/ha. Plants grown with sufficient soil-applied N (80kg N/ha) responded to the ethrel (200 $\mu$ L/L). However, if soil-applied N was less than 80kg N/ha, the effect of ethrel spray was not prominent. The increased vegetative growth due to ethrel (200 $\mu$ L/L) spray made the plant to extract more of the soil N and was reflected in N content and accumulation and increased growth, physiological, biochemical and yield characteristics. The calculated nitrogen uptake efficiency, nitrogen utilization efficiency and nitrogen-use efficiency showed that nitrogen use was better when plants were treated with the 200 $\mu$ L/L ethrel spray. Seed yield, oil yield and nitrogen yield potential were increased and were maximal with 200 $\mu$ L/L ethrel x 80kg N/ha.

Experiment 4 (1999-2000) was a factorial randomized complete block design conducted simultaneously with the Experiment 3 under non-irrigated

conditions. The scheme of the treatments, ethrel spray concentrations and soil-applied nitrogen were same as in Experiment 3. The data on growth, physiological, biochemical, yield and quality characteristics recorded were those mentioned for Experiment 3. Individual effects of 200 $\mu$ L/L ethrel spray, soil-applied 80kg N/ha and their interaction proved best for most of the plant characteristics. The response of the plant to ethrel and nitrogen treatments was similar as found in Experiment 3.

Combined analysis of Experiments 3 and 4 showed that the response of the plant to ethrel spray, soil-applied nitrogen and their interaction was uniform irrespective of the irrigation conditions. The data on plant characteristics under irrigated and non-irrigated conditions were non-significant. The data suggests that 200 $\mu$ L/L ethrel spray on plants grown with soil-applied 80kg N/ha may be used for improving mustard cultivation irrespective of the conditions of irrigation.

Experiment 5 (2000-2001) was a factorial conducted according to randomized complete block design. In this experiment applications of 0 and 200 $\mu$ L/L ethrel or 1mM silver thiosulphate were done as foliar spray at 60d after sowing (flowering stage) on mustard cultivar Alankar grown under irrigated and non-irrigated conditions. Plants were raised with uniform soil application of 80kg N/ha. This experiment was based on the findings of Experiments 3 and 4. The response of the plants to ethrel spray treatment was confirmed in this experiment with the use of silver thiosulphate spray treatment, as silver thiosulphate application inhibits ethylene action. The observations recorded at 80, 100 and 120d after sowing included growth (plant leaf area plant dry weight), physiological, biochemical, yield and quality characteristics were similar to Experiments 3 and 4. Maximum response was noted with 200 $\mu$ L/L ethrel spray treatment. However, silver thiosulphate spray inhibited the ethylene action and ethrel effect was not observed. The interaction effect of spray and irrigation was found non-significant. The results of the

experiments suggest that response of the plant to ethrel treatment seen in Experiment 1–4 was manifested through the action of ethylene.

The present chapter is followed by bibliography of the literature cited in the thesis. Appendix is given in the last to show the preparation of various chemicals used in the study.

# ***BIBLIOGRAPHY***

## BIBLIOGRAPHY

---

1. Abeles, F.B.; Morgan, P.W. and Saltveit Jr., M.E. 1992. Ethylene in plant biology. 2<sup>nd</sup> ed. Academic Press, San Diego.
2. Ado-Quaye, A.A.; Scarisbrick, D.H. and Daniel, R.W. 1986. Assimilation and distribution of <sup>14</sup>C photosynthate in oilseed rape (*Brassica napus*). *Field Crops Research* **13**: 205-215.
3. Afridi, M.M.R.K.; Wasiuddin; Afaq, S.H.; Samiullah and Parvaiz, M.A. 1983. Effect of nitrogen, phosphorus and potassium on the growth and yield of fennel (*Foeniculum vulgare* Mill.). *Agricultural Science Progress* **1**: 63-73.
4. Aharoni, N. 1978. Relationship between leaf water status and endogenous ethylene in detached leaves. *Plant Physiology* **61**: 658-662.
5. Al Jaloud Ali, A.; Hussain, G.; Karimullah, S. and Al Hamidi, A.H. 1996. Effect of irrigation and nitrogen on yield and yield components of two rapeseed cultivars. *Agricultural Water Management* **30**: 57-68.
6. Allen, E.J. and Morgan, D.J. 1972. A quantitative analysis of the effects of nitrogen on growth, development and yield of oilseed rape. *Journal of Agricultural Science Cambridge* **78**: 315-324.
7. Amanico, S. and Santosh, H. 1992. Nitrate and ammonium assimilation by root of maize (*Zea mays* L.) seedlings as investigated by *in vivo* <sup>15</sup>N-NMR. *Journal of Experimental Botany* **43**: 633-639.
8. Amoruwa, G.M.; Ogunlela, V.B. and Ologunde, O.O. 1987. Agronomic performance and nutrient concentration of maize (*Zea mays* L.) as influenced by nitrogen fertilization and plant density. *Journal of Agronomy and Crop Science* **159**: 226-231.
9. Anonymous. 1970. Pharmacopocia of India. Manager of Publication, Ministry of Health, Government of India, New Delhi.
10. Anonymous. 1993. Food and nutrition. In: *Manorma Year Book*. pp. 137-140. M. Mathen (Ed). Malayala Manorma, Kottayama, Kerala, India.
11. Ansari, H.R. 1996. Effect of some phytohormones and NPK on growth and metabolism of mustard. Ph.D. Thesis, Aligarh Muslim University, Aligarh, India.

12. Ansari, R. 1990. Growth and chemical composition of barley (*Hordeum vulgare*) cultivars on saline substrate as compared with salt tolerant variety of wheat (*Triticum aestivum*). In: *Plant Nutrition-Physiology and Application*. pp. 463-467. M.L. Van Beusichem (Ed.). Kluwer Academic Publishers.
13. Anten, N.P.R.; Schieving, F.; Medina, E.; Werger, M.J.A. and Schuffelen, P. 1995. Optimal leaf area indices in C<sub>3</sub> and C<sub>4</sub> mono<sup>and</sup> <sup>L-</sup>dicotyledonous species at low and high nitrogen availability. *Physiologia Plantarum* 95: 541-550.
14. Araus, José L.; Matthew, P. Reynolds and Acevedo, Edmundo. 1993. Leaf posture, grain yield, growth, leaf structure and carbon isotope discrimination in wheat. *Crop Science* 33: 1273-1279.
15. Arnozis, P.A. and Findenegg, G.R. 1986. Electrical charge balance in the xylem sap of beet and sorghum plants grown with either NO<sub>3</sub><sup>-</sup> or NH<sub>4</sub><sup>+</sup> nitrogen. *Journal of Plant Physiology* 125: 441-449.
16. Arora, A.; Singh, V.; Das, R.R. and Singh, V. 1994. Yield and quality of mustard as affected by rates of N and S in inceptisols. *Journal of Oilseeds Research* 11: 273-276.
17. Arshad, M. and Frenkenberger Jr., W.T. 1991. Effect of soil properties and trace elements on ethylene production in soils. *Soil Science* 151: 377-386.
18. Arteca, R.N. and Dong, C.N. 1981. Stimulation of photosynthesis by application of phytohormones to the root system of tomato plants. *Photosynthesis Research* 2: 243-249.
19. Arteca, R.N. 1997. Plant growth substances: Principles and applications. CBS Publishers & Distributors, Delhi
20. Arthamwar, D.N.; Shelke, V.B. and Ekshinge, B.S. 1996. The effect of nitrogen and phosphorus on yield attributes, seed and oil yield of India<sup>n</sup> mustard (*Brassica juncea*). *Indian Journal of Agronomy* 41: 282-285.
21. Asare, E. and Scarisbrick, D.H. 1995. Rate of nitrogen and sulphur fertilizers on yield, yield components and seed quality of oilseed rape (*Brassica napus* L.). *Field Crops Research* 44: 41-46.
22. Awasthi, U.D. and Surajbhan. 1994. Physiological responses of barley genotypes to nitrogen levels under moisture scare conditions on light



- textured soils of central Uttar Pradesh *Indian Journal of Plant Physiology* **37** 32-34
- 23 Ballare, C L , Scopel, A.L. and Sanchez, R.A. 1989. Photomodulation of axis extension in space canopies role of the stem in the perception of light-quality signals of stand density *Plant Physiology* **89**: 1324-1330.
  - 24 Bandel, V A , Dzienä, S. and Stanford, G. 1980. Composition of N fertilizers for no-till corn *Agronomy Journal* **72**: 337-341
  - 25 Barker, A and Corey, K 1990 Ethylene evolution by tomato plants receiving nitrogen nutrition from urea. *Horticultural Science* **25** 420-421
  - 26 Bengal, B D , Deshmukh, S N and Patil, V.A. 1982 Note on the effect of growth regulators and urea on yield attributes of gram (*Cicer arietinum*) *Legume Research* **5**: 54-56.
  - 27 Bergner, C and Teichmann, C 1993. A role for ethylene in barley plants responding to soil water storage. *Journal of Plant Growth Regulation* **12** 67-72.
  - 28 Beyer, E M 1976. Silver ion a potent inhibitor of ethylene action in plants *Plant Physiology* **56** 268-271.
  - 29 Bheemaiah, G T.; Madhusudhan, Subramanyan, M.V.R. and Ismail, S. 1998 Effect of green manuring and nitrogen application on growth and yield of rainfed castor (*Ricinus communis*) alley cropped with white popinae (*Leucaena leucocephala*) *Indian Journal of Agricultural Sciences* **68**. 722-725
  30. Boller, T. 1991. Ethylene in pathogenesis and disease resistance. In: *The Plant Hormone Ethylene* pp. 293-314. A.K. Matto and J.C. Stuttle (Eds). CRC Press, Boca Raton, FL
  31. Bolton, J.K. and Brown, R.H. 1980. Photosynthesis of grass species differing in carbon dioxide fixation pathways. V. Responses of *Panicum maximum*, *Panicum miliodes*, and tall fescue (*Festuca arundinacea*) to nitrogen nutrition. *Plant Physiology* **66**: 97-100.
  - 32 Broadbent, F.E. 1978. Transformations of soil nitrogen. In: *Nitrogen and Rice* pp 108-118. International Rice Institute, Philippines.
  - 33 Bruck, H ; Payne, W.A. and Sattelmacher, B. 2000. Effects of phosphorus and water supply on yield, transpirational water-use

- efficiency and carbon isotope discrimination of pearl millet. *Crop Science* **40**: 120-125.
34. Buehler, B., Drumm, H. and Mohr, H. 1978. Investigations on the role of ethylene in phytochrome-mediated photomorphogenesis. II. Enzyme levels and chlorophyll synthesis. *Planta (Berlin)* **142**: 119-122.
  35. Bulman, P. and Smith, D.L. 1993a. Accumulation and redistribution of dry matter and nitrogen by spring barley. *Agronomy Journal* **85**: 1114-1121.
  36. Bulman, P. and Smith, D.L. 1993b. Yield and grain protein response of spring barley to ethaphon and triadimefon. *Crop Science* **33**: 798-803.
  37. Burke, J.J.; Gamble, P.E.; Hatfield, J.L. and Quisenberry, J.E. 1985. Plant morphological and biochemical responses to field water deficits. *Plant Physiology* **79**: 415-419.
  38. Butler, D.R.; Pears, E.; Child, R.D. and Brain, P. 1989. Effects of triazole growth retardants on oilseed rape: Photosynthesis of single leaves. *Annals of Applied Biology* **114**: 331-337.
  39. Byrnes, B.H. and Freney, J.R. 1995. Recent development on the use of urease ~~of~~ inhibitors in the tropics. *Fertilizer Research* **42**: 251-259.
  40. Cajuste, L.J.; Sanchez, E. and Laird, R.J. 1996. Behaviour of urea and ammonium sulfate fertilizers and their N uptake relationship in calcareous soils. In: *Nitrogen Economy in Tropical Soils*. pp. 347-353. N. Ahmad (Ed.). Kluwer Academic Publishers, Netherlands.
  41. Campbell, C.A. and Souster, W. 1982. Loss of organic matter and potential mineralizable nitrogen from Saskatchewan soils due to cropping. *Canadian Journal of Soil Science* **62**: 651-656.
  42. Cao, Y.A., Glass, D.M. and Crawford, N.M. 1993. Ammonium inhibition of Arabidopsis root growth can be reversed by potassium and auxin resistance mutations aux1, axr1, axr2. *Plant Physiology* **102**: 983-989.
  43. Carlson, D.R., Dyer, D.J., Cotterman, C.D. and Durley, R.C. 1987. The physiological basis for cytokinin induced increased in pod set in IX93-100 soyabeans. *Plant Physiology* **84**: 233-239.

44. Carvalho, M. and Basch, G. 1996. Optimisation of nitrogen fertilization: Relationship between precipitation and expected yields. *Fertilizer Research* **43**: 127-130.
45. Cathey, H.M. 1964. Physiology of growth retarding chemicals. *Annual Rev. Plant Physiology* **15**: 271-302.
46. Chai, H.H. and Hou, T.T. 1975. Studies on the volatilization gaseous nitrogen from ammonium sulfate, urea and sodium nitrate applied to soils. *Soils Fertilizers (Taiwan)* **12**: 31.
47. Champigny, M.J. 1985. Regulation of photosynthetic carbon assimilation at the cellular level: A review. *Photosynthesis Research* **6**: 273-286.
48. Champigny, M.L.; Brauer, M.; Bismuth, E.; Thi Manh, C.; Siegi, G. Van Quy, L. and Stitt, M. 1992. The short-term effect of  $\text{NO}_3^-$  and  $\text{NH}_3$  assimilation on source synthesis in leaves. *Journal of Plant Physiology* **139**: 361-368.
49. Chaudhary, S.K.; Gogulwar, N.M. and Singh, A.K. 1992. Effect of nitrogen and sulphur on seed yield and oil content of 'Varuna' mustard (*Brassica juncea*). *Indian Journal of Agronomy* **37**: 839-840.
50. Cheema, S.S.; Singh, H. and Sahota, T.S. 1987. Growth regulator for improving crop productivity. In: *Crop Productivity*. pp. 91-115. Hindustan Liver Research Foundation, Oxford and IBH Publications.
51. Chi, N.T.; Andreeva, T.F.; Strongonova, L.E.; Shirinskaya, M.G.; Zhizneyskaya, G.Ya and Nichiporovich, A.A. 1983. Photosynthesis and fixation of atmospheric nitrogen by soybean plants. *Fizilologiya Rastini* **30**: 674-681.
52. Child, R.D., Arnold, G., Hislop, E.C., Huband, N.D.S. and Stinchcanbe, G.R. 1985. Effects of some experimental triazole retardants on yield of oil seed rape. *Proceedings of British Crop Protection Conference, Weeds* **2**: 561-567.
53. Child, R.D.; Chauvaux, N.; John, K.; Ulvskov, P. and Van Onckelen, H.A. 1998. Ethylene biosynthesis in oilseed rape pods in relation to pod shatter. *Journal of Experimental Botany* **49**: 829-838.
54. Chouhan, G.S.; Singh, I. and Singh, D. 1994. Nitrogen fertilization and irrigation scheduling in mustard under Jawai Command area of Rajasthan. *Madras Agricultural Journal* **81**: 646-666.

55. Chweya, J.A. 1990. Effect of nitrogen on thiocyanate content of *Brassica oleracea* var. *acephala* leaves. *Plant and Soil* **124**: 261-263.
56. Cliquet, J.B.; Boutin, J.P.; Deleens, E. and Morot-Gaurdy, J.F. 1991. Ethaphon effects on translocation and partitioning of assimilates in *Zea mays*. *Plant Physiology and Biochemistry* **29**: 623-630.
57. Coffelt, T.A. and Howell, R.K. 1986. Effect of ethrel seed treatment on growth, yield and grade of two Virginia type peanut cultivars. *Peanut Science* **13**: 60-63.
58. Connor, D.J.; Hall, A.J. and Sadras, V.O. 1993. Effect of nitrogen content on the photosynthetic characteristics of sunflower leaves. *Australian Journal of Plant Physiology* **20**: 251-263.
59. Corey, K.; Barker, A. and Craker, L. 1987. Ethylene evolution by tomato plants under stress of ammonium toxicity. *Horticultural Science* **22**: 471-473.
60. Cornertill, I.S. and Steele, K.W. 1981. Interpretation of maize leaf analysis in New Zealand. *Journal of New Zealand Experimental Agriculture* **8**: 91-96.
61. Crosby, K.E., Aung, L.H. and Buss, G.R. 1981. Influence of 6-benzylaminopurine on fruit-set and seed development in two soybean, *Glycine max* L. (Merr) genotypes. *Plant Physiology* **68**: 985-988.
62. Dahnous, K., Vigue, G.T., Law, A.G., Kanzak, C.F. and Miller, D.G. 1982. Height and yield response of selected wheat, barley and triticales cultivars to ethephon. *Agronomy Journal* **74**: 580-582.
63. Davies, W.J. and Zhang, J. 1991. Root signals and the regulation of growth and development of plants in drying soils. *Annual Review of Plant Physiology* **42**: 55-76.
64. Davis, T.D. and Curry, E.A. 1991. Chemical regulation of vegetative growth. *Critical Reviews in Plant Science* **10**: 151-188.
65. Davis, T.D., Steffens, G.L. and Sankhla, N. 1988. Triazole plant growth regulators. In: *Horticultural Reviews*. Vol. 10. pp. 63. J. Janick (Ed). Timber Press, Portland, OR.
66. Dayal, J.; Nandwal, A.S. and Kaur, H. 1993. Gas exchange and water relation studies in cycocel treated pigeonpea in response to water stress. *Indian Journal of Plant Physiology* **36**: 263-265.

67. De-Bouille, P.; Sotta, B.; Miginiac, E. and Merrien, A. 1989. Hormones and pod development in oilseed rape (*Brassica napus*). *Plant Physiology* **90**: 876-880.
68. DeJong, T.M. and Doyle, J.F. 1985. Seasonal relationship between leaf nitrogen content (photosynthetic capacity) and leaf canopy light exposure in peach (*Prunus persica*). *Plant Cell Environment* **8**: 701-706.
69. Denmead, O.T.; Freney, J.R. and Simpson, J.R. 1982. Dynamics of ammonia volatilization during furrow irrigation of maize. *Soil Science Society of American Journal* **46**: 149-155.
70. Dey, S.K.; Singh, V.P. and Murthy, K.S. 1989a. Effect of low light at different stages of development on growth and yield of rice. *Indian Journal of Plant Physiology* **32**: 74-77.
71. Dey, S.K.; Singh, V.P. and Murthy, K.S. 1989b. Influence of nitrogen levels on photosynthesis and productivity in early rice varieties. *Indian Journal of Plant Physiology* **32**: 364-368.
72. Dhakal, M.R. and Erdei, L. 1986. Long-term effects of plant hormones on  $K^+$  levels and transport in young wheat plants of different  $K^+$  status. *Physiologia Plantarum* **68**: 632-636.
73. Dharan, K.R., Bassi, P.K. and Spencer, M.S. 1981. Effects of carbon dioxide on ethylene production and action in sunflower plants. *Plant Physiology* **68**: 831-834.
74. Dhoble, V.M. 1998. Response of sunflower (*Helianthus annuus*) hybrids to nitrogen and phosphorus in rainfed conditions. *Indian Journal of Agronomy* **43**: 138-142.
75. Dijkstra, P. and Kuiper, P.J.C. 1989. Effects of exogenously applied growth regulators on shoot growth of inbred lines of *Plantago major* differing in relative growth rate: differential response to gibberellic acid and (2-chloroethyl)-trimethyl-ammonium chloride. *Physiologia Plantarum* **77**: 512-518.
76. Dodd, I.C. 2001. Do phytohormones control leaf growth of nitrogen deprived plants? In: *Phytohormones in Crop Productivity Under Different Environments*. pp. 1-24. N.A. Khan and Samiullah (Eds.). Scientific Publishers, Jodhpur, India.
77. Doran, J.W. 1980. Soil microbial changes associated with reduced tillage. *Soil Science Society of American Journal* **44**: 765-771.

78. Dowllet, H.N.A. and Kumamoto, J. 1972. The conversion of 2-hydroxyethylhydrazine to ethylene. *Plant Physiology* 49: 696-699.
79. Downey, R.K. and Rimmer, S.R. 1993. Agronomic improvement in oilseed Brassicas. *Advances in Agronomy* 50: 1-65.
80. Downey, R.K. and Robbelen, G. 1989. Brassica species. In: *Oilcrops of the World*. pp. 339-362. G. Robbelen, R.K. Downey and A. Ashri (Eds.). McGraw-Hill, New York.
81. Draber, W. 1977. Naturliche and synthetische pflanzenwachstum sregulatoren. In: *Chemie der Pflanzenschutz-und Schadlingsbekämpfungsmittel*. R. Wegler (Ed). Band 4. Springer-Verlag, Berlin, Heidelberg, New York.
82. Dubey, O.P. and Khan, R.A. 1993. Effect of nitrogen and sulphur on dry matter, grain yield and nitrogen content at different growth stages of mustard (*Brassica juncea*) under irrigated vertisol. *Indian Journal of Agronomy* 38: 270-276.
83. Dubey, O.P.; Sahu, T.R. and Garg, D.C. 1994. Response and economics in relation to nitrogen and sulphur in Indian mustard (*Brassica juncea*). *Indian Journal of Agronomy* 39: 49-53.
84. Dubey, O.P.; Sahu, T.R.; Garg, D.C. and Khan, R.A. 1993. Response of mustard to sulphur and nitrogen under irrigated vertisol condition. II. Effect of S and N on ancillary characters, yield and quality. *Journal of Oilseed Research* 10: 11-15.
85. Durand, J.L.; Onillon, B.; Schnyder, H. and Rademacher, I. 1995. Drought effects on cellular and spatial parameters of leaf growth in tall fescue. *Journal of Experimental Botany* 46: 1147-1155.
86. Ecker, J.R. 1995. The ethylene signal transduction pathway in plants. *Science* 268: 667-675.
87. El-Keltawi, N.F. and Rodney, C. 1986. Influence of ethephon and diaminozide on growth and essential oil content of peppermint (*Menta piperita*) and sage (*Saliva officinalis*). *Phytochemistry* 25: 1285-1288.
88. Epstein, E. 1972. Mineral nutrition of plants: principles and perspectives. Wiley, New York.
89. Erdei, L. and Dhakal, M.R. 1980. Investigations of the effect of phytohormones and growth regulators on the transpiration, stomata

aperture and photosynthesis of pepper (*Capsicum annum* L.) and tomato (*Lycopersicon esculentum* Mill.) plants. *Botany* **54**: 207-220. 8 8 2 c

90. Ericsson, T.; Rytter, L. and Linder, S. 1992. Nutritional dynamics and requirements of short rotation forests. In: *Ecophysiology of Short Rotation Forest Crops*. pp. 35-65. C.P. Mitchell; J.B. Ford-Robertson; T. Hinekley and L. Sennerby-Forsse (Eds.). Elsevier Applied Science, New York.
91. Evans, J.R. 1989. Photosynthesis and nitrogen relationships in leaves of C<sub>3</sub> plants. *Oecologia* **78**: 9-19.
92. Evans, J.R. and Seemann, J.R. 1989. The allocation of protein nitrogen in the photosynthetic apparatus: costs, consequences and control. In: *Photosynthesis Alan*. pp. 183-205. W. Briggs (Ed.). R. Liss, New York.
93. Evans, J.R. and Terashima, I. 1988. Photosynthetic characteristics of spinach leaves grown with different nitrogen treatments. *Plant Cell Physiology* **29**: 157-165.
94. Fageria, N.K. 1984. Fertilization and mineral nutrition of rice. EMBRAPA-CNPAP/Editora Campus, Rio de Janeiro.
95. Farooqi, A.H.A. and Sharma, S. 1988. Effect of growth retardants on flowering of *Rosa damascena* Mill. *Proceedings of the International Congress of Plant Physiology*, New Delhi.
96. Feiz, T.E.; Miller, B.C. and Pan, W.L. 1994. Assessment of specially variable nitrogen fertilizer management in winter wheat. *Journal of Production Agriculture* **7**: 86-94.
97. Feng, J. and Barker, A.V. 1992. Ethylene evolution and ammonium accumulation by tomato plants with various nitrogen forms and regimes of acidity. *JOURNAL Plant Nutr.* **15**: 2457-2469. 7
98. Feng, J. and Barker, A.V. 1993. Polyamine concentration and ethylene evolution in tomato plants under nutritional stress. *Hort. Science* **28**: 109-110.
99. Ferguson, R.B., Kissel, D.E., Koelliker, J.K. and Basal, W. 1984. Ammonia volatilization from surface-applied urea: effect of hydrogen ion buffering capacity. *Soil Science Society of American Journal* **48**: 578-582. e

100. Field, C. 1983. Allocating leaf nitrogen for the maximum of carbon gain: Leaf age as a carbon on the allocation programme. *Oecologia* **56**: 341-347.
101. Foroutan-Pour, K.; Ma, B.L. and Smith, D.L. 1997. Protein accumulation potential in barley seeds as affected by soil-and peduncle-applied N and peduncle-applied plant growth regulators. *Physiologia Plantarum* **100**: 190-201.
102. Foster, K.R., Taylor, J.S. and Reid, D.M. 1991. Tillering and yield responses to ethephon in three barley cultivars. *Crop Science* **31**: 130-134.
103. Francis, A.J. 1982. Effects of acidic precipitation and acidity on soil microbial process. *Water, Air and Soil Pollution* **18**: 375-394.
104. Freney, J.R., Leuning, R., Simpson, J.R., Denmead, O.T. and Muirhead, W.A. 1985. Estimating ammonia volatilisation from flooded rice fields by simplified techniques. *American Journal of Soil Science Society* **49**: 1049-1054.
105. Freney, J.R., Simpson, J.R. and Denmead, O.T. 1983. Volatilization of ammonia. In: *Gaseous Loss of Nitrogen from Plant-Soil Systems*. pp. 1-32. J.R. Freney (Ed). Martinus Nijhoff/or W. Jink Publishers, The Hague, Netherlands.
106. Friend, D.J.C. 1985. Brassica. In: *CRC Handbook of Flowering*. pp. 48-77. A.H. Halery (Ed).CRC Press, Boca Raton, FL.
107. Gajri, P.R.; Prihar, S.S. and Arora, V.K. 1993. Independence of nitrogen and irrigation effects on growth and input-use efficiencies in wheat. *Field Crops Research* **31**: 71-86.
108. Gardner, J.C.; Maranville, J.W. and Paparozzi, E.T. 1994. Nitrogen use efficiency among diverse sorghum cultivars. *Crop Science* **34**: 728-733.
109. Gashaw, L. and Mugwira, L.M. 1981. Ammonium-N and nitrate-N effects on the growth and mineral compositions of *triticale* wheat and rye. *Agronomy Journal* **73**: 45-51.
110. Gendy, A. and Marquard, R. 1989. Studies on the effect of nitrogen fertilization and growth regulators on seed yield and some quality criteria of oilseed-rape (*Brassica napus* L.). *Fett. Wiss. Technology* **91**: 353-357.



111. Ghatak, S.; Sounda, G. and Jana, P.K. 1992. Effect of irrigation and nitrogen on seed and oil content of Indian mustard (*Brassica juncea*). *Indian Journal of Agricultural Sciences* **62**: 664-668.
112. Ghosh, D.C.; Pand, P.K. and Sahoo, P.M. 1995. Response of rainfed rapeseed (*Brassica campestris* L.) to NPK. *Indian Journal of Agricultural Research* **29**: 5-9.
113. Gilmour, J.T. 1984. The effects of soil properties on nitrification and nitrification inhibition. *Soil Science Society of American Journal* **48**: 1262-1266.
114. Glimskar, A. and Ericsson, T. 1999. Relative nitrogen limitation at steady-state nutrition as a determinant of plasticity in five grassland plant species. *Annals of Botany* **84**: 413-420.
115. Gomez, K.A. and Gomez, A.A. 1984. Statistical procedure for agricultural research. Wiley International Publishers, New York.
116. Gopalkrishnana, S. and Srinivasan, P.S. 1975. Effect of planofix on NAA formation on groundnut. *Pesticides* **23**: 23-25.
117. Govindarajan, A.G. and Poovaiah, B.W. 1982. Effect of root zone carbon dioxide enrichment on ethylene inhibition of carbon assimilation in potato plants. *Physiologia Plantarum* **55**: 465-469.
118. Graan, T. and Boyar, J.S. 1990. Very high CO<sub>2</sub> partially restores photosynthesis in sunflower at lower water potentials. *Planta* **181**: 378-384.
119. Green, J. 1983. The effect of potassium and calcium on cotyledon expansion and ethylene evolution by cytokinins. *Plant Physiology* **57**: 57-61.
120. Greenwood, D.J. 1982. Nitrogen supply and crop yield. The global scene. *Plant and Soil* **67**: 45-59.
121. Grewal, H.S. and Kolar, J.S. 1990. Response of *Brassica juncea* to chlorocholine chloride and ethrel sprays in association with nitrogen application. *Journal of Agricultural Science* **114**: 87-91.
122. Grewal, H.S.; Kolar, J.S.; Cheema, S.S. and Singh, G. 1993. Studies on the use of growth regulators in relation to nitrogen for enhancing sink capacity and yield of gobhi-season (*Brassica napus*). *Indian Journal of Plant Physiology* **36**: 1-4.

123. Grodzinski, B.; Boesel, I. and Horton, R.F. 1982. Ethylene release from leaves of *Xanthium strumarium* and *Zea mays* L. *Journal of Experimental Botany* **33**: 344-354.
124. Groffman, P.M. 1995. A conceptual assessment of the importance of denitrification as a source of soil nitrogen loss in tropical agroecosystems. *Fertilizer Research* **42**: 139-148.
125. Guiamet, J.J., Willemoes, .G. and Montaldi, E.R. 1989. Modulation of progressive leaf senescence by the red: far-red ratio of incident light. *Botanical Gazzette* **150**: 148-151.
126. Guruprasad, A. and Guruprasad, K.N. 1988. Interaction of potassium ions and gibberellin in the control of hypocotyls growth in *Amaranthus caudatus*. *Physiologia Plantarum* **74**: 154-158.
127. Haru, K., Naitu, K. and Suzuki, H. 1982. Differential effects on benzyladenine and potassium on DNA, RNA, protein and chlorophyll contents and expansion growth of detached cucumber cotyledon in the dark and light. *Physiologia Plantarum* **55**: 247-254.
128. Heichel, G.L. 1980. Energy for agriculture. In: *Moving up the Yield Curve, Advances and Obstacles*. pp. 91-103. L.S. Murphy, E.C. Doll and L.F. Welsh (Eds.). American Society of Agronomy, Special Publication, No. 39.
129. Heilman, P.E. and Fu-Guang, X. 1993. Influence of nitrogen on growth and productivity of short-rotation *Populus trichocarpa* x *Populus deltoids* hybrids. *Canadian Journal of Forest Research* **23**: 1863-1869.
130. Heilman, P.E. and Fu-Guang, X. 1994. Effects of nitrogen fertilization on leaf area, light interception and productivity of short-rotation *Populus trichocarpa* x *Populus deltoids* hybrids. *Canadian Journal of Forest Research* **24**: 166-173.
131. Hewitt, E.J. 1963. The essential nutrient elements: requirements and interaction in plants. In: *Plant Physiology: A Treatise*. pp. 137-329. F.C. Steward (Ed.). Academic Press, New York.
132. Holmes, M.G. 1981. Spectral distribution of reaction within plant canopies. In: *Plants and Daylight Spectrum*. pp. 147-158. H. Smith (Ed). Academic Press, London.

133. Howard, D.D. and Tyler, D.D. 1989. Nitrogen source application method for no-tillage corn. *Soil Society of American Journal* **53**: 1573-1577.
134. Hua, J., Chang, C., Sun, Q. and Meyerowitz, E.M. 1995. Ethylene insensitivity conferred by arabidopsis ERS gene. *Science* **269**: 1712-1714.
135. Huber, S.C.; Sugiyama, T. and Alberte, R.S. 1989. Photosynthetic determinants of growth in maize plants: effects of nitrogen nutrition on growth, carbon fixation and photochemical features. *Plant Cell Physiology* **30**: 1063-1072.
136. Hunter, S.C. and Ohlrogge, J.B. 1998. Regulation of spinach chloroplast acetyl-CoA carboxylase. *Archives of Biochemistry and Biophysics* **15**: 170-178
137. Hussain, A., Black, C.R., Taylor, I.B. and Roberts, J.A. 1999. Soil compaction: a role for ethylene in regulating leaf expansion and shoot growth in tomato. *Plant Physiology* **121**: 1227-1237.
138. levinsh, G. and Kreicbergs, O. 1992. Endogenous rhythmicity of ethylene production in growing intact cereal seedlings. *Plant Physiology* **100**: 1389-1391.
139. Imsande, J. 1992. Agronomic characteristics that identify high yield, high soybean genotypes. *Agronomy Journal* **84**: 409-414.
140. Ingestad, T. 1980. Growing nutrition, and nitrogen fixation in gray alder at varied rate of nitrogen addition. *Physiologia Plantarum* **50**: 353-364.
141. Jana, B.K. and Kabir, K. 1991. Effect of ethrel on seed production of cauliflower cv. Dania. *Crop Research* **4**: 222-224.
142. Jansson, S.L. and Perrson, J. 1982. Mineralization and immobilization of soil nitrogen. In: *Nitrogen in Agricultural Soils*. pp. 229-252. F.J. Stevenson (Ed). Society of Agronomy, Madison, Wisc, USA. *Tables*
143. Jaworski, E.G. 1971. Nitrate reductase assay in intact plant tissues. *Biochemistry and Biophysics Research Communication* **43**: 1274-1279.
144. Jensen, C.R. and Tophoj, H. 1985. Potassium induced improvement of yield response in barley exposed to soil water stress. *Irrigation Science* **6**: 118-129.

145. Jeschke, W.D.; Wolf, O. and Hartung, W. 1992. Effect of NaCl salinity on flows and partitioning of C, N and mineral ions in whole plants of white lupin (*Lupinus albus* L.) *Journal of Experimental Botany* **43**: 777-788.
146. Johnson, J.D. 1984. The effects of ethylene on stomatal conductance and transpiration at various levels of water stress. *Plant Physiology* **75**: S-129.
147. Joshi, R.K.; Mishra, S.D. and Gaur, B.K. 1987. Productivity of dormant groundnut as affected by ethrel and benzyladenine. *Indian Journal of Agricultural Science* **57**: 179-182.
148. Kao, C.H. and Yang, S.F. 1982. Light inhibition of the conversion of 1-aminocyclopropane-1-carboxylic acid to ethylene in leaves is mediated through carbon dioxide. *Planta* **155**: 261-266.
149. Kasele, I.N.; Shanahan, J.F. and Nielsen, D.C. 1995. Impact of growth retardants on corn leaf morphology and gas exchange traits. *Crop Science* **35**: 190-194.
150. Katyal, J.C., Singh, B., Vlek, P.L.G. and Craswell, E.T. 1985. Fate and efficiency of nitrogen fertilizers applied to wet land rice II. Punjab, India. *Soil Science* **140**: 194-201.
151. Kays, S.J. and Pallas Jr., J.E. 1980. Inhibition of photosynthesis by ethylene. *Nature* **385**: 51-52.
152. Kerr, P.S.; Israel, D.W.; Huber, S.C. and Rufty, T.W. 1986. Effect of supplemental NO<sub>3</sub> on plant growth and components of photosynthetic carbon metabolism in soybeans (*Glycine max*). *Canadian Journal of Botany* **64**: 2020-2027.
153. Khan, G.M. and Agarwal, S.K. 1988. Physiological basis of growth analysis of mustard (*Brassica juncea* Coss.) in relation to sowing methods, water stress and nitrogen levels. *Haryana Journal of Agronomy* **4**: 91-96.
154. Khan, N.A. 1996a. Effect of gibberellic acid on carbonic anhydrase, photosynthesis, growth and yield of mustard. *Biologia Plantarum* **38**: 145-147.
155. Khan, N.A. 1996b. Response of mustard to ethrel spray and basal and foliar application of nitrogen. *Journal of Agronomy and Crop Science* **176**: 331-334.

156. Khan, N.A. 1997. Evaluation of gibberellic acid for increasing yield efficiency of mustard cultivars. *Test of Agrochemicals and Cultivars* **18**: 26-27.
157. Khan, N.A. 1998. Effect of ethrel spray on yield of mustard cultivars under non irrigated conditions. *Test of Agrochemicals and Cultivars* **19**: 38-39.
158. Khan, N.A., Ansari, H.R., Mobin, M. and Samiullah. 1999. Effect of gibberellic acid on morphophysiology and yield of mustard (*Brassica juncea* L.) grown with basal levels of nitrogen and phosphorus. In: *Plant Physiology of Sustainable Agriculture*. pp. 214-224. G.C. Srivastava, K. Singh and M. Pal (Eds.). Pointer Publishers, Jaipur, India.
159. Khan, N.A.; Ansari, H.R. and Mobin, M. 1996. Effect of gibberellic acid and nitrogen on carbonic anhydrase activity and mustard biomass. *Biologia Plantarum* **38**: 601-603.
160. Khan, N.A.; Ansari, H.R. and Samiullah. 1997. Effect of gibberellic acid spray and basal nitrogen and phosphorus on productivity and fatty acid composition of rapeseed-mustard. *Journal of Agronomy and Crop Science* **179**: 29-33.
161. Khan, N.A.; Ansari, H.R. and Samiullah. 1998. Effect of gibberellic acid spray during ontogeny of mustard on growth, nutrient uptake and yield characteristics. *Journal of Agronomy and Crop Science* **181**: 61-63.
162. Khan, N.A.; Lone, N.A. and Samiullah. 2000. Response of mustard (*Brassica juncea* L.) to applied nitrogen with or without ethrel spray under non-irrigated conditions. *Journal of Agronomy and Crop Science* **183**: 1-4.
163. Khan, N.A.; Mobin, M.; Mir, R. and Samiullah. 2001. Phytohormones and nutrients interaction on growth and productivity of crops. In: *Phytohormones in Crop Productivity Under Different Environments*. pp. 167-178. N.A. Khan and Samiullah (Eds.). Scientific Publishers, Jodhpur, India.
164. Khan, N.A.; Samiullah; Afridi, M.M.R.K. and Ansari, S.A. 1990. Response of six mustard varieties to different combinations of nitrogen and phosphorus. *Indian Journal of Agronomy* **35**: 412-414.
165. Khanpara, V.D.; Porwal, B.L.; Sahu, M.P. and Patel, C. 1993. Effect of sulphur and nitrogen on yield attributes and seed yield of Indian mustard (*Brassica juncea*) on vertisol. *Indian Journal of Agronomy* **38**: 588-592.

- 166 Kieber, J.J.; Rathenberg, M.; Roman, G.; Feldmann, K.A. and Ecker, J.R. 1993. CTR1, a negative regulator of ethylene response pathway in arabidopsis, encodes a member of the Raf family of protein kinases. *Cell* **72**: 427-441.
- 167 Kirkby, E.A. 1981. Plant growth in relation to nitrogen supply. In: *Terrestrial nitrogen cycles, processes, ecosystem strategies and management impacts*. pp. 249-267. F.E. Clarke and T. Ross Wall (Eds.). Ecological Bulletin. Stockholm 33.
- 168 Kleinhoffs, A. and Warner, R.L. 1990. Advances in nitrate assimilation. In: *The Biochemistry of Plants: A Comprehensive Treatise*. pp. 89-120. P.K. Stumpf and E.E. Conn (Eds.). Vol. 16, Academic Press, New York.
- 169 Konings, H. and Jackson, M.B. 1979. A relationship between rates of ethylene production by roots and the promoting or inhibiting effects of exogenous ethylene and water on root elongation. *Zeitschrift Pflanzenphysiol* **92**: 385-398.
- 170 Koritsas, V.M. 1988. Effect of ethylene and ethylene precursors on protein phosphorylation and xylogenesis in tuber explants of *Helianthus tuberosus* (L.). *Journal of Experimental Botany* **39**: 375-386.
- 171 Kozaki, A. and Sasaki, Y. 1999. Light dependent changes in redoxstatus of the plastidic acetyl CoA carboxylase and its regulatory component. *Biochemistry Journal* **339**: 541-546.
172. Krishnamoorthy, H.N. 1993. Physiology of plant growth and development. Atma Ram and Sons, Delhi.
173. Kuiper, D. 1988. Growth response of *Plantago major* L. ssp. Pleiosperma (pilger) to change in mineral supply. *Plant Physiology* **87**: 557-567.
174. Kuiper, D.; Schuit, J. and Kuiper, P.J.C. 1989. Effect of internal and external cytokinins concentration on root growth and shoot to root ratio of *Plantago major* ssp. Pleiosperma at different nutrient conditions. In: *Structural and Functional Aspects of Transport in Roots*. pp. 183-188. B.C. Loughman, O. Gasparikova and J. Kolek (Eds.). Kluwer Academic Publishers, London.
175. Kulkarni, S.S.; Chetti, M.B. and Uppar, O.S. 1995. Influence of growth retardants on biochemical parameter in sunflower. *Journal of Maharashtra Agricultural Universities* **20**: 352-354.

- 176 Kullimann, A , Ogunlela, V B and Geisler, G. 1989 Concentrations and distribution of some mineral elements in oilseed rape (*Brassica napus* L.) plants in relation to nitrogen supply *Agronomy and Crop Science* **163**: 225-235
- 177 Kumar, A and Elston, J 1993 Leaf expansion in *Brassica* species in response to water stress. *Indian Journal of Plant Physiology* **36**: 220-222
- 178 Kumar, A , Singh, D P., Singh, P and Singh, D 1994 Influence of water stress on photosynthesis, transpiration, water use efficiency and yield of *Brassica juncea* L. *Field Crops Research* **37**: 95-101.
- 179 Kumar, P , Dhingra, P K , Singh, V P and Pannu, R K 1995. Oil yield, oil content and nitrogen uptake in different *Brassica* genotypes as affected by irrigation and nitrogen application *Crop Research* **10**: 145-147
- 180 Kumar, R , Singh, V.P. and Faroda, A S 1989. Studies on the oil yield and nitrogen uptake by raya cultivars as influenced by nitrogen levels under irrigated and conserved soil moisture conditions. *Haryana Journal of Agronomy* **5**: 182-184
- 181 Kumar, S and Bharti, S. 1988. Effect of CCC and FAP on water status and yield of sunflower (*Helianthus annuus* L.) under stimulated drought conditions *Indian Journal of Plant Physiology* **31**: 381-387.
- 182 Kurvitis, A and Kirkby, E A 1980 The uptake of nutrients by sunflower plants (*Helianthus annuus*) growing in continuous flowing culture system, supplied with nitrate or ammonium as nitrogen source. *Zeitschrift fur Pflanzenernahrung Bodenkunde* **143**: 140-149.
- 183 Kushad, M.H. and Pooviah, B.W 1984 Deferral of senescence and abscission by chemical inhibition of ethylene synthesis and action in bean explants *Plant Physiology* **76**: 293-296.
- 184 Ladd, J.N. and Jackson, R.B. 1982. Biochemistry of ammonification. In: *Nitrogen in Agricultural Soils*. pp. 173.228. F.J. Stevenson (Ed). ASA-CSSA-SSS, Madison, Wisconsin.
- 185 Lauer, M.J. and Boyer, J.S. 1992 Internal CO<sub>2</sub> measures directly in leaves: abscisic acid and low leaf water potential cause opposing effects. *Plant Physiology* **98**: 1010-1016.

186. Lea, P.J.; Robinson, S.A. and Steward, G.R. 1990. The enzymology and metabolism of glutamine, glutamate and asparagines. In: *The Biochemistry of Plants. Intermediary Nitrogen Metabolism*. pp. 121-159. B.J. Mifflin and P.J. Lea (Eds.). Vol. 16. Academic Press, San Diego, CA.
187. Leary, W.P. and Oplinger, E.S. 1983. Effect of plant growth regulators and grain crops. *Proc. Plant Growth Regulators Society of America* **10**: 277-286.
188. Lecoeur, J.; Wery, J.; Ture, O. and Tardieu, F. 1995. Expansion of pea leaves subjected to short water deficit: cell number and cell size are sensitive stress at different periods of leaf development. *Journal of Experimental Botany* **46**: 1093-1101.
189. Lee, S.H. and Reid, D.M. 1997. The role of endogenous ethylene in the expansion of *Helianthus annuus* leaves. *Canadian Journal of Botany* **75**: 501-508.
190. Leitch, M.H. and Kuat, O. 1999. Effect of plant growth regulators on stem extension and yield components of Linseed (*Linum usitatissimum*). *Journal of Agricultural Science* **132**: 189-199.
191. Lemcoff, J.H. and Loomis, R.S. 1986. Nitrogen influences on yield determination in maize. *Crop Science* **26**: 1017-1022.
192. Leopold, A.C. and Kriedemann, P.E. 1979. Chemical regulation of growth and development. In: *Plant Growth and Development*. pp. 433-458. 2<sup>nd</sup> Ed. Mc Graw-Hill Inc., New York.
193. Lincoln, J.E. and Fischer, R.L. 1988. Diverse mechanisms for the regulation of ethylene-inducible gene expression. *Molecular Gene Genetics* **212**: 71-75.
194. Lindner, R.C. 1944. Rapid analytical methods for some of the more common inorganic constituents of plant tissues. *Plant Physiology* **19**: 70-89.
195. Liu, H.S.; Peng, W.B.; Meng, F.T.; Yuan, J.P. and Wang, D.Q. 1993. Effects of S-3307 on morphological and some physiological characteristics of wheat seedlings. *Plant Physiology Communication* **29**: 354-355.



196. Liu, J.H., Mukherjee, I. and Reid, D.M. 1990. Adventitious rooting in hypocotyls of sunflower (*Helianthus annuus*). III. The role of ethylene. *Physiologia Plantarum* **78**: 268-276.
197. Liu, Z. and Dickmann, D.I. 1992. Absciscic acid accumulation in leaves of two contrasting hybrid poplar clones affected by nitrogen fertilization plus cyclic flooding and drought. *Tree Physiology* **11**: 109-122.
198. Lizada, M.C.C. and Yang, S.F. 1979. A simple and sensitive assay for 1-aminocyclopropane-1-carboxylic acid. *Anal. of Biochemistry* **100**: 140-145.
199. Lloyd, D. 1993. Aerobic denitrification in soils and sediments: from fallacies to facts. *Trends in Ecological Evolution* **8**: 352-356.
200. Longstreth, D.J. and Nobel, P.S. 1980. Nutrient influences on leaf photosynthesis. *Plant Physiology* **65**: 541-543.
201. Lovatt, C.J. 1986. Salinity stress results in ammonia intoxication. *Plant Physiology* **80**(Suppl.): 10.
202. Lu, C. and Zhang, J. 1999. Effects of water stress on photosystem II photochemistry and its thermostability in wheat plants. *Journal of Experimental Botany* **50**: 1199-1206.
203. Lu, Z. and Neumann, P.M. 1998. Water-stressed maize, barley and rice seedlings show species diversity in mechanisms of leaf growth inhibition. *Journal of Experimental Botany* **49**: 1945-1952.
204. Ludlow, M.M. and Muchow, R.C. 1990. A critical evaluation of traits for improving crop yields in water-limited environments. *Advances in Agronomy* **43**: 107-153.
205. Lurie, S.; Ronen, R.; Lipsker, Z. and Aloni, B. 1994. Effects of paclobutrazol and chilling temperatures on lipids, antioxidants and ATPase activity of plasma membrane isolated from green bell pepper fruits. *Physiologia Plantarum* **91**: 593-598.
206. Lurssen, K. 1982. Manipulation of crop growth by ethylene and some implications of the mode of generation. In: *Chemical Manipulation of Crop Growth and Development*. J.S. (Ed). Butterworth Scientific, London.
207. Lutman, P.J.W. 1993. The occurrence and persistence of volunteer oilseed rape (*Brassica napus*). *Aspects of Applied Biology* **35**: 29-37.

- 208 Lyster, S , Morgan, M A and O'Toole, P. 1980 Ammonia volatilization from soil fertilization from soils fertilized with urea and ammonium nitrate *Journal of Life Science Royal Dublin Society* **1**: 167-176.
- 209 Ma, B L and Smith, D.L 1992 Apical development of spring barley under field conditions in Northeastern North America *Crop Science* **32**: 144-149
- 210 Ma, B L , Dwyer, L M and Smith, D.L. 1994a Evaluation of peduncle perfusion for in vivo studies of carbon and nitrogen distribution in cereal crops *Crop Science* **34**: 1584-1588
- 211 Ma, B L , Leibovitch, S and Smith, D L. 1994b Plant growth regulator effects on protein content and yield of spring barley and wheat. *Journal of Agronomy and Crop Science* **172**: 9-18
- 212 Madhulety, T Y and Prakash, V 1988. Gigas bread wheat in respective of nitrogen economy I Comparative response of gigas wheat to single staggered application of nitrogen in respect of growth and yield *Indian Journal of Plant Physiology* **28**: 349-356
- 213 Mai, V V , Andreeva, T F and Nichiporovich, A A 1987 Growth and gas exchange of Indian mustard in the absence of exogenous nitrogen *Soviet Plant Physiology*. **34**: 194-203
- 214 Makeev, A V , Krendeleva, T.E. and Mokronsov, A T. 1992. Photosynthesis and abscisic acid *Soviet Plant Physiology* **39**: 118-126
- 215 Marschner, H 1983 General introduction to the mineral nutrition of plants In: *Encyclopedia of Plant Physiology*. pp. 5-60. A. Lauchli and R L Bielecki (Eds) New Ser , vol 15A Springer-Verlag, New York
- 216 Marschner, H. 1986. Mineral nutrition of higher plants. Academic Press, New York
- 217 Mathias, J.N , Bradburone, J A. and Dupree, M. 1989. GA effects on greening in pea seedlings. *Plant Physiology* **91**: 19-22.
218. Mathur, D. and Wattal, P.N. 1996. Physiological analysis of growth and development in three species of rape-mustard (*Brassica juncea*, *B. campestris* and *B. napus*) under irrigated and unirrigated conditions. *Indian Journal of Plant Physiology* **1**: 171- 174.
- 219 Mattoo, A.K. and White, W.B. 1991. Regulation of ethylene biosynthesis. In: *The Plant Hormone Ethylene*. pp. 21-42. A.K. Mattoo

- and J.C. Suttle (Eds). CRC Press, Boca Raton Ann Arbor, Boston, London.
220. Mattoo, A.K., Mehta, R.A. and Chung, K.H. 1997. Molecular regulation of ethylene biosynthesis and fruit ripening. *Plant Molecular Biology and Biochemistry* **15**: 53-67.
  221. Mc Keon, T.A., Hoffman, N.E. and Yang, S.F. 1982. The effect of plant hormone pretreatments on ethylene production and synthesis of 1-aminocyclopropane-1-carboxylic acid in water-stressed wheat leaves. *Planta* **155**: 437-443.
  222. McGrath, S.P. and Zhao, F.J. 1996. Sulphur uptake, yield responses and the interactions between nitrogen and sulphur in winter oilseed rape (*Brassica napus*). *Journal of Agricultural Sciences Cambridge* **126**: 53-62.
  223. Menon, K.K.G. and Srivastava, H.C. 1984. Increasing plant productivity through improved photosynthesis. *Plant Science* **93**: 359-378.
  224. Mhatre, M.; Mirza, M and Rao, P.S. 1998. Stimulating effect of ethylene inhibitors and ortho-caumeric acid in tissue cultures of cucumber (*Cucumis sativas* L.). *Indian Journal of Experimental Biology* **3**: 104-107.
  225. Michelozzi, M.; Johnson, J.D. and Warrag, E.I. 1995. Response of ethylene and chlorophyll in two eucalyptus clones during drought. *New Forests* **9**: 197-204.
  226. Migge, A. and Becker, T.W. 1996. In tobacco leaves, the genes encoding the nitrate-reducing or the ammonium assimilating enzymes are regulated differently by external nitrogen-sources. *Plant Physiology and Biochemistry* **34**: 665-671.
  227. Mishra, P.C.; Choudhury, N.K. and Biswal, U.C. 1985. A critical evaluation of the role of minerals on plant development. *Indian Journal of Plant Nutrition* **4**: 41-45.
  228. Mishra, S.D., Joshi, R.K. and Gaur, B.K. 1984. Preferential effect of gibberellic acid, benzyladenine and ethrel at pegging stage in groundnut (*Arachis hypogaeae*). *Acta Botanica Indica* **12**: 123-128.
  229. Mishra, S.N. and Srivastava, H.S. 1985. Role of inorganic nitrogen in synthesis and degradation of protein in maize leaves. *Indian Journal of Plant Physiology* **28**: 43-52.

230. Mobin, M. 1999. Morphophysiology and productivity of mustard in relation to gibberellic acid and sulphur application. Ph.D. Thesis, Aligarh Muslim University, Aligarh, India.
231. Moes, J. and Stobbe, E.H. 1991a. Barley treated with ethephon: I. Yield components and net grain yield. *Agronomy Journal* **83**: 86-90.
232. Moes, J. and Stobbe, E.H. 1991b. Barley treated with ethephon: II. Tillering pattern and its impact on yield. *Agronomy Journal* **83**: 90-94.
233. Mohan Ram, H.Y. and Rina, S. 1982. Modification of growth and sex expression in *Cannabis sativa* by aminoethoxy vinyl-glycine and ethephon. *Zeitschrift Pflanzen Physiol* **105**: 165-172.
234. Moll, R.H.; Kamprath, E.J. and Jackson, W.A. 1982. Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agronomy Journal* **74**: 562-564.
235. More, S.D. and Varade, S.B. 1976. Volatilization losses of ammonia from different nitrogen carriers as affected by soil-moisture, organic matter and method of fertilizer application. *Journal of Indian Society of Soil Sciences* **26**: 112-115.
236. Morgan, D.G. 1980. Factors affecting fruit and seed development in field beans and oilseed rape. In: *Joint DPGRG and BPGRG symposium: aspects and prospects of plant growth regulators*. Monograph **6**: 151-164.
237. Morgan, D.G., Kieller, D.R. and Prynne, A.O. 1983. Nitrogen, growth regulators and pod development in oilseed rape. In: *Interaction Between Nitrogen and Growth Regulators in Control of Plant Development*. M.B. Jackson, (Ed). British Plant Growth Regulator Group Monograph **9**: 75-86.
238. Morgan, P.W., Sarquis, J.L., He, C., Jordan, W.R. and Drew, M.C. 1993. Regulation of ethylene biosynthesis in maize root response to stress. In: *Cellular and Molecular Aspects of the Plant Hormone Ethylene*. pp. 232-237. J.C. Pech, A. Latche and C. Balague (Eds.). Kluwer Academic Publishers, Dordrecht, The Netherlands.
239. Morris, C.F.; Ferguson, D.L. and Paulsen, G.M. 1989. Nitrogen fertilizer management with foliar fungicide and growth regulator for hard winter wheat production. *Applied Agronomical Research* **4**: 135-140.

240. Muchow, R.C. and Sinclair, T.R. 1994. Nitrogen response of leaf photosynthesis and canopy radiation use efficiency in field-grown maize and sorghum. *Crop Science* **34**: 721-727.
241. Mulvaney, R.L. and Bremner, J.M. 1981. Control of urea transformations in soils. In: *Soil Biochemistry*. pp. 153-196. E.A. Paul and J.N. Ladd (Eds). Marcel Dekker, Inc. New York, USA.
242. Nafziger, E.D.; Wax, L.M. and Brown, C.M. 1986. Response of five winter wheat cultivars to growth regulators and increased nitrogen. *Crop Science* **26**: 767-770.
243. Narang, R.S.; Singh, S. and Singh, S. 1985. Influence of irrigation and nitrogen management on the oil quality of Indian mustard (*Brassica juncea* Linn) Czern. & Coss. *Journal of Research Punjab Agricultural University* **22**: 229-234.
244. Naylor, R.E.L. and Stephen, N.H. 1993. Effects of nitrogen and the plant growth regulator chloromequat on grain size, nitrogen content and amino acid composition of triticale. *Journal of Agricultural Science* **120**: 159-169.
245. Neljubow, D. 1901. Uber die horizontale nutrition der stengel von *Pisum sativum* and einiger anderen. *Pflazen Beitrage and Botanik Zentralblatt* **10**: 128-139.
246. Nickell, L.G. 1982. Plant growth regulators: agricultural uses. Springer-Verlag, Berlin.
247. Nilsen, E.T. and Orcutt, D.N. 1996. The physiology of plants under stress: Abiotic factors. John Wiley and Sons Inc., New York.
248. Nonami, H.; Wu, Y. and Boyer, J.S. 1997. Decreased growth induced water potential. A primary cause of growth inhibition at low water potentials. *Plant Physiology* **114**: 501-509. h l
249. Novoa, R. and Loomis, R.S. 1981. Nitrogen and plant production. *Plant and Soil* **58**: 177-204.
250. O'Toole, P., McGarry, S.J. and Morgan, M.A. 1985. Ammonia volatilization from urea-treated pasture and tillage soils: effects of soil properties. *Journal of Soil Science* **36**: 613-620.
251. Oak, A. and Hirel, B. 1985. Nitrogen metabolism in roots. *Annual Review of Plant Physiology* **36**: 345-365.

- 252 Oritant, T and Yoshida, R 1971. Studies on nitrogen metabolism in crop plants XI The changes in abscisic acid and cytokinin like activity accompanying with growth and senescence in the crop plants *Proceedings of Crop Science Society of Japan* **40**: 325-331
- 253 Ortuno, A J.A , Del Rio, J L ; Casas, M ; Serrano, M.A. and Sanchez-Brava, J 1993 Influence of ACC and ethephon on cell growth in etiolated lupin hypocotyls dependence on cell growth stage *Biologia Plantarum* **33**: 81-90
- 254 Pallaghy, C K and Raschke, K 1972 No stomatal response to ethylene. *Plant Physiology* **49**: 275-276
- 255 Pallas Jr., J E and Kays, S.J. 1982 Inhibition of photosynthesis by ethylene a stomatal effect *Plant Physiology* **70**: 598-601
- 256 Palmer, R L , Lewis, L N , Hield, H Z and Kumamoto, J. 1967 Abscission induced by betahydroxyethylhydrazine: conversion of betahydroxyethylhydrazine to ethylene *Nature* **216**: 1216-1217.
- 257 Pando, S B and Srivastava, G C 1985 Physiological studies on seed set in sunflower III Significance of dwarfing the plant size using growth regulator. *Indian Journal of Plant Physiology* **28**: 72-80. 8 r
- 258 Pankovic, D., Sakac, Z , Kevresan, S and Plesnicar, M 1999. Acclimation to long-term water deficit in the leaves <sup>of</sup> two sunflower hybrids: photosynthesis, electron transport and carbon metabolism. *Journal of Experimental Botany* **50**: 127-138 2 of
- 259 Pannu, R K , Singh, D.P , Singh, P ; Sangwan, V.P. and Chaudhary, B D. 1992. Effect of moisture stress on growth partitioning of biomass and harvest index of oil seed brassicas *Crop Research Hisar* 5(supplement): 31-43
- 260 Panwar, J D.S ; Abbas, S , Ram, S and Sirohi, G.S. 1988. Effect of BAP, ethrel and stem girdling on growth and partitioning of photosynthates in Y-shaped mungbean *Indian Journal of Plant Physiology* **31**: 418-422.
- 261 Paparozzi, E.T.; Darrow, P.O.; McCallister, D.E. and Stroup, W.W. 1994. Effect of varying the nitrogen and sulphur supply on the following of poinsettia. *Journal of Plant Nutrition* **17**: 593-606. fl. ....
262. Parry, M. 1992. The potential effect on climate change on agriculture and land use *Advancement in Ecological Research* **22**: 63-91.

263. Pate, J.S. 1980. Transport and partitioning of nitrogenous solutes. *Annual Review of Plant Physiology* **31**: 313-340.
264. Patil, B.N. and Bhargava, S.C. 1987. Seed quality studies in rapeseed mustard in relation to nitrogen nutrition. *Annals of Plant Physiology* **1**: 81-87.
265. Patra, A.K.; Tripathy, S.K. and Samui, R.C. 1995. Effect of post-flowering foliar application of nutrients on growth, yield and economics of rainfed groundnut (*Arachis Hypogaea* L.). *Indian Journal of Plant Physiology* **38**: 203-206.
266. Patrick, J.W. 1982. Hormonal control<sup>#</sup> of assimilate transport. In: *Plant Growth Substances*. pp. 669. P.F. Wareing (Ed). Academic Press, New York.
267. Patrick, J.W. and Steains, K.H. 1987. Auxin promoted transport of metabolites in stem of *Phaseolus vulgaris*<sup>15</sup>; auxin dose response curves and effect of inhibitors of polar auxin transport. *Journal of Experimental Botany* **38**: 203-210. ?
268. Paul, E.A. and Clark, F.E. 1988. Soil microbiology and biochemistry. Academic Press, New York.
269. Paulpandi, V.K.; Paleniappan, S.P. and Solaiappan, U. 1998. Effect of cycocel on different crop geometry and nutrient levels in sorghum <sup>Se</sup> (*Sorghum bicolor*). *Journal of Agronomy* **43**: 524-527.
270. Payne, W.A.; Drew, M.C.; Hossner, L.R.; Lascano, R.J.; Onken, A.B. and Wendt, C.W. 1992. Soil phosphorus availability and pearl millet water-use efficiency. *Crop Science* **32**: 1010-1015.
271. Pearson, C.H.; Golus, H.M. and Tindall, T.A. 1989. Ethaphon<sup>e</sup> application and nitrogen fertilization of irrigated winter barley in an arid environment. *Agronomy Journal* **81**: 717-719.
272. Pechan, P.M. and Morgan, D.G. 1983. The use of radiography in studies of plant development *in vivo*. *Planta* **159**: 476-481.
273. Pereto, J.G. and Beltran, J.P. 1987. Hormone directed sucrose transport during fruit set induced by gibberellins in *Pisum sativum*. *Physiologia Plantarum* **69**: 356-360.

274. Perumal, N.K. 1999. Effect of different nitrogen levels on morpho-physiological characters and yield in rainfed cotton. *Indian Journal of Plant Physiology* **4**: 65-67.
275. Pilbeam, D.J. and Kirkby, E.A. 1992. Some aspects of the utilization of nitrate and ammonium by plants. In: *Nitrogen Metabolism of Plants*. pp. 55-70 K. Mengel and D.J. Pilbeam (Eds.). Clarendon Press, Oxford, U.K.
276. Pincus, G. and Thimann, K.C. 1948. The Hormones. Academic Press, New York.
277. Pinkerton, A. 1991. Critical phosphorus concentration in oilseed rape (*Brassica napus*) and Indian mustard (*Brassica juncea*) as affected by nitrogen and plant age. *Australian Journal of Experimental Agriculture* **31**: 107-115.
278. Platt, S.G.; Plaut, Z. and Bascham, J.A. 1977. Ammonia regulation of carbon metabolism in photosynthesizing alfalfa leaf discs. *Plant Physiology* **60**: 739-742.
279. Polley, H.W.; Johnson, H.B.; Tischer, C.R. and Torbert, H.A. 1999. Links between transpiration and plant nitrogen: vibration with atmospheric CO<sub>2</sub> concentration and nitrogen availability. *International Journal of Plant Science* **160**: 535-542.
280. Post-Beittenmiller, D., Roughan, G. and Ohlrogge, J. 1992. Regulation of plant fatty acid biosynthesis. *Plant Physiology* **100**: 923-930.
281. Pradhan, A.C.; Sarkar, S.K. and Roy, S.K. 1994. Growth and yield of rape seed-mustard varieties as influenced by nitrogen and phosphorus fertilization. *Environment and Ecology* **12**: 166-170.
282. Prasad, N.K.; Kumar, S.; Kumar, P. and Singh, R.S. 2000. Components of nutrients-use efficiency as influenced by legume-wheat (*Triticum aestivum*) sequences. *Indian Journal of Agricultural Sciences* **70**: 503-506.
283. Prasad, S. and Shukla, D.N. 1992. Accumulation of nitrogen, phosphorus and potassium as affected by N, K<sub>2</sub>O and CCC interaction in mustard. *Annals of Agriculture Research* **13**: 25-31.
284. Prasad, S. and Shukla, D.N. 1993. Effect of interactions of nitrogen, potassium and cycocel on growth characters in relation to grain yield of



- mustard (*Brassica juncea* L.) var. T<sub>59</sub>. *Indian Journal of Agricultural Research* **27**: 13-20.
285. Pregtizer, K.S.; Dickmann, D.I.; Hendrick, R. and Nguyen, P.V. 1990. Whole-tree carbon and nitrogen partitioning in young hybrid poplars. *Tree Physiology* **7**: 79-93.
  286. Pua, E.C. and Chi, G.L. 1993. De novo shoot morphogenesis and plant growth of mustard (*Brassica juncea*) in vitro in relation to ethylene. *Physiologia Plantarum* **88**: 467-474.
  287. Raghuvanshi, R.K.S.; Paradhar, V.K.; Gupta, V.K. and Jain, S.C. 1987. Evaluation of irrigation and N fertilization requirement for linseed grown on a sodic clay soil. *Indian Journal of Agricultural Science* **57**: 479-482.
  288. Rajala, A. and Peltonen-Sainio, P. 2001. Plant growth regulator effects on spring cereal root and shoot growth. *Agronomy Journal* **93**: 936-943.
  289. Ramans, M. 1989. Double low oilseed rape in U.K. exharvest 1988. In: *Proceedings of Double-low Oilseed Rape for the 1990s Conference*. pp. 16-53. C. Green and G. Beamont (Eds.). Peterborough, U.K.
  290. Ramanujan, T. and Mohan Naidu, K. 1995. Light interception and energy conservation in sugarcane genotypes. *Indian Journal of Plant Physiology* **38**: 189-192.
  291. Ramos, J.M.; Garcial de Moral, L.F.; Molina-cano, J.L.; Salamanca, P. and Roca de Togores, F. 1989. Effect of an early application of sulphur or ethephon as foliar sprays on the growth and yield of spring barley in a Mediterranean environment. *Agronomy and Crop Science* **163**: 129-137.
  292. Rana, D.S.; Singh, M.P. and Ahlawat, I.P.S. 1991. Oil yield and nitrogen uptake in mustard (*Brassica juncea*) as affected by irrigation, plant density and nitrogen application. *Indian Journal of Agronomy* **36**: 143-146.
  293. Raskin, I. and Kende, H. 1984. The role of gibberellin in the growth response of submerged deep water rice. *Plant Physiology* **76**: 947-950.
  294. Rathore, P.S. and Manohar, S.S. 1989. Response of mustard to nitrogen and sulphur. I. Effects of nitrogen and sulphur on ancillary characters and yield of mustard. *Indian Journal of Agronomy* **34**: 333-336.

295. Reddy, B.N. and Sinha, M.N. 1988. Yield and phosphorus utilization of mustard as influenced by phosphorus at various levels of nitrogen and moisture regimes. *Fertilizer News* **33**: 31-37.
296. Reddy, B.S.; Reddy, S.R. and Subbaih, G. 1991. Effect of supplemental nutrition during post flowering phase of groundnut. *The Andhra Agricultural Journal* **38**: 4-7.
297. Reddy, D.S. 1983. Response of linseed varieties to varying fertility levels under irrigation. *Mysore Agricultural Science* **17**: 91.
298. Reddy, G.K. and Reddy, K.A. 1986. Uptake and apparent recovery of nitrogen by rice as influenced by nitrogen levels and forms of urea. *Journal Research of A.P.A.U.* **14**: 106-110.
299. Reddy, K.R. and Patrick, W.H. 1986. Denitrification losses in flooded rice fields. In: *Nitrogen Economy of Flooded Rice Soils*. pp. 99-116. S.K. De Datta and W.H. Patrick Jr. (Eds). Martinus Nijhoff, Dordrecht, Netherlands.
300. Reddy, S.N.; Singh, B.G. and Reddy, R.V. 1996. Effect of sulphur and benzyladenine on quality and nutrient uptake in sunflower. *Annals of Plant Physiology* **10**: 166-170.
301. Reid, M.S. 1987. Ethylene in plant growth, development, and senescence. In: *Plant Hormones and their Role in Plant Growth and Development*. pp. 257-279. P.J. Davies (Ed). Martinus Nijhoff Publishers, Dordrecht, The Netherlands.
302. Richards, W.J. IV, Bender, L. and Thorpe, T.A. 1994. Nitrogen metabolisms in cultured cotyledon explants of *Pinus radiata* during de novo organogenesis. *Physiologia Plantarum* **92**: 681-688.
303. Ridge, I. and Osborne, D.J. 1971. Role of peroxidase when proline-rich protein in plant cell walls is increased by ethylene. *Nature New Biology* **229**: 205-208.
304. Ries, S.K., Wert, V., Sweely, C. and Leavitt, R.A. 1977. Triacntanol regulator. *Science* **195**: 1331-1341.
305. Rigaud, J. 1981. Comparison of the efficiency of nitrate and nitrogen fixation in crop yield. In: *Nitrogen and Carbon Metabolism*. pp. 17-48. J.D. Bewlly and Martinus Nijhoff (Eds.). W. Junk Publishers. The Hague-Boston-London.

306. Robertson, .A. and Kuenen, J.G. 1991. Physiology of nitrifying and denitrifying bacteria. In: *Microbial production and Consumption of Greenhouse Gases*. pp. 189-199. J.E. Rogers and Whitman, W.B. (Eds). American Society for Microbiology, Washington DC.
307. Robertson, G.P. 1982. Factors regulating nitrification in primary and secondary succession. *Ecology* **63**: 1561-1573.
308. Rodriguez-Pousida, R.A.; De Rycke, R.; Dedondar, A.; Van Caeneghem, W.; Engler, E.; Van Montagu, M. and Van Der Straeten, D. 1993. The arabidopsis 1-amino-cyclopropane-1-carboxylate synthase gene 1 is expressed during early development. *Plant Cell* **5**: 897-911.
309. Rogers, G.S.; Milham, P.J.; Grillings, M. and Conroy, J.P. 1996. Sink strength may be the key to growth and nitrogen responses in N-deficient wheat at elevated CO<sub>2</sub>. *Australian Journal of Plant Physiology* **23**: 253-264.
310. Rood, S.B. and Major, D.J. 1984. Influence of plant density, nitrogen, water supply and pod leaf removal on growth of oilseed rape. *Field Crop Research* **8**: 323-332.
311. Roth-Bejerano, N. and Lips, S.H. 1970. Hormonal regulation of nitrate reductase activity in leaves. *New Phytologist* **69**: 165-169.
312. Ryan, J., Curtain, D. and Safi, I. 1981. Ammonia volatilization as influenced by calcium carbonate particle size and iron oxides. *Soil Science Society American Journal* **45**: 338-341.
313. Sage, R.F.; Pearey, R.W. and Seemann, J.R. 1987. The nitrogen efficiency of C<sub>3</sub> and C<sub>4</sub> plants III: Leaf nitrogen effect on activity of carboxylating enzymes in *Chenopodium album* L. and *Amaranthus retroflexus* L. *Plant Physiology* **85**: 355-359.
314. Sage, R.F.; Sharky, T.D. and Pearcy, R.W. 1990. The effect of leaf nitrogen and temperature on the CO<sub>2</sub> response of photosynthesis in the C<sub>3</sub> dicot *Chenopodium album* L. *Australian Journal of Plant Physiology* **17**: 135-148.
315. Sahrawat, K.L. 1978. Nitrification inhibitors for efficient use of fertilizer nitrogen: A scheme for development of nitrification inhibitors. In: *Plant Nutrition*. pp. 431-438. A.R. Ferguson, R.L. Bielecki and I.R. Ferguson (Eds). *Proceedings of 8<sup>th</sup> Int. Colloq. Plant Analysis and Fert. Problems*, Auckland, New Zealand, 28 Aug-1 Sept., 1978. DSIR Inform Series 134, Govt. Printer, Wellington, New Zealand.

316. Sahrawat, K.L. 1980a. Control of urea hydrolysis and nitrification in soils by chemicals-prospects and problems. *Plant Soil*. 57: 335-352. *and*
317. Sahrawat, K.L. 1980b. On the criteria for comparing the ability of compounds for retardation of nitrification in soil. *Plant Soil* 55: 487-490. *and*
318. Sahrawat, K.L. 1989. Effects of nitrification inhibitors on nitrogen transformations, other than nitrification, in soils. *Advances in Agronomy* 42: 279-309.
319. Sahrawat, K.L. 1996. Nitrification, inhibitors with emphasis on natural products and persistence of fertilizer nitrogen in the soil. In: *Nitrogen Economy in Tropical Soils*. pp. 379-388. N. Ahmad (Ed.). Kluwer Academic Publishers, Netherlands.
320. Sahrawat, K.L. and Keeney, D.R. 1984. Effects of nitrification inhibitors on chemical composition of plants: A review. *Journal of Plant Nutrition* 7: 1251-1288.
321. Saini, J.S.; Jolly, R.S. and Dhillon, A.S. 1984. Effect of some growth regulators on the pod yield and quality in peanut (*Arachis hypogaea*). *Crop Improvement* 11: 61-63.
322. Sairam, R.K.; Deshmukh, P.S. and Shukla, D.S. 1991. Influence of chloromequate chloride on photosynthesis and nitrate assimilation in wheat genotypes under water stress. *Indian Journal of Plant Physiology* 34: 222-227.
323. Sairam, R.K.; Deshmukh, P.S.; Shukla, D.S.; Wasnik, K.G. and Kushwaha, S.R. 1989. Effect of abscisic acid and triadimefon on photosynthesis and nitrate reductase activity during water stress in wheat. *Indian Journal of Physiology* 32: 51-56. *Plant*
324. Salama, A.M. and Bazas, I. 1987. Effect of growth regulators and nitrogen, phosphorus, potassium fertilizers on the trace element contents of sunflower in calcareous soils. *Acta Agronomica Hungarica* 36: 37-42.
325. Salisbury, F.B. and Ross, C.W. 1994. *Plant Physiology*. 4<sup>th</sup> ed. Wadsworth Publishing Company, Belmont, C.A.
326. Salsac, L.; Chaillou, S; Morot-Gaurdy, J.F. and Lesaint, C. 1987. Nitrate and ammonium in plants. *Plant Physiology and Biochemistry* 25: 805-812.

327. Samimy, C. 1978. Influence of cobalt on soybean hypocotyl growth and its ethylene evolution. *Plant Physiology* **62**: 1005-1006.
328. Samiullah, Khan, N.A., Ansari, S.A. and Afridi, M.M.R.K. 1991. Pyridoxine augment growth, yield and quality of mustard through efficient utilization of soil applied NP fertilizers. *Acta Agronomica Hungarica* **40**: 111-116.
329. Samiullah; Khan, N.A.; Afridi, M.M.R.K. and Ansari, S.A. 1990. Response of ten varieties of mustard (*Brassica juncea* L. Czern & Coss.) to nitrogen and phosphorus Plant dry weight and NAR. *Bangladesh Journal of Botany* **19**: 83-86.
330. Saniewske, M. and Ludhika, K. 1989. Silver thiosulphate on tulip stem of the inhibitory effect of ethephon on tulip stem elongation induced by auxin. *Bulletin Polish Academic Sciences* **36**: 265-270.
331. Santakumari, M. and Berkowitz, G.A. 1990. Correlation between the maintenance of photosynthesis and situ protoplast volume at low water potentials in droughted wheat. *Plant Physiology* **92**: 733-739.
332. Sanvicente, P.; Lozarevitch, S.; Blouet, A. and Guckert, A. 1999. Morphological and anatomical modification in winter barley culm after late plant growth regulator treatment. *European Journal of Agronomy* **11**: 45-51.
333. Sanvicente, P.; Lozarevitch, S.; Blouet, A. and Guckert, A. 1999. Morphological and anatomical modification in winter barley culm after late plant growth regulator treatment. *European Journal of Agronomy* **11**: 45-51. repetition
334. Saran, G. and Giri, G. 1990. Influence of nitrogen, phosphorus and sulphur on mustard under semi-arid rain fed conditions of North-west India. *Indian Journal of Agronomy* **35**: 131-136.
335. Sardadevi, K.; Padamsree, K. and Raghavendra, A.S. 1996. Correlation between the inhibition of photosynthesis and the decrease in area of detached leaf discs or volume/absorbance of protoplasts under osmotic stress in pea (*Pisum sativum*). *Physiologia Plantarum* **96**: 395-400.
336. Sarkar, R.K.; Saha, A. and Chakraborty, A. 1999. Analysis of growth and productivity of sunflower (*Helianthus annuus*) in relation to crop geometry, nitrogen and sulphur. *Indian Journal of Plant Physiology* **4**: 28-31.

- 337 Satyavan, K S ; Panwar, S.; Punia, S. and Dhindwal, A.S. 1999. Dry matter production, N uptake and oil content in Indian rape (*Brassica campestris* L.) genotypes as influenced by nitrogen levels and plant densities *Annals of Biology (Hissar)* **15**: 63-65.
- 338 Sauerbrey, E , Grossmann, K. and Jung, J. 1988. Ethylene production by sunflower cell suspension effects of plant growth retardants. *Plant Physiology* **87**: 510
- 339 Sawage, L J and Ohlrogge, J.B. 1999. Phosphorylation of pea chloroplast acetyl-CoA carboxylase. *Plant Journal* **18**: 521-527.
- 340 Sawan, Z M , Gregg, B.R. and Yousef, S.E. 1998. Influence of nitrogen fertilization and foliar applied growth retardants and zinc on cotton seed yield, viability and seedling vigour. *Seed Science and Technology* **26**: 393-404
- 341 Sawhney, V ; Singal, H.R.; Singh, P.; Sawhney, S.K. and Singh, D.P. 1996. Pattern of plant relations and carbon dioxide exchange rates in contrasting genotypes of *Brassica juncea* under water deficit conditions. *Indian Journal of Plant Physiology* **1**: 203-206.
342. Schmidt, E.L. 1982. Nitrification in soil. In: *Nitrogen in agricultural soils*. pp. 252-288. F.J Stevenson, (Ed). American Society of Agronomy, Madison, WI.
- 343 Sekhon, N.K and Singh, G. 1994. Effect of growth regulators and date of sowing on grain development in wheat. *Indian Journal of Plant Physiology* **37**: 1-4.
- 344 Shanahan, J.F. and Nielsen, D.C. 1987. Influence of growth retardants (anti-gibberellins) on corn vegetative growth, water use and grain yield under different levels of water stress. *Agronomy Journal* **79**: 103-109.
- 345 Shand, C.R. 1996. Response of sugarcane to nitrogen fertilization practices on some sugarcane soils in Trinidad. In: *Nitrogen Economy in Tropical Soils*. pp. 399-413. N. Ahmad (Ed.). Kluwer Academic Publishers, Netherlands.
346. Sharma, A.K.; Sharma, A.M. and Sharma, Y.M. 1994. Effect of irrigation, nitrogen and sulphur application on seed yield, quality and sulphur uptake by Indian mustard (*Brassica juncea*). *Agricultural Science Digest (Karnal)* **14**: 63-67.

347. Sharma, D.K. 1992. Physiological analysis of yield variations in mustard varieties under water stress and non-water stress conditions. *Annals of Agricultural Research* **13**: 174-176.
348. Sharma, J.P. 1994. Response of Indian mustard (*Brassica juncea*) to different irrigation schedules and nitrogen and sulphur levels. *Indian Journal of Agronomy* **39**: 421-425.
349. Sharma, R.; Grewal, M.K. and Singh, G. 1986. Effect of ethrel on lipid metabolism in soybean germinating under moisture stress. *Indian Journal of Plant Physiology* **29**: 207-210.
350. Sharma, R.; Sharma, B. and Singh, G. 1982. Effect of growth regulators on some biochemical changes during pod development in ground-nut (*Arachis hypogaea*). *Indian Journal of Botany* **5**: 102-106.
351. Sharpe, R.R. and Harper, L.A. 1995. Soil, plant and atmospheric conditions as they relate to ammonia volatilization. *Fertilizer Research* **42**: 149-158.
352. Shelp, B.J. 1987. Plant characteristics and nutrient composition and mobility of broccoli (*Brassica oleracea* var. *italica*) supplied with  $\text{NH}_4^+$ ,  $\text{NO}_3^-$  or  $\text{NH}_4\text{NO}_3$ . *Journal of Experimental Botany* **38**: 1603-1618.
353. Shibaoka, H. 1994. Plant hormone-induced changes in the orientation of cortical microtubules: alterations in the crosslinking between microtubules and plasma membrane. *Annual Review of Plant Physiology* **45**: 527-544.
354. Shukla, A. and Kumar, A. 1994. Effect of some Indian mustard varieties and nitrogen level on growth, yield and quality. *Annals of Agricultural Research* **15**: 76-80.
355. Silk, W.K. 1992. Steady form from changing cells. *International Journal of Plant Science* **153**: 49-58.
356. Simmons, S.R.; Orlka, E.A.; Wiersma, J.V.; Lueschen, W.E. and Warners, D.D. 1988. Spring wheat and barley responses to ethephon. *Agronomy Journal* **80**: 829-834.
357. Sinclair, T.R. 1990. Nitrogen influence on the physiology of crop yield. In: *Theoretical Production Ecology: Reflections and Prospects*. R. Rabbinge; J. Gourdiaan; H. van Keulen; F.W.T. Penning de Vries and H.H. van Laar (Eds.). Pudoc, Wageningen.

358. Sinclair, T.R. and Horie, T. 1989. Leaf nitrogen, photosynthesis and crop radiation use efficiency: A review. *Crop Science* **29**: 90-98.
359. Singh, B and Kumar, V. 1991. Response of Indian mustard (*Brassica juncea*) to nitrogen and sulphur application under rain fed condition. *Indian Journal of Agronomy* **41**: 286-289.
360. Singh, B.; Singh, B.P.; Faroda, A.S. and Gupta, S.K. 1994. Effect of irrigation and nitrogen levels on the quality and oil yield of Brassica species. *Indian Journal of Agronomy* **39**: 362-365.
361. Singh, H.; Chandra, S.; Jolly, R.S. and Kolar, I.S. 1987. Influence of chlorocholine chloride and ethrel on the grain yield and uptake of nitrogen, phosphorus and potassium in soybean. *Crop Improvement* **14**: 79-83.
362. Singh, J.; Nandwal, A.S.; Kuhad, M.S.; Verma, S.K. and Singh, J. 1996. Studies on CO<sub>2</sub> exchange and plant water status under consecutive occurrence of salt and water stress. *Annals of Biology* **12**: 71-76.
363. Singh, O.P. and Srivastava, R.D.L. 1996. Utilization and response to fertilizer N and productive efficiency of wheat grown under varying levels of nitrogen. *Indian Journal of Plant Physiology* **1**: 210-211.
364. Singh, R.B. and Mishra, P.H. 1994. Effect of nitrogen and sulphur on uptake and dry matter accumulation in linseed (*Linum usitatissimum* L.). *Indian Journal of Plant Physiology* **37**: 104-108.
365. Singh, R.P. and Kumar, A. 1991. Effect of phytohormones on yield attributes and seed yield of mustard (*Brassica juncea*). *Indian Journal of Agronomy* **36**: 379-381.
366. Singh, R.P.; Yazdani, S.S.; Verma, G.D. and Singh, V.N. 1995. Effect of different levels of nitrogen, phosphorus and potash on aphid infestation and yield of mustard. *Indian Journal of Entomology* **57**: 18-21.
367. Singh, S. 1996. Physiological effects of growth <sup>promoters</sup> ~~parameter~~ (GA<sub>3</sub>) and retardants (CCC) in diverse rice genotypes. *Plant Physiology and Biochemistry* **23**: 153-158.
368. Singh, S. 1997. Growth and yield response of different crop species to low light and high temperature-humidity stress. *Indian Journal of Plant Physiology* **2**: 151-155.



369. Singh, S. and Saran, G. 1987. Effect of sulphur and nitrogen on growth yield, quality and nutrient uptake of Indian rape. *Indian Journal of Agronomy* **32**: 474-475.
370. Singh, S.; Singh, K. and Singh, S.P. 1995. Effect of hormones on growth and yield characters of seed crop of kharif onion (*Allium cepa* L.). *Indian Journal of Plant Physiology* **38**: 193-196.
371. Singh, S.R.P. and Bansal, K.N. 1999. Effect of different organic manures and nitrogen levels on growth and yield of sunflower (*Helianthus annuus*). *Indian Journal of Agricultural Sciences* **69**: 8-9.
372. Sinha, S.K. 1992. Climatic change and agriculture. *Indian Farming* **5**: 13-18.
373. Sircar, S.M. 1971. Plant hormone research in India. IARI, New Delhi.
374. Sisler, E.C. Yang, S.F. 1984. Anti-ethylene effects of *cis*-2-butene and cyclic olefins. *Phytochemistry* **23**: 2765-2768.
375. Sisler, E.C.; Dupille, E. and Serek, M. 1996a. Effect of 1-methylecyclopropene and methylenecyclopropene on ethylene binding and ethylene action on cut carnations. *Plant Growth Regulators* **18**: 79-86.
376. Sisler, E.C.; Serek, M. and Dupille, E. 1996b. Comparison of cyclopropene, 1-methylecyclopropene and 3,3-dimethylecyclopropene as ethylene antagonists in plants. *Plant Growth Regulators* **18**: 169-174.
377. Sivasankar, S. and Oak, A. 1996. Nitrate assimilation in higher plants: the effect of metabolites and light. *Plant Physiology and Biochemistry* **34**: 609-620.
378. Slife, F.W. and Earley, E.B. 1970. Effect of ethrel on growth and yield of soybean. *Agronomy Journal* **62**: 434-435.
379. Smalle, J.; Heagman, M.; Kurepa, J.; Van Montagu, M. and Van Der Straden, D. 1997. Ethylene can stimulate arabidopsis hypocotyls elongation in light. *Proceedings of the National Academy of Sciences of the United States of America*. **94**: 2756-2761.
380. Smith, C.J.; Wright, G.C. and Woodroffe, M.R. 1988. The effect of irrigation and nitrogen fertilizer on rapeseed (*Brassica napus*) production in south-eastern Australia. II. Nitrogen accumulation and oil yield. *Irrigation Science* **9**: 15-25.

- 381 Solomonson, L.P. and Barber, M.J. 1990. Assimilatory nitrate reductase: functional properties and regulation. *Annual Review of Plant Physiology and Plant Molecular Biology* **41**: 225-253.
382. Soper, R.J., Raez, G.J. and Fehr, P.I. 1971. Nitrate nitrogen in the soil as a means of predicting the nitrogen fertilizer requirements of barley. *Canadian Journal of Soil Sciences* **51**: 45-49.
- 383 Sowers, K.E.; Pan, W.L.; Miller, B.C. and Smith, J.L. 1994. Nitrogen use efficiency of split nitrogen application in soft white winter wheat. *Agronomy Journal* **86**: 942-948.
384. Stecker, J.A.; Buchholz, D.D.; Hanson, R.G.; Wollenhaupt, N.C. and McVay, K.A. 1993. Broadcast nitrogen sources for no-till continuous corn and corn following soybean. *Agronomy Journal* **85**: 893-897.
385. Steer, B., Hocking, P.J., Kortt, A.A. and Roxburgh, C.M. 1984. Nitrogen nutrition of sunflower (*Helianthus annuus* L.): yield components, the timing of their establishment and seed characteristics in response to nitrogen supply. *Field Crops Research* **9**: 219-236. / n
- 386 Steer, B.T.; Coaldrake, P.D.; Pearson, C.J. and Canty, P.J. 1986. Effects of nitrogen supply and population density on plant development and yield components of irrigated sunflower (*Helianthus annuus* L.). *Field Crops Research* **13**: 99-115
387. Subrahmanyam, D. and Rathore, V.S. 1992a. Influence of ethylene on carbon-14 labeled carbondioxide assimilation and partitioning in mustard. *Plant Physiology and Biochemistry* **30**: 81-86.
388. Subrahmanyam, D. and Rathore, V.S. 1992b. Plant growth regulators influence <sup>14</sup>CO<sub>2</sub> assimilation and translocation of assimilates in Indian mustard. *Journal of Agronomy and Crop Science* **168**: 145-152. / n
389. Sugiharto, B.; Burnell, J.N. and Sugiyama. 1992. Cytokinin is required to induce the nitrogen dependant accumulation of RNAs for phosphoenolpyruvate carboxylase and carbonic anhydrase in detached maize leaves. *Plant Physiology* **100**: 153-156.
390. Sundqvist, C.; Bjorn, L.O. and Virgin, H.I. 1980. Factors in chloroplast differentiation. In: *Chloroplast*. pp. 201-224. J. Reinert (Ed.). Springer-Verlag, Berlin.
391. Taiz, L. 1984. Plant cell expansion: regulation of cell wall mechanical properties. *Annual Review of Plant Physiology* **35**: 585-657.

392. Tandon, H.L.S. 1986. Sulphur research and agricultural production in India. 2<sup>nd</sup> Ed. Fertilizer Development and Consultation Organisation, New Delhi, 76.
393. Tanguilig, V.C.; Yambao, E.B.; O'Toole, J.C. and De Datta, S.K. 1987. Water stress effects on leaf elongation, leaf water potential, transpiration and nutrient uptake of rice, maize and soybean. *Plant and Soil* **103**: 155-168.
394. Tate, R.L. 1992. Soil organic matter: Biological and ecological effects. Krieger Publishing Company, Malabor, Florida, USA.
395. Tayo, T.O. and Morgan, D.G. 1978. Factors influencing flower and pod development in oilseed rape (*Brassica napus* L.). *Journal of Agricultural Science Cambridge* **92**: 363-373.
396. Terman, G.L. 1979. Volatilization losses of nitrogen as ammonia from surface-applied fertilizers, organic amendments and crop residues. *Advances in Agronomy* **31**: 189-223.
397. Thakral, S.K.; Singh, B.P. and Faroda, A.S. 1997. Consumptive use, water use efficiency and soil moisture extraction pattern as influenced by irrigation levels, fertility levels and Brassica species. *Indian Journal of Agricultural Research* **31**: 65-70.
398. Thakral, S.K.; Singh, B.P.; Faroda, A.S. and Gupta, S.K. 1996. Effect of irrigation and fertility levels on the oil yield and quality of Brassica species. *Annals of Agricultural Research* **17**: 416-418.
399. Thakral, S.K.; Singh, B.P.; Faroda, A.S.; Gupta, S.K. and Kumar, S. 1995. Fatty-acid composition in seeds of Brassica species as affected by moisture and fertility levels. *Crop Research (Hissar)* **10**: 137-140.
400. Thakuria, K. and Gogoi, R.K. 1996. Response of rainfed Indian mustard (*Brassica juncea*) to nitrogen and row spacing. *Indian Journal of Agronomy* **41**: 279-281.
401. Thomas, T.H. 1986. Hormonal control of assimilate movement and compartmentation. In: *Plant Growth Substances*. pp. 350-359. M. Bopp (Ed.). Springer-Verlag, Heidelberg.
402. Thorsteinssen, B. and Eliasson, L. 1990. Growth retardation induced by nutritional deficiency or abscisic acid in *Lemna gibba*. The relationship of growth rate and endogenous cytokinin content. *Plant Growth Regulation* **9**: 171-181.

403. Tiedje, J.M. 1988. Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In: *Biology of Anaerobic Microorganisms*. pp. 179-244. A.J.B. Zehnder (Ed.). John Wiley & Sons, New York. A e
404. Tisdale, S.L. and Nelson, W.L. 1967. Soil fertility and fertilizers. The Macmillan Co., New York.
405. Tisdale, S.L., Nelson, W.L. and Beaton, J.D. 1985. Soil fertility – past and present. In: *Soil Fertility and Fertilizers*. pp. 5-18. 4<sup>th</sup> ed. Macmillan, New York.
406. Tomar, J.S. and Soper, R.J. 1981. Fate of tagged urea N in the field with different methods of N and organic matter placement. *Agronomy Journal* 73: 991-995.
407. Tomar, R.K.S.; Chourasia, S.C.; Raghu, J.S. and Singh, V.B. 1996. Growth, yield and net returns of mustard under different levels of nitrogen and sulphur application on clay loam soils. *Journal of Oilseed Research* 13: 13-17.
408. Tomar, T.S.; Singh, Subey; Kumar, Sandeep; Tomar, Savita; Singh, S.; Kumar, S. and Tomar, S. 1997. Response of Indian mustard (*Brassica juncea*) to nitrogen phosphorus and sulphur fertilization. *Indian Journal of Agronomy* 42: 148-151.
409. Towes, W.H. and Soper, R.J. 1978. Effect of nitrogen sources, method of placement and soil type on seedling emergence and barley crop yield. *Canadian Journal of Soil Science* 58: 311-320. of
410. Tripathi, H.P. and Singh, H.R. 1992. Nutrient uptake and quality of mustard varieties (*Brassica juncea* L. Czern & Coss) as influenced by nitrogen and row spacing. *Narendra Deva Journal of Agricultural Research* 7: 185-187.
411. Tudela, D. and Primo Millo, E. 1992. 1-aminocyclopropene-1-carboxylic acid transported from roots to shoots promotes leaf abscission in Cleopatra mandarin (*Citrus rashni* Hort. Extan.) seedlings rehydrated after water stress. *Plant Physiology* 100: 131-137.
412. Uhart, S.A. and Andrade, F.H. 1995a. Nitrogen deficiency in maize. I. Effects on crop growth, development, dry matter partition and kernel set. *Crop Science* 35: 1376-1383.

413. Uhart, S.A. and Andrade, F.H. 1995b. Nitrogen deficiency in maize: II. carbon-nitrogen interaction effects on kernel number and grain yield. *Crop Science* **35**: 1384-1389.
414. Urwiller, M.J. and Stuttle, C.A. 1986. Influence of ethaphon on soybean reproductive development. *Crop Science* **26**: 976-979.
415. Van der Werf, A.; Nuenen, M. van; Visser, A.J. and Lambers, H. 1993. Effects of N supply on the rates of photosynthesis and shoot and root respiration of inherently fast and slow growing monocotyledonous species. *Physiologia Plantarum* **89**: 563-569.
416. Van Quy, L; Lamaze, T and Champigny, M.L. 1991. Short-term effects of nitrate on sucrose synthesis in wheat leaves. *Planta* **185**: 53-57.
417. Van Sanford, D.A.; Grove, J.H.; Grabau, J. and MacKown, C.T. 1989. Ethephon and nitrogen use in winter wheat. *Agronomy Journal* **81**: 951-954.
418. Vance, C.P. 1990. Simbiotic nitrogen fixation. Recent genetics advances. In: *The Biochemistry of Plants. Intermediary Nitrogen Metabolism*. pp. 43-88. Vol. 16. B.J. Mifflin and P.J. Lea (Eds.). Academic Press, San Diego, CA.
419. Vander Boogard, R. Alwynse, D., Veneklaas, E.J. and Lambers, S.H. 1996. The association of biomass allocation with growth and water use efficiency of two *Triticum aestivum* cultivars. *Australian Journal of Plant Physiology* **23**: 751-761.
420. Vander Boogard, R., Kostadirova, S., Veneklaas, E.J. and Lambess, H. 1995. Association of water use efficiency and nitrogen use efficiency with photosynthetic characteristics of two wheat cultivars. *Journal of Experimental Botany* **46**: 1429-1438.
421. Varade, P.B.; Wasnik, K.G. and Bagga, A.K. 1995. Effect of low nitrate applied at preflowering stage on nodular <sup>PEP</sup> ~~pep~~ carboxylase activity in soybean (*Glycine max* L. Merr). *Indian Journal of Plant Physiology* **38**: 148-150. PEP
422. Veen, H. 1986. A theoretical model for anti-ethylene effects of silver thiosulphate and 2,5-norbornadiene. *Acta Horticulture* **181**: 129-134.
423. Vioque, B. and Castellano, J.M. 1994. Extraction and biochemical characterization of 1-aminocyclopropane-1-carboxylic acid oxidase from pear. *Physiologia Plantarum* **90**: 334-338. AO

424. Vlek, P.L.G. and Byrnes, B.H. 1986. The efficiency and loss of fertilizer N I lowland rice. *Fertilizer Research* **9**: 131-147.
425. Von Cammerer, S., Farquhar, G.D. 1981. Some relationship between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta* **153**: 376-387.
426. Vreugdenhil, D. and Harro, J.B. 1989. Effects of ethylene on tuberization in radish (*Raphanus sativus*). *Plant Growth Regulators* **8**: 21-30.
427. Vyas, S.P.; Garg, B.K.; Kathju, S. and Lahiri, A.N. 1995. Influence of nitrogen on Indian mustard grown under different levels of stored soil moisture. *Journal of Arid Environment* **29**: 173-184.
428. Vyas, S.P.; Kathju, S.; Garg, B.K. and Lahiri, A.N. 1999. Influence of supplemental irrigation and urea application on productivity and nitrogen metabolism of sesame. *Indian Journal of Plant Physiology* **4**: 197-201.
429. Wareing, P.F. and Phillips, I.D.J. 1981. The control of growth and differentiation in plants. Pergaman Press, New York.
430. Warner, H.L. and Leopold, A.C. 1969. Ethylene evolution from 2-chloroethyl phosphonic acid. *Plant Physiology* **44**: 156-158.
431. Watson, D.J. 1958. The dependence of net assimilation rate of leaf area index. *Annals of Botany* **22**: 37-54.
432. Westerman, R.L. and Toker, T.C. 1987. Soil fertility concepts: past, present and future. In: *Future Developments in Soil Science Research*. pp. 171-179. Soil Sci. Soc. Am., Madison, Wisconsin.
433. Whenham, R.J., Burn, I.G., Stone, D.A. and Fraser, R.S.S. 1989. Effect of nitrogen nutrition and water regime on abscisic acid, phaseic and dihydrophaseic acid metabolism in leaves of field grown kale (*Brassica oleracea*): Consequences for plant growth and crop yield. *Journal of Science Food and Agriculture* **49**: 143-155.
434. Wiersma, D.W., Oplinger, E.S. and Guy, S.O. 1986. Environment and cultivar effects on winter wheat response to ethephon plant growth regulator. *Agronomy Journal* **78**: 761-764.
435. Wong, S.C.; Cowan, I.R. and Farquhar, G.D. 1985. Leaf conductance in relation to rate of CO<sub>2</sub> assimilation. I. Influence of nitrogen nutrition,

phosphorus nutrition, photon flux density and ambient partial pressure of CO<sub>2</sub> during ontogeny. *Plant Physiology* **78**: 821-825.

436. Yang, D.Q.; Yang, J.X. and Hu, Y.W. 1994. Effects of S-3307 on some physiological characteristics of rape seedlings. *Plant Physiology Communication* **30**: 182-185.
437. Yang, S. and Hoffman, N.E. 1984. Ethylene biosynthesis and its regulation in higher plants. *Annual Review of Plant Physiology* **35**: 155-189.
438. Yang, S.F. 1987. Regulation of biosynthesis and action of ethylene. *Acta Horticulture* **201**: 53-59.
439. Zaman, A. and Choudhri, S.K. 1998. Water use efficiency and moisture extraction patterns of safflower in semi-arid region of Indian sub-tropics. *International Journal of Tropical Agriculture* **16**: 157-163.
440. Zerbe, R. and Wild, A. 1981. The effect of indole-3-acetic acid on photosynthetic apparatus of *Sinapis alba*. *Photosynthesis Research* **1**: 71-81.
441. Zhang, J. and Davies, W.J. 1990. Does ABA in the xylem control the rate of leaf growth in soil-dried maize and sunflower plants? *Journal of Experimental Botany* **41**: 1125-1132.
442. Zhou, W. and Ye, Q. 1996. Physiological and yield effects of uniconazole on winter rape (*Brassica napus* L.). *Journal of Plant Growth Regulation* **15**: 69-73.
443. Zhou, W.J. and Xi, H.F. 1993. Effects of mixtalol and paclobutrazol on photosynthesis and yield of rape (*Brassica napus*). *Journal of Plant Growth Regulation* **12**: 157-161.
444. Zobel, R.W. and Roberts, L.W. 1978. Effects of low concentration of ethylene on cell division and cytodifferentiation in lettuce pit explants. *Canadian Journal of Botany* **56**: 987-990.

# ***APPENDIX***



## PREPARATION OF REAGENTS

Reagents used in various determinations were prepared as follows:

### 1. Phosphate buffer (0.1M)

This was prepared by weighing 27.2g of  $\text{KH}_2\text{PO}_4$  and 45.6g of  $\text{K}_2\text{HPO}_4$  and dissolving each separately in 1 litre double distilled water. A 16ml of the  $\text{KH}_2\text{PO}_4$  solution and 84ml of the  $\text{K}_2\text{HPO}_4$  solution were then mixed and diluted to 200ml with double distilled water.

### 2. $\text{KNO}_3$ solution (0.2M)

This was made by weighing 20.2g of potassium nitrate in 1 litre of double distilled water.

### 3. Isopropanol solution (5%)

5ml of isopropanol was added to distilled water to make 100ml of solution.

### 4. Chloramphenical solution (0.5%)

0.5g of chloroamphenical was dissolved to get 100ml with double distilled water.

### 5. Sulphanilamide in 3NHCl (1%)

1g sulphanilamide was dissolved in 10ml 3NHCl.

### 6. N-1-Naphthyl-ethylene diamine dihydrochloride (NED-HCl) solution (0.2%)

0.2g of N-1-Naphthyl-ethylene diamine dihydrochloride was dissolved in double distilled water for the final volume of 100ml.

### 7. Sodium hydroxide (2.5N NaOH)

This was prepared by dissolving 10.0g of NaOH in double distilled water and the final volume was made 100ml.

**8. Nessler's reagent**

3.5g of potassium iodide was dissolved in 100ml of double distilled water to which 4 per cent mercuric chloride solution was added with stirring until a slight red precipitate remained. Thereafter, 120g of sodium hydroxide with 250ml of double distilled water were added. The volume was made to one litre. The mixture was decanted and kept in an amber-coloured bottle.

**9. Hydrochloride acid (0.5N)**

Hydrochloride acid (21.49ml) was mixed with 478.51ml of double distilled water to get 500ml.

**10. Iodine monochloride solution**

Iodine (13g) was dissolved in a mixture of 300ml of carbontetrachloride and 700ml of glacial acetic acid and the resulting solution was divided in solution A and B. To 20ml of solution A, 15ml of potassium iodide solution and 100ml of double distilled water was added and titrated against 0.1N sodium thiosulphate solution using starch solution as an indicator. Chlorine gas was passed through solution B until the amount of 0.1N sodium thiosulphate solution required for the titration was not more than double of that needed in solution A.

**11. Phenolphthalein solution**

Phenolphthalein (10g) was dissolved in 95 percent ethanol and the volume was made to 1 litre.

**12. Potassium hydroxide (0.1N KOH)**

5.6g of KOH was dissolved in 95 per cent ethanol and the volume was made to 1 litre.

**13. Potassium hydroxide solution (0.5N)**

Potassium hydroxide (28g) was dissolved in 95 per cent ethanol and the volume was made to 1 litre.

**14. Potassium iodide solution**

Potassium iodide (150g) was dissolved in double distilled water and the volume was made to 1 litre.

#### **15. Sodium thiosulphate solution (0.1N)**

Sodium thiosulphate (24.8g) was dissolved in double distilled water and the volume was made to 1 litre.

#### **16. Solvent mixture**

Ethanol 95 percent was mixed in diethyl ether in 1:1 ratio. This mixture of solvent was neutralized just before use of 0.1N KOH solution in the presence of phenolphthalein solution as an indicator.

#### **17. Starch solution**

Soluble starch (1g) was dissolved in 100ml of boiling double distilled water.

#### **18. Silver thiosulphate solution (1mM)**

540mg of  $\text{AgNO}_3$  was dissolved in 1 litre of double distilled water to get 2.0mM  $\text{AgNO}_3$  and 1264mg of  $\text{Na}_2\text{S}_2\text{O}_3$  was dissolved in 1 litre of double distilled water to get 8mM  $\text{Na}_2\text{S}_2\text{O}_3$ . The above two solutions were mixed together, resulted in 1mM of silver thiosulphate.

# ***TABLES***

Table 7 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant height (cm plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)															
80															
Cultivar	Ethrel concentration (µL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	90 17	94 43	103 80	101 37	98 03	97 56	94 70	100 63	109 27	106 30	104 20	103 02			
PBM16	86 30	90 57	99 00	95 70	93 67	93 05	91 37	95 90	105 77	102 20	99 33	98 91			
Mean	88 23	92 50	101 40	98 53	95 85		93 03	98 27	107 52	104 25	101 77				
120															
Alankar	101 43	106 70	114 90	112 50	110 43	109 19	80						100	120	
PBM16	97 30	100 77	111 60	109 50	106 40	105 11	Spray						1 65	1 28	1 28
Mean	99 37	103 73	113 25	111 00	108 42		Cultivar						1 05	0 81	0 81
							Interaction						NS	NS	NS

Table 8. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf area ( $\text{cm}^2 \text{ plant}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Cultivar	Sampling stage (days after sowing)												L.S.D. at 5%			
	80						100									
	Ethrel concentration (μL/L)															
	0	100	200	400	600	Mean	0	100	200	400	600	Mean				
Alankar	3077.33	3225.67	3907.00	3584.67	3458.67	3450.67	3295.33	3470.33	4242.00	3993.67	3866.67	3773.60				
PBM16	2900.00	3108.00	3684.67	3444.33	3346.00	3296.60	3171.67	3321.00	4096.67	3845.33	3731.00	3633.13				
Mean	2988.67	3166.83	3795.83	3514.50	3402.33		3233.50	3395.67	4169.33	3919.50	3798.83					
120																
Alankar	1880.67	1990.33	2546.00	2394.67	2303.67	2223.07								80	100	120
PBM16	1799.67	1884.00	2453.67	2292.00	2232.33	2132.33								Spray		
Mean	1840.17	1937.17	2499.83	2343.33	2268.00									Cultivar		
														Interaction		
														NS		



Table 10 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on specific leaf area ( $\text{cm}^2\text{g}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80							100					
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	288.00	291.67	303.67	299.67	295.00	295.60	399.33	406.00	422.33	415.33	410.00	410.60
PBM16	284.33	287.33	300.67	295.33	291.33	291.80	395.67	399.33	416.67	411.00	405.33	405.60
Mean	286.17	289.50	302.17	297.50	293.17		397.50	402.67	419.50	413.17	407.67	
L.S.D. at 5%												
120							80					
Alankar	428.67	433.00	447.33	441.33	436.00	437.27						
PBM16	425.00	428.33	441.67	436.33	432.00	432.67						
Mean	426.83	430.67	444.50	438.83	434.00							
							Spray		0.99	1.17	1.07	
							Cultivar		0.62	0.74	0.67	
							Interaction		NS	NS	NS	



Table 11. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on specific leaf weight ( $\text{mg cm}^{-2}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
Cultivar	80						100					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
	Alankar	3.47	3.42	3.29	3.33	3.38	3.38	2.50	2.46	2.37	2.41	2.44
PBM16	3.52	3.47	3.32	3.38	3.43	3.42	2.52	2.50	2.40	2.43	2.47	2.46
Mean	3.49	3.45	3.31	3.36	3.41		2.51	2.48	2.38	2.42	2.45	

L.S.D. at 5%															
Cultivar	120						80						100	120	
	Spray												0.012	0.010	0.010
	Cultivar												0.007	0.006	0.006
	Interaction												NS	NS	NS

Table 12. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
Cultivar	80						100					
	Ethrel concentration (µL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	32.51	33.48	37.78	35.35	34.78	34.78	34.78	35.78	40.01	38.51	37.91	37.40
PBM16	31.17	32.80	36.18	34.58	34.24	33.79	34.02	35.01	39.33	37.67	37.38	36.68
Mean	31.84	33.14	36.98	34.97	34.51		34.40	35.40	39.67	38.09	37.65	37.40
Cultivar	120						L.S.D. at 5%					
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	36.99	38.05	43.11	41.61	40.99	40.15						
PBM16	36.21	37.15	42.46	40.77	40.49	39.42						
Mean	36.60	37.60	42.79	41.19	40.74							
							Spray	0.31	0.32	0.32	0.32	0.32
							Cultivar	0.20	0.20	0.20	0.20	0.20
							Interaction	0.44	NS	NS	NS	NS

Table 13 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L ) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Cultivar	Sampling stage (days after sowing)														
	80						100								
	Ethrel concentration (µL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	10.68	11.05	12.87	11.95	11.71	11.65	8.25	8.54	10.04	9.61	9.43	9.17			
PBM16	10.19	10.78	12.25	11.65	11.48	11.27	8.01	8.31	9.83	9.35	9.20	8.94			
Mean	10.44	10.92	12.56	11.80	11.60		8.13	8.43	9.94	9.48	9.31				
L.S.D at 5%															
120															
Alankar	4.39	4.59	5.69	5.42	5.28	5.07	80						100	120	
PBM16	4.23	4.40	5.55	5.25	5.17	4.92	Spray						0.10	0.08	0.04
Mean	4.31	4.50	5.62	5.34	5.22		Cultivar						0.06	0.05	0.02
							Interaction						0.14	NS	NS

Table 14 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stem dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)															
Cultivar	80						100								
	Ethrel concentration (µL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	19 10	19 48	21 07	19 87	19 63	19 83	20 78	21 20	22 74	22 08	21 80	21 72			
PBM16	18 39	19 18	20 27	19 54	19 44	19 36	20 46	20 88	22 42	21 72	22 05	21 51			
Mean	18 75	19 33	20 67	19 70	19 54		20 62	21 04	22 58	21 90	21 93				
L S D at 5%															
Cultivar	120						80						100	120	
	Spray												0 20	0 19	0 23
	Cultivar												0 12	0 12	0 14
Interaction												0 28	0 27	NS	

Table 15. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on pod dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80												
Cultivar	Ethrel concentration (µL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
	Alankar	2.73	2.94	3.85	3.53	3.43	3.30	5.75	6.04	7.23	6.83	6.68
PBM16	2.58	2.84	3.66	3.40	3.32	3.16	5.55	5.82	7.08	6.60	6.14	6.24
Mean	2.66	2.89	3.75	3.47	3.38		5.65	5.93	7.16	6.71	6.41	
L.S.D. at 5%												
120												
Alankar	8.81	9.12	10.79	10.29	10.06	9.81						
PBM16	8.59	8.85	10.58	10.02	9.89	9.59						
Mean	8.70	8.98	10.69	10.16	9.98							
							Spray					
							Cultivar					
							Interaction					
							NS					
							0.09					
							NS					

Table 16. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80							100					
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	32.85	33.02	34.05	33.81	33.68	33.48	23.71	23.88	25.10	24.95	24.86	24.50
PBM16	32.70	32.87	33.85	33.68	33.52	33.32	23.54	23.73	24.99	24.82	24.68	24.35
Mean	32.77	32.94	33.95	33.74	33.60		23.63	23.81	25.05	24.89	24.77	
L.S.D. at 5%												
120												
Alankar	11.86	12.07	13.20	13.03	12.90	12.61						
PBM16	11.68	11.84	13.08	12.87	12.76	12.44						
Mean	11.77	11.96	13.14	12.95	12.83							
							Spray			0.08	0.07	0.05
							Cultivar			0.05	0.04	0.03
							Interaction			NS	NS	NS

Table 17 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stem dry weight (%) of mustard (*Brassica juncea* L ) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Cultivar	Sampling stage (days after sowing)														
	80						100								
	Ethrel concentration (μL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	58.75	58.19	55.76	56.19	56.45	57.07	59.75	59.25	56.83	57.32	57.51	58.13			
PBM16	59.01	58.47	56.04	56.50	56.77	57.36	60.14	59.64	57.00	57.66	58.97	58.68			
Mean	58.88	58.33	55.90	56.35	56.61		59.95	59.45	56.91	57.49	58.24				
L S D at 5%															
Alankar	64.33	63.96	61.76	62.24	62.57	62.97	80						100	120	
PBM16	64.59	64.35	62.00	62.54	62.80	63.26	Spray						0.12	0.10	0.09
Mean	64.46	64.15	61.88	62.39	62.69		Cultivar						0.08	0.07	0.06
							Interaction						NS	0.15	NS

Table 18. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on pod dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)																		
80																		
Cultivar	Ethrel concentration (μL/L)																	
	0	100	200	400	600	Mean	0	100	200	400	600	Mean						
	Alankar	8.40	8.79	10.19	9.99	9.87	9.45	16.53	16.87	18.07	17.72	17.62	17.36					
PBM16	8.29	8.66	10.11	9.82	9.70	9.32	16.32	16.62	18.01	17.52	16.41	16.98						
Mean	8.34	8.72	10.15	9.91	9.79		16.43	16.75	18.04	17.62	17.02							
L.S.D. at 5%																		
120																		
Alankar	23.81	23.97	25.04	24.73	24.54	24.42							80	100	120			
PBM16	23.73	23.82	24.92	24.58	24.43	24.29							Spray			0.07	0.06	0.07
Mean	23.77	23.89	24.98	24.66	24.49								Cultivar			0.04	0.04	0.04
													Interaction			NS	0.09	NS



Table 19. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf fresh weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	44.86	47.00	55.71	51.32	49.70	49.72	32.22	33.68	40.51	38.54	37.34	36.46
PBM16	42.40	45.06	52.52	49.59	48.62	47.64	30.87	32.33	39.14	37.03	35.98	35.07
Mean	43.63	46.03	54.12	50.45	49.16		31.55	33.00	39.82	37.79	36.66	
L.S.D. at 5%												
120												
Alankar	16.26	17.17	21.84	20.71	19.87	19.17						
PBM16	15.49	16.23	20.95	19.74	19.17	18.31						
Mean	15.88	16.70	21.40	20.22	19.52							
							Spray		0.40		0.32	
							Cultivar		0.25		0.20	
							Interaction		0.57		NS	
											NS	

Table 20. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf turgid weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)																		
80																		
Cultivar	Ethrel concentration (µL/L)																	
	100																	
	0	100	200	400	600	Mean	0	100	200	400	600	Mean						
Alankar	66.22	68.02	76.55	72.32	70.87	70.79	54.09	55.96	63.26	61.51	60.33	59.03						
PBM16	63.70	66.84	73.48	71.05	70.01	69.01	53.26	54.85	62.40	60.30	59.34	58.03						
Mean	64.96	67.43	75.02	71.68	70.44		53.68	55.40	62.83	60.91	59.83							
L.S.D. at 5%																		
120																		
Alankar	30.27	31.35	37.55	36.33	35.37	34.17							80	100	120			
PBM16	29.40	30.34	36.93	35.44	34.88	33.40							Spray			0.60	0.65	0.25
Mean	29.83	30.84	37.24	35.89	35.13								Cultivar			0.38	0.41	0.16
												Interaction			0.84	NS	NS	

Table 21. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf relative water content (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)															
80															
Cultivar	Ethrel concentration (µL/L)														
	100														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	61.55	63.11	67.27	65.21	64.18	64.26	51.91	53.01	57.23	55.75	54.83	54.54			
PBM16	60.19	61.16	65.71	63.88	63.45	62.88	50.51	51.61	55.76	54.32	53.41	53.12			
Mean	60.87	62.13	66.49	64.54	63.82		51.21	52.31	56.50	55.03	54.12				
L.S.D. at 5%															
120															
Alankar	45.89	47.00	50.68	49.45	48.47	48.30	80						100	120	
PBM16	44.73	45.60	49.08	47.99	47.12	46.91	Spray						0.25	0.23	0.30
Mean	45.31	46.30	49.88	48.72	47.80		Cultivar						0.16	0.15	0.19
							Interaction						0.35	NS	NS

Table 22. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on rate of photosynthesis ( $\mu \text{ mol m}^{-2} \text{ s}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
100												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	17.15	18.33	21.16	20.12	19.24	19.20	15.08	16.30	19.36	18.12	17.24	17.22
PBM16	16.22	17.28	20.14	19.14	18.25	18.21	14.35	15.27	18.50	17.30	16.36	16.36
Mean	16.69	17.81	20.65	19.63	18.75		14.72	15.78	18.93	17.71	16.80	
L.S.D. at 5%												
80												
100												
Spray												
0.19												
0.23												
Cultivar												
0.12												
0.14												
Interaction												
NS												
NS												

Table 23. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stomatal conductance ( $\text{mol m}^{-2}\text{s}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	0.44	0.45	0.48	0.46	0.45	0.45	0.41	0.42	0.45	0.44	0.43	0.43
PBM16	0.42	0.44	0.46	0.45	0.44	0.44	0.41	0.42	0.44	0.43	0.42	0.42
Mean	0.43	0.44	0.47	0.46	0.45		0.41	0.42	0.44	0.43	0.42	

L.S.D. at 5%			
		80	100
Spray		0.006	0.006
Cultivar		0.004	0.004
Interaction		NS	NS

Table 24. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on internal CO<sub>2</sub> concentration ( $\mu$  mol mol<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
100												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	273.03	283.53	312.13	301.50	292.67	292.57	263.43	272.43	302.20	292.20	281.40	282.33
PBM16	267.03	275.60	301.43	292.77	283.83	284.13	258.07	267.33	294.43	284.80	275.30	275.99
Mean	270.03	279.57	306.78	297.13	288.25		260.75	269.88	298.32	288.50	278.35	

L.S.D. at 5%			
		80	100
Spray		2.96	3.35
Cultivar		1.87	2.12
Interaction		NS	NS

Table 25. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on transpiration rate ( $\text{kg m}^{-2} \text{ day}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)													
80													
100													
Cultivar	Ethrel concentration (µL/L)												
	0	100	200	400	600	Mean	0	100	200	400	600	Mean	
	Alankar	12.79	12.87	13.28	13.20	13.16	13.06	10.44	10.59	11.02	10.75	10.70	10.70
	PBM16	12.64	12.85	13.20	12.98	12.97	12.93	10.42	10.49	10.92	10.69	10.66	10.64
	Mean	12.71	12.86	13.24	13.09	13.07		10.43	10.54	10.97	10.72	10.68	
L.S.D. at 5%													
80													
100													
Spray													
0.04													
Cultivar													
0.03													
Interaction													
0.06													
NS													

Table 26 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on carboxylation efficiency (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80							100					
Cultivar	Ethiel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	6.28	6.44	6.77	6.67	6.57	6.55	5.72	5.98	6.40	6.20	6.12	6.08
PBM16	6.40	6.26	6.68	6.53	6.43	6.46	5.56	5.71	6.28	6.07	5.61	5.84
Mean	6.34	6.35	6.73	6.60	6.50		5.64	5.84	6.34	6.14	5.86	

L.S D. at 5%		
	80	100
Spray	0.19	0.25
Cultivar	NS	0.16
Interaction	NS	NS



Table 27. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on photosynthetic water use efficiency ( $\mu\text{molmol}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
100												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	39.28	41.05	44.38	43.74	42.44	42.18	36.48	38.49	43.02	41.51	40.09	39.92
PBM16	38.32	39.58	44.10	42.54	41.80	41.26	35.29	36.65	42.37	40.54	39.26	38.82
Mean	38.80	40.31	44.24	43.14	42.12		35.89	37.57	42.70	41.03	39.67	
L.S.D. at 5%												
80												
100												
Spray												
0.32												
0.48												
Cultivar												
0.20												
0.30												
Interaction												
0.45												
NS												

Table 28. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant water use efficiency ( $\text{mg g}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
100												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	2.54	2.60	2.84	2.68	2.64	2.66	3.30	3.38	3.63	3.58	3.54	3.49
PBM16	2.46	2.55	2.74	2.66	2.64	2.61	3.27	3.34	3.60	3.52	3.50	3.44
Mean	2.50	2.58	2.79	2.67	2.64		3.28	3.36	3.62	3.55	3.52	
L.S.D. at 5%												
80												
100												
Spray												
0.02												
Cultivar												
0.01												
Interaction												
0.03												
NS												

Table 29 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrogen content (%) of mustard (*Brassica juncea* L ) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)															
80							100								
Cultivar	Ethrel concentration (μL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	2.22	2.32	2.41	2.38	2.37	2.34	1.99	2.16	2.32	2.27	2.24	2.20			
PBM16	2.13	2.22	2.37	2.33	2.31	2.27	1.85	2.00	2.23	2.13	2.12	2.06			
Mean	2.18	2.27	2.39	2.35	2.34		1.92	2.08	2.28	2.20	2.18				
L S D at 5%															
120															
Alankar	1.74	1.85	2.06	1.95	1.90	1.90	80						100	120	
PBM16	1.64	1.74	1.95	1.82	1.80	1.79	Spray						0.02	0.04	0.03
Mean	1.69	1.79	2.01	1.88	1.85		Cultivar						0.01	0.02	0.02
							Interaction						0.02	NS	NS

Table 30. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrogen accumulation (mg plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
Cultivar	80						100					
	Ethrel concentration (µL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	722.67	775.33	901.33	841.33	824.33	813.00	691.00	771.33	929.67	874.33	849.00	823.07
PBM16	664.00	727.00	856.33	804.33	791.00	768.53	629.67	700.33	875.33	801.33	791.00	759.53
Mean	693.33	751.17	878.83	822.83	807.67		660.33	735.83	902.50	837.83	820.00	
L.S.D at 5%												
120												
Alankar	642.33	702.67	889.67	810.00	777.33	764.40						
PBM16	593.67	645.00	826.00	741.67	728.67	707.00						
Mean	618.00	673.83	857.83	775.83	753.00							
Spray												
Cultivar												
Interaction												
NS												

Table 31. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on number of pods plant<sup>-1</sup>, number of seeds pod<sup>-1</sup> and 1000 seed weight (g) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16

Cultivar	Pods plant <sup>-1</sup>						Seeds pod <sup>-1</sup>					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	144.33	148.67	160.33	156.33	153.67	152.67	12.90	13.27	14.77	14.33	14.32	13.92
PBM16	140.33	144.67	158.33	150.67	150.00	148.80	12.75	13.03	14.62	14.20	14.17	13.75
Mean	142.33	146.67	159.33	153.50	151.83		12.83	13.15	14.69	14.27	14.24	
L.S.D. at 5%												
1000 seed weight												
Alankar	4.44	4.45	4.58	4.54	4.51	4.51						
PBM16	4.41	4.44	4.56	4.51	4.51	4.48						
Mean	4.43	4.45	4.57	4.53	4.51							
							Spray	1.67	0.08	0.01		
							Cultivar	1.05	0.05	0.01		
							Interaction	NS	NS	NS	NS	NS

Table 32. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on seed yield ( $\text{q ha}^{-1}$ ), biological yield ( $\text{q ha}^{-1}$ ) and harvest index (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16

Sampling stage (days after sowing)													
Cultivar	Seed yield						Biological yield						
	Ethrel concentration (μL/L)												
	0	100	200	400	600	Mean	0	100	200	400	600	Mean	
Alankar	10.33	10.98	13.55	12.73	12.41	12.00	46.23	47.57	53.89	52.02	51.23	50.19	
PBM16	9.87	10.46	13.19	12.05	11.97	11.51	45.27	46.43	53.08	50.97	50.62	49.27	
Mean	10.10	10.72	13.37	12.39	12.19		45.75	47.00	53.49	51.49	50.93		
L.S.D. at 5%													
Alankar	Harvest index						Seed yield						Har-vest index
	22.35	23.07	25.15	24.46	24.21	23.85	Spray						0.20
	21.80	22.52	24.85	23.64	23.65	23.29	Cultivar						0.13
	22.07	22.80	25.00	24.05	23.93		Interaction						NS
PBM16													NS
Mean													NS

Table 33. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on oil yield (q ha<sup>-1</sup>) and oil content (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16

Sampling stage (days after sowing)												
Cultivar	Oil yield						Oil content					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
	Alankar	3.63	3.90	5.02	4.65	4.45	4.33	35.19	35.52	37.07	36.55	35.91
PBM16	3.43	3.66	4.77	4.32	4.22	4.08	34.72	35.00	36.15	35.85	35.28	35.40
Mean	3.53	3.78	4.90	4.48	4.34		34.96	35.26	36.61	36.20	35.60	

L.S.D. at 5%		
Oil yield    Oil content		
Spray	0.07	0.09
Cultivar	0.04	0.06
Interaction	NS	0.13

Table 34. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on acid value, iodine value and Saponification value of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16

Sampling stage (days after sowing)												
Cultivar	Acid value						Iodine value					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	3.26	3.56	3.97	3.84	3.68	3.66	96.40	97.51	99.28	98.62	97.72	97.91
PBM16	3.04	3.21	3.53	3.40	3.29	3.30	95.84	96.33	97.53	97.14	96.79	96.73
Mean	3.15	3.39	3.75	3.62	3.49		96.12	96.92	98.41	97.88	97.25	
L.S.D. at 5%												
Saponification value												
Alankar	142.80	145.22	147.46	147.02	146.71	145.84						
PBM16	140.33	143.59	146.30	145.30	144.37	143.98						
Mean	141.57	144.40	146.88	146.16	145.54							
							Spray	0.06	NS	0.53		
							Cultivar	0.04	NS	0.33		
							Interaction	0.09	NS	NS		



Table 35. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant height (cm plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)															
Cultivar	80						100								
	Ethrel concentration (μL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	64.37	67.27	74.93	71.87	69.60	69.61	69.17	71.97	79.17	76.87	75.13	74.46			
PBM16	60.43	62.97	70.13	67.30	64.33	65.03	64.47	67.83	74.40	72.43	69.63	69.75			
Mean	62.40	65.12	72.53	69.58	66.97		66.82	69.90	76.78	74.65	72.38				
L.S.D. at 5%															
120															
Alankar	74.07	77.27	84.27	83.00	80.20	79.76	80						100	120	
PBM16	67.97	70.23	79.80	76.33	73.23	73.51	Spray						1.38	1.45	1.20
Mean	71.02	73.75	82.03	79.67	76.72		Cultivar						0.87	0.92	0.76
							Interaction						NS	NS	NS

Table 36. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf area (cm<sup>2</sup> plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Cultivar	Sampling stage (days after sowing)																	
	80						100											
	Ethrel concentration (µL/L)																	
	0	100	200	400	600	Mean	0	100	200	400	600	Mean						
Alankar	2160.33	2356.67	2950.00	2760.67	2634.33	2572.40	2317.67	2531.33	3244.33	3046.00	2929.00	2813.67						
PBM16	2042.33	2234.67	2755.33	2610.00	2512.67	2431.00	2222.33	2414.67	3060.33	2908.33	2816.33	2684.40						
Mean	2101.33	2295.67	2852.67	2685.33	2573.50		2270.00	2473.00	3152.33	2977.17	2872.67							
L.S.D. at 5%																		
120																		
Alankar	1355.33	1456.33	2001.33	1869.00	1789.33	1694.27							80	100	120			
PBM16	1251.00	1394.00	1886.67	1766.00	1701.00	1599.73							Spray			26.46	26.41	24.35
Mean	1303.17	1425.17	1944.00	1817.50	1745.17								Cultivar			16.73	16.70	15.40
												Interaction			37.41	37.35	NS	

Table 37. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf area index of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80												
100												
Cultivar	Ethrel concentration (µL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	2.70	2.95	3.69	3.45	3.29	3.22	2.90	3.16	4.06	3.81	3.41	3.47
PBM16	2.55	2.79	3.44	3.26	3.14	3.04	2.78	3.02	3.83	3.64	3.52	3.36
Mean	2.63	2.87	3.57	3.36	3.22		2.84	3.09	3.94	3.72	3.47	
L.S.D. at 5%												
120												
Alankar	1.69	1.82	2.50	2.34	2.24	2.12						
PBM16	1.56	1.74	2.36	2.21	2.13	2.00						
Mean	1.63	1.78	2.43	2.27	2.18							
						Spray		0.03	0.17	0.03		
						Cultivar		0.02	0.11	0.02		
						Interaction		0.05	NS	NS		

Table 38. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on specific leaf area ( $\text{cm}^2 \text{g}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80							100					
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
	Alankar	257.33	261.33	273.00	268.67	263.67	264.80	361.00	365.00	376.67	372.00	368.00
PBM16	252.67	257.00	267.67	263.33	259.67	260.07	357.00	361.33	371.67	366.00	363.00	363.80
Mean	255.00	259.17	270.33	266.00	261.67		359.00	363.17	374.17	369.00	365.50	
120							L.S.D. at 5%					
Alankar	399.67	406.00	423.67	417.00	412.00	411.67						
PBM16	392.00	399.67	416.33	409.33	404.00	404.27						
Mean	395.83	402.83	420.00	413.17	408.00							
							Spray			0.73	1.07	1.51
							Cultivar			0.46	0.67	0.96
							Interaction			NS	NS	NS

Table 39. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on specific leaf weight ( $\text{mg cm}^{-2}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
	Alankar	3.88	3.82	3.66	3.72	3.79	3.77	2.77	2.74	2.65	2.69	2.71
PBM16	3.95	3.88	3.73	3.79	3.85	3.84	2.80	2.77	2.69	2.73	2.75	2.74
Mean	3.92	3.85	3.70	3.76	3.82		2.79	2.75	2.67	2.71	2.73	
L.S.D. at 5%												
120												
Alankar	2.50	2.46	2.36	2.40	2.42	2.43						
PBM16	2.55	2.50	2.40	2.44	2.48	2.47						
Mean	2.53	2.48	2.38	2.42	2.45							
						Spray		0.011	0.010	0.010	0.010	
						Cultivar		0.007	0.006	0.006	0.007	
						Interaction		NS	NS	NS	NS	

Table 40. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)																		
80																		
Cultivar	Ethrel concentration (µL/L)																	
	100																	
	0	100	200	400	600	Mean	0	100	200	400	600	Mean						
Alankar	26.37	28.15	32.46	31.04	30.32	29.67	28.33	30.19	35.63	34.07	33.43	32.31						
PBM16	25.46	27.26	31.06	30.04	29.55	28.68	27.53	29.30	34.17	33.16	32.74	31.38						
Mean	25.92	27.71	31.76	30.54	29.94		27.88	29.74	34.90	33.62	33.09							
L.S.D. at 5%																		
120																		
Alankar	31.09	32.24	38.81	37.20	36.35	35.14							80	100	120			
PBM16	29.62	31.77	37.63	36.27	35.83	34.22							Spray			0.29	0.29	0.45
Mean	30.36	32.00	38.22	36.74	36.09								Cultivar			0.19	0.18	0.29
												Interaction			NS	NS	NS	

Table 41 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Cultivar	Sampling stage (days after sowing)											
	80						100					
	Ethrel concentration (µL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	8.40	9.01	10.80	10.27	9.99	9.69	6.42	6.93	8.61	8.18	7.96	7.62
PBM16	8.08	8.69	10.28	9.90	9.67	9.32	6.22	6.68	8.23	7.94	7.76	7.37
Mean	8.24	8.85	10.54	10.08	9.83		6.32	6.81	8.42	8.06	7.86	
L.S.D at 5%												
	120											
Alankar	3.39	3.59	4.72	4.48	4.34	4.10						
PBM16	3.19	3.49	4.53	4.31	4.21	3.95						
Mean	3.29	3.54	4.63	4.40	4.28							
							Spray			0.09	0.06	0.06
							Cultivar			0.05	0.04	0.04
							Interaction			NS	0.09	NS

Table 42. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stem dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80							100					
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	15.95	16.89	18.47	17.98	17.65	17.39	17.36	18.43	20.80	20.14	19.86	19.32
PBM16	15.47	16.44	17.76	17.48	17.33	16.90	17.01	17.97	20.01	19.65	19.56	18.84
Mean	15.71	16.67	18.12	17.73	17.49		17.19	18.20	20.41	19.90	19.71	
L.S.D. at 5%												
120												
Alankar	20.54	21.12	25.01	23.76	23.30	22.74						
PBM16	19.65	20.90	24.00	23.28	23.08	22.18						
Mean	20.10	21.01	24.50	23.52	23.19							
							Spray			0.19	0.19	0.35
							Cultivar			0.12	0.12	0.22
							Interaction			NS	NS	NS



Table 43 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on pod dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)															
80															
Cultivar	Ethrel concentration (μL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	2.02	2.25	3.19	2.79	2.68	2.59	4.45	4.83	6.21	5.75	5.60	5.37			
PBM16	1.91	2.14	3.02	2.66	2.55	2.45	4.30	4.64	5.93	5.56	5.43	5.17			
Mean	1.97	2.20	3.11	2.72	2.62		4.37	4.74	6.07	5.66	5.52				
L.S.D. at 5%															
120															
Alankar	7.16	7.53	9.42	8.96	8.71	8.36	80						100	120	
PBM16	6.78	7.38	9.10	8.68	8.54	8.10	Spray						0.03	0.04	0.10
Mean	6.97	7.46	9.26	8.82	8.63		Cultivar						0.02	0.03	0.07
											Interaction		NS	0.06	NS

Table 44. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)													
80													
Cultivar	Ethrel concentration (μL/L)												
	100												
	0	100	200	400	600	Mean	0	100	200	400	600	Mean	
Alankar	31.84	32.00	33.28	33.07	32.94	32.63	22.73	22.97	24.17	24.02	23.81	23.54	
PBM16	31.72	31.86	33.10	32.95	32.73	32.47	22.60	22.80	24.10	23.95	23.69	23.43	
Mean	31.78	31.93	33.19	33.01	32.84		22.67	22.89	24.14	23.99	23.75		
120													
L.S.D. at 5%													
Alankar	10.90	11.12	12.17	12.04	11.94	11.63	80 100 120						
PBM16	10.77	10.98	12.04	11.89	11.75	11.48	Spray 0.23 0.08 0.07						
Mean	10.84	11.05	12.10	11.97	11.85		Cultivar NS 0.05 0.05						
							Interaction NS NS NS						

Table 45. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stem dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80							100					
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	60.50	60.00	56.88	57.93	58.27	58.72	61.51	61.05	58.38	59.10	59.42	59.89
PBM16	60.76	60.29	57.18	58.20	58.65	59.02	61.79	61.35	58.57	59.27	59.72	60.14
Mean	60.63	60.15	57.03	58.07	58.46		61.65	61.20	58.48	59.18	59.57	
L.S.D. at 5%												
120												
Alankar	66.06	65.51	63.88	63.87	64.09	64.68						
PBM16	66.35	65.80	63.77	64.18	64.41	64.90						
Mean	66.21	65.66	63.83	64.02	64.25							
							Spray		0.13	0.11	0.25	
							Cultivar		0.08	0.07	0.16	
							Interaction		NS	NS	NS	

Table 46. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on pod dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
80							100					
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	7.66	7.99	9.84	8.99	8.85	8.66	15.75	15.99	17.44	16.88	16.77	16.57
PBM16	7.51	7.85	9.71	8.84	8.62	8.51	15.60	15.85	17.35	16.78	16.58	16.43
Mean	7.59	7.92	9.77	8.92	8.73		16.68	15.92	17.39	16.83	16.68	
L.S.D. at 5%												
120												
Alankar	23.03	23.37	24.26	24.08	23.96	23.74						
PBM16	22.83	23.22	24.19	23.93	23.83	23.61						
Mean	22.96	23.29	24.22	24.01	23.90							
							Spray		0.07	0.05	0.06	
							Cultivar		0.04	0.03	0.04	
							Interaction		NS	NS	NS	

Table 47 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf fresh weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L ) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)															
80															
Cultivar	Ethrel concentration (μL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	29.74	32.36	38.68	36.93	35.65	34.67	20.70	22.67	29.05	27.43	26.27	25.22			
PBM16	28.27	30.80	36.69	35.36	33.79	32.98	19.84	21.66	27.44	26.24	25.18	24.07			
Mean	29.01	31.58	37.69	36.15	34.72		20.27	22.16	28.24	26.84	25.72				
120															
Alankar	9.65	10.26	13.70	13.02	12.53	11.83	L S.D at 5%						80	100	120
PBM16	9.02	10.41	13.06	12.39	11.97	11.37	Spray						0.35	0.26	0.35
Mean	9.34	10.34	13.38	12.70	12.25		Cultivar						0.22	0.16	0.22
						Interaction						NS	NS	NS	

Table 48. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf turgid weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)													
Cultivar	80						100						
	Ethrel concentration (μL/L)												
	0	100	200	400	600	Mean	0	100	200	400	600	Mean	
Alankar	54.40	58.11	67.52	65.19	63.42	61.73	43.95	47.25	56.85	54.83	53.33	51.24	
PBM16	52.90	56.46	64.78	63.34	61.91	59.88	42.94	45.76	54.73	53.62	52.36	49.88	
Mean	53.65	57.29	66.15	64.27	62.66		43.45	46.50	55.79	54.22	52.85		
L.S.D. at 5%													
	120												
Alankar	24.24	25.46	32.57	31.36	30.51	28.83							120
PBM16	22.92	24.93	31.48	30.41	29.68	27.88							
Mean	23.58	25.20	32.03	30.88	30.10								
							Spray			0.57	0.43	0.43	
							Cultivar			0.36	0.27	0.27	
							Interaction			NS	NS	NS	

Table 49. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf relative water content (%) leaf area index of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
Cultivar	80						100					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	46.43	47.55	49.15	48.58	48.04	47.95	37.79	39.12	42.36	41.26	40.34	40.18
PBM16	45.05	46.29	48.22	47.65	46.17	46.68	37.07	38.32	41.31	40.06	39.06	39.17
Mean	45.74	46.92	48.69	48.11	47.11		37.43	38.72	41.84	40.66	39.70	

L.S.D. at 5%															
Cultivar	120						80						100	120	
	Spray														
	Cultivar														
Alankar	30.02	30.49	32.20	31.76	31.25	31.15									
PBM16	29.57	30.02	31.65	30.95	30.48	30.53									
Mean	29.80	30.25	31.92	31.36	30.88										
							Interaction						0.29	0.26	NS

Table 50. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on rate of photosynthesis ( $\mu \text{ mol m}^{-2} \text{ s}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
	Alankar	15.49	16.42	19.20	18.40	17.27	17.36	14.42	15.32	18.33	17.13	16.31
PBM16	14.23	15.37	18.61	17.42	16.34	16.39	13.36	14.44	17.27	16.45	15.21	15.35
Mean	14.86	15.90	18.90	17.91	16.80		13.89	14.88	17.80	16.79	15.76	

L.S.D. at 5%		
	80	100
Spray	0.36	0.27
Cultivar	0.23	0.17
Interaction	NS	NS



Table 51 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stomatal conductance ( $\text{mol m}^{-2} \text{s}^{-1}$ ) of mustard (*Brassica juncea* L ) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
100												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	0.41	0.42	0.45	0.44	0.43	0.43	0.40	0.41	0.44	0.43	0.42	0.42
PBM16	0.40	0.41	0.44	0.43	0.42	0.42	0.39	0.40	0.42	0.42	0.40	0.41
Mean	0.41	0.42	0.44	0.43	0.42		0.40	0.41	0.43	0.42	0.41	
L S D at 5%												
80												
100												
Spray												
0.007												
0.006												
Cultivar												
0.005												
0.004												
Interaction												
NS												
NS												

Table 52 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on intake carbon dioxide of mustard (*Brassica juncea* L ) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80							100					
Cultivar	Ethiel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	262.87	272.83	301.53	291.60	283.17	282.40	253.33	262.53	290.97	280.50	271.90	271.85
PBM16	254.00	263.03	294.20	283.17	272.90	273.46	245.13	253.17	283.87	273.93	263.27	263.87
Mean	258.43	267.93	297.87	287.38	278.03		249.23	257.85	287.42	277.22	267.58	

L S D at 5%		
	80	100
Spray	3.73	3.41
Cultivar	2.36	2.16
Interaction	NS	NS

Table 53. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on transpiration rate ( $\text{kg m}^{-2} \text{ day}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
100												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	11.40	11.77	12.23	12.13	12.05	11.91	9.28	9.62	10.20	10.09	10.01	9.84
PBM16	11.32	11.68	12.01	11.94	11.90	11.77	9.19	9.46	10.07	10.00	9.95	9.74
Mean	11.36	11.73	12.12	12.03	11.98		9.24	9.54	10.14	10.05	9.98	
L.S.D at 5%												
80												
100												
Spray												
0.04												
0.05												
Cultivar												
0.03												
0.03												
Interaction												
0.06												
NS												

Table 54. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on carboxylation efficiency (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	5.89	6.02	6.36	6.28	6.09	6.13	5.69	5.83	6.30	6.10	5.99	5.98
PBM16	5.60	5.84	6.32	6.15	5.98	5.98	5.45	5.70	6.08	6.00	5.77	5.80
Mean	5.75	5.93	6.34	6.21	6.04		5.57	5.76	6.19	6.05	5.88	

L.S.D. at 5%		
	80	100
Spray	0.16	0.12
Cultivar	0.10	0.07
Interaction	NS	NS

Table 55. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on photosynthetic water use efficiency ( $\mu\text{molmol}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)													
Cultivar	80						100						
	Ethrel concentration (μL/L)												
	0	100	200	400	600	Mean	0	100	200	400	600	Mean	
	Alankar	37.48	38.78	42.99	42.13	39.85	40.24	35.75	37.06	41.98	40.49	38.82	38.82
PBM16	35.27	37.18	42.61	40.51	39.21	38.96	33.96	35.80	40.79	39.48	38.02	37.61	
Mean	36.38	37.98	42.80	41.32	39.53		34.86	36.43	41.38	39.99	38.42		
L.S.D. at 5%													
							80						100
							Spray		0.53		0.51		
							Cultivar		0.33		0.32		
							Interaction		0.75		NS		

Table 56. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant water use efficiency ( $\text{mg g}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80 and 100 DAS

Sampling stage (days after sowing)												
80												
Cultivar	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	2.31	2.39	2.65	2.56	2.51	2.49	3.04	3.14	3.49	3.38	3.34	3.28
PBM16	2.25	2.33	2.59	2.52	2.48	2.43	2.97	3.10	3.39	3.31	3.29	3.21
Mean	2.28	2.36	2.62	2.54	2.50		3.01	3.12	3.44	3.35	3.32	

L.S.D. at 5%		
	80	100
Spray	0.02	0.02
Cultivar	0.01	0.01
Interaction	NS	NS

Table 57 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrogen content (%) of mustard (*Brassica juncea* L ) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Sampling stage (days after sowing)												
Cultivar	80						100					
	Ethiel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	2 10	2 18	2 29	2 25	2 24	2 21	1 75	1 94	2 12	2 05	2 03	1 98
PBM16	1 82	1 91	2 20	2 08	2 06	2 01	1 64	1 72	2 10	1 89	1 85	1 84
Mean	1 96	2 05	2 24	2 17	2 15		1 69	1 83	2 11	1 97	1 94	
L S D at 5%												
Cultivar	120											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
	Alankar	1 52	1 63	1 79	1 73	1 70	1 67					
PBM16	1.44	1 51	1 74	1.69	1 67	1 61						
Mean	1 48	1 57	1 77	1.71	1 68							
L S D at 5%												
Spray							0.02					
Cultivar							0.01					
Interaction							0.03					

Table 58. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrogen accumulation (mg plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16 at 80, 100 and 120 DAS

Cultivar	Sampling stage (days after sowing)														
	80						100								
	Ethrel concentration (µL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
Alankar	553.67	614.67	742.00	698.00	680.00	657.67	493.00	585.33	755.33	697.00	678.33	641.80			
PBM16	463.33	519.33	682.33	624.33	609.00	579.67	450.33	502.67	717.33	626.67	605.67	580.53			
Mean	508.50	567.00	712.17	661.17	644.50		471.67	544.00	736.33	661.83	642.00				
L.S.D. at 5%															
	120						80						100	120	
Alankar	471.67	524.33	696.00	644.67	616.67	590.67	Spray						9.18	14.00	10.75
PBM16	425.33	478.67	654.67	612.67	599.67	554.20	Cultivar						5.81	8.85	6.80
Mean	448.50	501.50	675.33	628.67	608.17		Interaction						12.99	19.80	NS



Table 59. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on number of pods plant<sup>-1</sup>, number of seeds pod<sup>-1</sup> and 1000 seed weight (g) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16

Cultivar	Pods plant <sup>-1</sup>						Seeds pod <sup>-1</sup>					
	Ethrel concentration (µL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	125.00	129.00	144.00	139.00	136.67	134.73	11.87	11.91	13.28	12.99	12.80	12.57
PBM16	122.00	125.67	139.00	136.00	133.67	131.27	11.80	11.85	13.20	12.83	12.71	12.48
Mean	123.50	127.33	141.50	137.50	135.17		11.84	11.88	13.24	12.91	12.75	
L.S.D. at 5%												
1000 seed weight												
Alankar	4.11	4.21	4.46	4.37	4.36	4.30						
PBM16	3.94	4.14	4.36	4.29	4.28	4.20						
Mean	4.03	4.18	4.41	4.33	4.32							
							Spray		1.36	0.05	0.05	
							Cultivar		0.86	0.03	0.03	
							Interaction		NS	NS	NS	

Table 60. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on seed yield ( $\text{q ha}^{-1}$ ), biological yield ( $\text{q ha}^{-1}$ ) and harvest index (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16

Sampling stage (days after sowing)													
Cultivar	Seed yield					Biological yield							
	Ethrel concentration (µL/L)												
	0	100	200	400	600	Mean	0	100	200	400	600	Mean	
Alankar	7.62	8.08	10.67	9.86	9.53	9.15	38.87	40.30	48.52	46.50	45.43	43.92	
PBM16	7.10	7.71	10.00	9.36	9.09	8.65	37.03	39.71	47.04	45.34	44.78	42.78	
Mean	7.36	7.90	10.34	9.61	9.31		37.95	40.01	47.78	45.92	45.11		
L.S.D. at 5%													
Alankar	Harvest index												
	19.60	20.05	21.99	21.21	20.98	20.77	Seed yield						Har-vest index
	19.16	19.42	21.26	20.61	20.30	20.15	0.13						0.57
	19.38	19.38	21.63	20.91	20.64		0.08						0.36
PBM16	Spray												0.17
Mean	Cultivar												0.11
Interaction													NS
													NS

Table 61. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on oil yield (q ha<sup>-1</sup>) and oil content (%) of mustard (*Brassica juncea* L.) cultivar Alankar and PBM16

Sampling stage (days after sowing)												
Cultivar	Oil yield						Oil content					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	2.51	2.69	3.67	3.36	3.23	3.09	32.98	33.33	34.42	34.08	33.83	33.73
PBM16	2.26	2.51	3.39	3.12	3.01	2.86	31.81	32.60	33.90	33.35	33.13	32.96
Mean	2.38	2.60	3.53	3.24	3.12		32.40	32.97	34.16	33.72	33.48	

L.S.D. at 5%

Oil yield			Oil content		
Spray	0.20	0.15			
Cultivar	0.13	0.09			
Interaction	NS	0.21			

Table 62 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on acid value, iodine value and Saponification value of mustard (*Brassica juncea* L ) cultivar Alankar and PBM16

Sampling stage (days after sowing)													
Cultivar	Acid value						Iodine value						
	Ethrel concentration (µL/L)												
	0	100	200	400	600	Mean	0	100	200	400	600	Mean	
	Alankar	2.46	2.65	2.98	2.84	2.78	2.74	92.41	93.34	95.17	94.12	93.88	93.79
PBM16	2.16	2.40	2.81	2.67	2.55	2.52	91.72	92.58	94.07	93.65	93.06	93.02	
Mean	2.31	2.53	2.89	2.76	2.66		92.06	92.96	94.62	93.89	93.47		
L S D at 5%													
Cultivar	Saponification value						Acid value						Saponif ication value
	Alankar	133.98	136.93	139.41	138.15	137.85	137.26						
	PBM16	131.68	133.91	136.42	135.15	135.10	134.45						
	Mean	132.83	135.42	137.91	136.65	136.48							
							Spray		0.06	NS	0.29		
							Cultivar		0.04	NS	0.19		
							Interaction		NS	NS	NS		

Table 63. Pooled analysis of plant dry weight in Experiment 1 and 2

Cultivar	Sampling stage (days after sowing)																													
	80						100																							
	Ethrel concentration (μL/L)																													
	0	100	200	400	600	Mean	0	100	200	400	600	Mean																		
Alankar	32.51	33.48	37.78	35.35	34.78	34.78	34.78	35.78	40.01	38.51	37.91	37.40																		
													31.17	32.80	36.18	34.58	34.24	33.79	34.02	35.01	39.33	37.67	37.38	36.68						
Non-Irrigated						Non-Irrigated																								
Alankar	26.37	28.15	32.46	31.04	30.32	29.67	28.23	30.19	35.63	34.07	33.43	32.31																		
PBM16	25.46	27.26	31.06	30.04	29.55	28.68	27.53	29.30	34.17	33.16	32.74	31.38																		
Mean	28.88	30.42	34.37	32.75	32.22		31.14	32.57	37.28	35.86	35.37																			
Alankar	36.99	38.05	43.11	41.61	40.99	40.15	40.15	Irrigation (A)	NS	NS	NS	NS																		
													36.21	37.15	42.46	40.77	40.49	39.42	0.15	0.18	0.18									
																						31.09	32.24	38.81	37.20	36.35	35.14	0.00024	0.00022	0.00059
33.48	34.80	40.50	38.97	38.44	0.0008	NS	NS	NS																						
									A X B X C	A X B X C	A X B X C	A X B X C	NS	NS	NS	NS														

Table 64. Pooled analysis of leaf fresh weight in Experiment 1 and 2

Sampling stage (days after sowing)												
Cultivar	80						100					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	44.86	47.00	55.71	51.32	49.70	49.72	32.22	33.68	40.51	38.54	36.34	36.26
PBM16	42.40	45.06	52.52	49.59	48.62	47.64	30.87	32.33	39.14	37.03	35.98	35.07
Alankar	29.74	32.36	36.68	36.93	35.65	34.67	20.70	22.67	29.05	27.43	26.27	25.22
PBM16	28.27	30.80	36.69	35.36	33.79	32.98	19.84	21.66	27.44	26.24	25.18	24.07
Mean	36.32	38.81	45.90	43.30	41.94		25.91	27.58	34.03	32.31	30.94	
L.S.D. at 5%												
Alankar	16.26	17.17	21.84	20.71	19.87	19.17	19.17	18.31	11.83	11.37	15.88	120
PBM16	15.49	16.23	20.95	19.74	19.17	18.31						
Alankar	9.65	10.26	13.70	13.02	12.53	11.83	11.83	11.37	0.0006	0.0027	0.00013	NS
PBM16	9.02	10.41	13.06	12.39	11.97	11.37						
Mean	12.61	13.52	17.39	16.46								

Table 65. Pooled analysis of leaf turgid weight in Experiment 1 and 2

[illegible]

Table 66. Pooled analysis of leaf relative water content in Experiment 1 and 2

Sampling stage (days after sowing)															
Cultivar	80														
	Ethrel concentration (µL/L)														
	0	100	200	400	600	Mean	0	100	200	400	600	Mean			
	Irrigated														
Alankar	61.55	63.11	67.27	65.21	64.18	64.26	51.91	53.01	57.23	55.75	54.83	54.54			
PBM16	60.19	61.16	65.71	63.88	63.45	62.88	50.51	51.61	55.76	54.32	53.41	53.12			
Alankar	46.43	47.55	49.15	48.58	48.05	47.95	37.79	39.12	42.36	41.26	40.34	40.18			
PBM16	45.05	46.29	48.22	47.65	46.17	46.68	37.07	38.32	41.31	40.06	39.06	39.17			
Mean	53.30	54.53	57.59	56.33	55.46		44.32	45.51	49.17	47.85	46.91				
120															
L.S.D. at 5%															
Alankar	45.89	47.00	50.68	49.45	48.47	48.30	Irrigation (A)						NS	NS	120
							Cultivar (B)								
PBM16	44.73	45.60	49.08	47.99	47.12	46.91	A X B						0.15	0.05	0.12
Alankar	30.02	30.49	32.20	31.76	31.28	31.15	Spray (C)						NS	NS	NS
PBM16	29.57	30.02	31.65	30.95	30.48	30.53	A X C						NS	NS	NS
Mean	37.55	38.28	40.90	40.04	39.34		B X C						0.0003	NS	NS
							A X B X C						NS	NS	NS



Table 67. Pooled analysis of rate of photosynthesis in Experiment 1 and 2

Cultivar	Sampling stage (days after sowing)											
	80						100					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	17.15	18.33	21.16	20.12	19.24	19.20	15.08	16.30	19.36	18.12	17.24	17.22
PBM16	16.22	17.28	20.14	19.14	18.25	18.21	14.35	15.27	18.50	17.30	16.36	16.36
Alankar	15.49	16.42	19.20	18.40	17.27	17.36	14.42	15.32	18.33	17.13	16.31	16.30
PBM16	14.23	15.37	18.61	17.42	16.34	16.39	13.36	14.44	17.27	16.45	15.21	15.35
Mean	15.77	16.85	19.78	18.77	17.77		14.30	15.33	18.36	17.25	16.28	

L.S.D. at 5%		
Irrigation (A)	NS	NS
Cultivar (B)	0.10	0.10
A X B	NS	NS
Spray (C)	0.0002	0.0001
A X C	NS	NS
B X C	NS	NS
A X B X C	NS	NS

Table 68. Pooled analysis of carboxylation efficiency in Experiment 1 and 2

Sampling stage (days after sowing)												
80												
Cultivar	Ethrel concentration (μL/L)											
	100											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	6.28	6.44	6.77	6.67	6.57	6.55	5.72	5.98	6.40	6.20	6.12	6.08
PBM16	6.40	6.26	6.68	6.53	6.43	6.46	5.56	5.71	6.28	6.07	5.61	5.84
Alankar	5.89	6.02	6.37	6.28	6.09	6.13	5.69	5.83	6.30	6.10	5.99	5.98
PBM16	5.60	5.84	6.32	6.15	5.98	5.98	5.45	5.70	6.08	6.00	5.77	5.80
Mean	6.04	6.14	6.53	6.41	6.27		5.60	5.80	6.26	6.09	5.87	

L.S.D. at 5%		
	80	100
Irrigation (A)	NS	NS
Cultivar (B)	0.08	0.08
A X B	NS	NS
Spray (C)	NS	NS
A X C	NS	NS
B X C	NS	NS
A X B X C	NS	NS

Table 69. Pooled analysis of photosynthetic water use efficiency in experiment 1 and 2

Cultivar	Sampling stage (days after sowing)											
	80						100					
	Ethrel concentration (μL/L)											
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	39.28	41.05	44.38	43.74	42.44	42.18	36.48	38.49	43.02	41.51	40.09	39.92
PBM16	38.32	39.58	44.10	42.54	41.80	41.26	35.29	36.65	42.37	40.54	39.26	38.82
Alankar	37.48	38.78	42.99	42.13	39.85	40.24	35.75	37.06	41.98	40.49	38.82	38.82
PBM16	35.27	37.18	42.94	40.51	39.21	39.02	33.96	35.80	40.79	39.48	38.02	37.61
Mean	37.59	39.14	43.60	42.23	40.82		35.37	37.00	42.04	40.51	39.05	

L.S.D. at 5%		
Irrigation (A)	NS	NS
Cultivar (B)	0.26	0.06
A X B	NS	NS
Spray (C)	0.001	0.002
A X C	NS	NS
B X C	0.003	NS
A X B X C	NS	NS

Table 70. Pooled analysis of plant water use efficiency in Experiment 1 and 2

Sampling stage (days after sowing)													
Cultivar	80												
	Ethrel concentration (μL/L)												
	0	100	200	400	600	Mean	0	100	200	400	600	Mean	
Alankar	2.54	2.60	2.84	2.68	2.64	2.66	3.30	3.38	Irrigated				3.49
									3.63	3.58	3.54	3.44	
PBM16	2.46	2.55	2.74	2.66	2.64	2.61	3.27	3.34					3.60
									Non-Irrigated				
Alankar	2.31	2.39	2.65	2.49	2.51	2.47	3.04	3.14	Non-Irrigated				3.28
									3.49	3.38	3.34	3.21	
PBM16	2.25	2.33	2.59	2.52	2.48	2.43	2.97	3.10					3.39
									Non-Irrigated				
Mean	2.39	2.47	2.71	2.59	2.57		3.14	3.24	3.53	3.45	3.42		

L.S.D. at 5%		
	80	100
Irrigation (A)	NS	NS
Cultivar (B)	0.02	0.01
A X B	NS	NS
Spray (C)	NS	NS
A X C	NS	NS
B X C	NS	NS
A X B X C	NS	NS

Table 71. Pooled analysis of pod number, seed yield and oil yield in Experiment 1 and 2

Cultivar	Pod number					Seed yield						
						Ethrel concentration (μL/L)						
	0	100	200	400	600	Mean	0	100	200	400	600	Mean
Alankar	144.33	148.67	160.33	156.33	153.67	152.67	10.33	10.98	13.55	12.73	12.41	12.00
PBM16	140.33	144.67	158.33	150.67	150.00	148.80	9.87	10.46	13.19	12.05	11.97	11.51
Alankar	125.00	129.00	144.00	139.00	136.67	134.73	7.62	8.08	10.67	9.86	9.53	9.15
PBM16	122.00	125.67	139.00	136.00	133.67	131.27	7.10	7.71	10.00	9.36	9.09	8.65
Mean	132.92	137.00	150.42	145.50	143.50		8.73	9.31	11.85	11.00	10.75	

L.S.D. at 5%											
Oil yield						Pod					
						number					
						yield					
						yield					
Alankar	3.63	3.90	5.02	4.65	4.45	4.33	Irrigation (A)	NS	NS	NS	NS
PBM16	3.43	3.66	4.77	4.32	4.22	4.08	A X B	NS	NS	NS	
											Non-Irrigated
Alankar	2.51	2.69	3.67	3.36	3.23	3.09	Spray (C)	0.17	NS	NS	
PBM16	2.26	2.51	3.39	3.12	3.01	2.86	A X C	NS	NS	NS	NS
Mean	2.96	3.19	4.21	3.87	3.73		B X C	NS	NS	NS	NS

Table 72. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant height (cm plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (µL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	84.63	92.47	101.83	92.98	84.37	99.67	113.47	99.17
N <sub>40</sub>	108.17	113.13	120.30	113.87	105.73	116.97	129.00	117.23
N <sub>60</sub>	114.70	121.37	128.37	121.48	113.87	125.30	138.73	125.97
N <sub>80</sub>	123.27	130.27	138.33	130.62	123.73	135.13	147.70	135.52
Mean	107.69	114.31	122.21		106.73	119.27	132.23	
L.S.D. at 5%								
N <sub>0</sub>	90.30	104.00	116.50	103.60		80	100	120
N <sub>40</sub>	109.77	122.07	134.73	122.19	Spray (S)	0.83	0.76	0.98
N <sub>60</sub>	119.37	131.13	143.67	131.39	Nitrogen (N)	0.96	0.88	1.14
N <sub>80</sub>	128.73	140.63	152.93	140.77	S x N	1.66	1.53	NS
Mean	112.04	124.46	136.96					

Table 73. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf area plant<sup>-1</sup> (cm<sup>2</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (µL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	2249.33	2764.67	3243.33	2752.44	2381.67	3113.33	3777.00	3090.67
N <sub>40</sub>	3260.33	3869.00	4425.33	3851.55	3501.33	4207.67	4968.33	4225.78
N <sub>60</sub>	3833.00	4483.33	5095.00	4470.44	4312.67	5123.67	5946.67	5127.67
N <sub>80</sub>	4625.67	5268.33	5817.00	5237.00	5203.67	6022.33	6790.00	6005.33
Mean	3412.08	4096.33	4645.17		3849.83	4616.75	5370.50	
L.S.D. at 5%								
	120							
N <sub>0</sub>	1313.33	1814.00	2311.00	1812.78		80	100	120
N <sub>40</sub>	2030.00	2552.67	3082.33	2555.00	Spray (S)	24.38	23.66	23.66
N <sub>60</sub>	2565.33	3107.00	3699.67	3124.00	Nitrogen (N)	28.15	27.32	27.32
N <sub>80</sub>	3282.33	3826.67	4380.33	3829.78	S x N	48.77	47.33	47.31
Mean	2297.75	2825.08	3368.33					

Table 74. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf area index of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	2.81	3.46	4.05	3.44	2.98	3.89
N <sub>40</sub>	4.07	4.84	5.53	4.81	4.38	5.26
N <sub>60</sub>	4.79	5.60	6.37	5.59	5.39	6.40
N <sub>80</sub>	5.78	6.59	7.27	6.55	6.50	7.53
Mean	4.36	5.12	5.80		4.81	5.77
120						
L.S.D. at 5%						
N <sub>0</sub>	1.64	2.27	2.89	2.27	80	100
N <sub>40</sub>	2.54	3.19	3.85	3.19	Spray (S)	0.030
N <sub>60</sub>	3.21	3.88	4.62	3.90	Nitrogen (N)	0.035
N <sub>80</sub>	4.10	4.78	5.47	4.78	S x N	0.061
Mean	2.87	3.53	4.21			0.059



Table 75 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on specific leaf area ( $\text{cm}^2\text{g}^{-1}$ ) of mustard (*Brassica juncea* L ) cultivar Alankai grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethiel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	280 46	296 00	309 18	295 21	387 89	420 33	443 83	417 35
N <sub>40</sub>	296 39	315 33	324 43	312 05	414 85	435 67	453 31	434 61
N <sub>60</sub>	315 73	325 00	332 57	324 43	428 27	446 00	460 98	445 08
N <sub>80</sub>	325 52	333 00	340 17	332 90	440 98	457 00	470 87	456 28
Mean	304 52	317 33	326 59			439 75	457 25	
L S D at 5%								
120								
N <sub>0</sub>	419 59	441 00	453 14	437 91		80	100	120
N <sub>40</sub>	454 14	465 33	473 48	464 32	Spray (S)	0 67	0 67	0 88
N <sub>60</sub>	471.56	478 33	485 52	478 49	Nitrogen (N)	0 76	0 77	1.02
N <sub>80</sub>	486 99	492 33	496 09	491 80	S x N	NS	NS	NS
Mean		469 25	477 06					

Table 76 Effect of ethiel spray at 60d after sowing (DAS, flowering stage) on specific leaf weight (mg cm<sup>2</sup>) of mustard (*Brassica juncea* L ) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	3 56	3 37	3 23	3 39	2 58	2 38
N <sub>40</sub>	3 37	3 17	3 08	3 21	2 41	2 29
N <sub>60</sub>	3 17	3 07	3 01	3 08	2 33	2 24
N <sub>80</sub>	3 07	3 00	2 94	3 00	2 27	2 19
Mean	3 29	3 15	3 06		2 40	2 27
120						
L S D at 5%						
N <sub>0</sub>	2 38	2 27	2 21	2 29	80	100
N <sub>40</sub>	2 20	2 15	2 11	2 15	Spray (S)	0 006
N <sub>60</sub>	2 12	2 09	2 06	2 09	Nitrogen (N)	0 007
N <sub>80</sub>	2 05	2 03	2 01	2 03	S x N	NS
Mean	2 19	2 13	2 10			
					NS	NS
						0 005
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006
						0 006

Table 77. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	25.35	29.18	32.59	29.04	28.45	32.34	35.68	32.16
N <sub>40</sub>	33.68	37.22	40.92	37.27	36.78	40.33	43.99	40.37
N <sub>60</sub>	35.81	39.90	43.97	39.89	39.77	43.83	47.87	43.82
N <sub>80</sub>	39.67	43.53	47.03	43.41	44.77	48.63	52.12	48.51
Mean	33.63	37.46	41.13		37.44	41.28	44.92	
L.S.D. at 5%								
120								
N <sub>0</sub>	32.84	36.39	39.81	36.35		80	100	120
N <sub>40</sub>	40.84	44.39	48.09	44.44	Spray (S)	0.18	0.18	0.14
N <sub>60</sub>	44.73	48.76	52.77	48.75	Nitrogen (N)	0.21	0.21	0.17
N <sub>80</sub>	50.82	54.65	58.45	54.64	S x N	0.37	0.36	0.29
Mean	42.31	46.05	49.78					

Table 78. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	8.02	9.33	10.49	9.28	6.14	7.40	8.51	7.35
N <sub>40</sub>	11.00	12.26	13.64	12.30	8.44	9.66	10.96	9.68
N <sub>60</sub>	12.14	13.79	15.32	13.75	10.07	11.50	12.90	11.49
N <sub>80</sub>	14.21	15.82	17.10	15.71	11.80	13.20	14.42	13.14
Mean	11.34	12.80	14.14		9.11	10.44	11.70	
L.S.D. at 5%								
120								
N <sub>0</sub>	3.13	4.12	5.10	4.12		80	100	120
N <sub>40</sub>	4.47	5.48	6.51	5.49	Spray (S)	0.08	0.06	0.05
N <sub>60</sub>	5.44	6.49	7.62	6.52	Nitrogen (N)	0.09	0.06	0.06
N <sub>80</sub>	6.74	7.77	8.83	7.78	S x N	0.15	0.11	NS
Mean	4.94	5.97	7.02					

Table 79. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stem dry weight (g) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	15.37	17.25	18.86	17.16	17.96	19.48	20.84	19.43
N <sub>40</sub>	19.36	20.92	22.50	20.93	21.77	23.09	24.38	23.08
N <sub>60</sub>	19.75	21.36	23.07	21.39	22.20	23.64	25.11	23.65
N <sub>80</sub>	20.45	22.04	23.64	22.04	23.89	25.25	26.54	25.23
Mean	18.74	20.39	22.02		21.45	22.87	24.22	
L.S.D. at 5%								
120								
N <sub>0</sub>	22.49	23.76	24.99	23.75		80	100	120
N <sub>40</sub>	26.55	27.80	29.08	27.81	Spray (S)	0.09	0.09	0.06
N <sub>60</sub>	27.86	29.28	30.68	29.28	Nitrogen (N)	0.11	0.11	0.07
N <sub>80</sub>	30.24	31.54	32.80	31.53	S x N	NS	NS	0.11
Mean	26.79	28.10	29.39					

Table 80. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on pod dry weight (g) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	1.95	2.60	3.24	2.60	4.35	5.45	6.43	5.41
N <sub>40</sub>	3.32	4.04	4.78	4.05	6.58	7.58	8.66	7.60
N <sub>60</sub>	3.92	4.78	5.58	4.76	7.50	8.69	9.86	8.69
N <sub>80</sub>	5.01	5.68	6.29	5.66	9.08	10.18	11.15	10.14
Mean	3.55	4.28	4.97		6.88	7.98	9.03	
L.S.D. at 5%								
120								
N <sub>0</sub>	7.22	8.51	9.72	8.49		80	100	120
N <sub>40</sub>	9.83	11.10	12.50	11.14	Spray (S)	0.03	0.04	0.06
N <sub>60</sub>	11.43	12.98	14.47	12.96	Nitrogen (N)	0.04	0.05	0.06
N <sub>80</sub>	13.84	15.34	16.82	15.33	S x N	0.06	0.09	0.11
Mean	10.58	11.98	13.38					

Table 81. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	31.64	31.97	32.19	31.93	21.58	22.90	23.85	22.78
N <sub>40</sub>	32.66	32.94	33.33	32.98	22.95	23.95	24.91	23.94
N <sub>60</sub>	33.90	34.53	34.84	34.42	25.32	26.24	26.95	26.02
N <sub>80</sub>	35.82	36.33	36.36	36.17	26.36	27.15	27.96	27.16
Mean	33.50	33.94	34.18		24.05	25.06	25.92	
L.S.D. at 5%								
	120							
N <sub>0</sub>	9.52	11.32	12.81	11.22		80	100	120
N <sub>40</sub>	10.94	12.35	13.54	12.28	Spray (S)	0.07	0.06	0.08
N <sub>60</sub>	12.16	13.32	14.44	13.31	Nitrogen (N)	0.08	0.07	0.09
N <sub>80</sub>	13.26	14.22	15.11	14.20	S x N	0.15	0.12	NS
Mean	11.47	12.80	13.97					

Table 82. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stem dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	60.63	59.12	57.87	59.21	63.13	60.25	58.41	60.60
N <sub>40</sub>	57.48	56.20	54.98	56.22	59.19	57.25	55.42	57.29
N <sub>60</sub>	55.15	53.49	52.47	53.70	57.82	53.92	52.45	54.73
N <sub>80</sub>	51.55	50.62	50.26	50.81	53.36	51.92	50.92	52.09
Mean	56.20	54.86	53.89		58.37	55.84	54.30	
L.S.D. at 5%								
120								
N <sub>0</sub>	68.48	65.28	62.77	66.51		80	100	120
N <sub>40</sub>	65.01	62.64	60.49	62.71	Spray (S)	0.10	0.10	0.11
N <sub>60</sub>	62.28	60.06	58.14	60.16	Nitrogen (N)	0.12	0.11	0.13
N <sub>80</sub>	60.14	57.72	56.12	57.99	S x N	0.21	0.19	NS
Mean	63.98	61.43	59.38					



Table 83. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on pod dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	7.69	8.91	9.94	8.85	15.29	16.85	18.02	16.72
N <sub>40</sub>	9.85	10.85	11.68	10.79	17.89	18.79	19.68	18.79
N <sub>60</sub>	10.94	11.98	12.69	11.87	19.61	19.83	20.60	20.01
N <sub>80</sub>	12.63	13.04	13.37	13.01	20.28	20.93	21.39	20.87
Mean	10.28	11.20	11.92		18.27	19.10	19.92	
L.S.D. at 5%								
120								
N <sub>0</sub>	21.90	23.39	24.41	23.23		80	100	120
N <sub>40</sub>	24.07	25.01	26.00	25.02	Spray (S)	0.05	0.05	0.06
N <sub>60</sub>	25.55	26.62	27.42	26.53	Nitrogen (N)	0.05	0.06	0.07
N <sub>80</sub>	27.23	28.06	28.78	28.02	S x N	0.09	0.10	0.13
Mean	24.71	25.77	58.45					

Table 84 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf fresh weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethiel concentration (µL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	34.67	41.34	47.34	41.12	25.26	30.58	35.26	30.37
N <sub>40</sub>	49.77	56.22	63.45	56.48	36.26	41.43	47.50	41.73
N <sub>60</sub>	56.08	64.43	72.37	64.29	44.14	50.39	56.48	50.34
N <sub>80</sub>	66.77	75.38	82.06	74.74	52.82	59.11	64.39	58.77
Mean	51.82	59.34	66.31		39.62	45.38	50.91	
120								
L S D at 5%								
N <sub>0</sub>	12.32	16.17	20.02	16.17		80	100	120
N <sub>40</sub>	18.41	22.41	26.51	22.45	Spray (S)	0.47	0.33	0.23
N <sub>60</sub>	22.86	27.09	31.70	27.22	Nitrogen (N)	0.54	0.38	0.26
N <sub>80</sub>	28.22	32.37	36.66	32.42	S x N	0.94	0.65	NS
Mean	20.45	24.51	28.73					

Table 85. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf turgid weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (µL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	53.05	59.71	65.46	59.41	45.33	50.34	54.29	49.99
N <sub>40</sub>	70.30	76.37	83.68	76.78	58.61	63.27	69.07	63.65
N <sub>60</sub>	76.09	84.79	92.77	84.55	68.18	74.18	79.95	74.10
N <sub>80</sub>	86.54	94.90	101.32	94.25	77.42	83.16	87.77	82.78
Mean	71.49	78.94	85.81		62.38	67.74	72.77	
L.S.D. at 5%								
	120							
N <sub>0</sub>	25.20	29.25	3.20	29.22		80	100	120
N <sub>40</sub>	33.38	37.56	41.81	37.58	Spray (S)	0.50	0.50	0.34
N <sub>60</sub>	39.40	43.83	48.76	44.00	Nitrogen (N)	0.57	0.58	0.39
N <sub>80</sub>	46.31	50.50	54.89	50.57	S x N	0.99	1.00	NS
Mean	36.07	40.29	44.67					

Table 86 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf relative water content (%) of mustard (*Brassica juncea* L ) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	59.02	63.53	67.04	63.20	48.84	53.97	58.43	53.75
N <sub>40</sub>	65.38	68.57	71.12	68.36	55.45	59.25	62.88	59.19
N <sub>60</sub>	68.74	71.32	73.66	71.24	58.63	62.03	65.00	61.88
N <sub>80</sub>	72.66	75.32	77.13	75.04	62.51	65.60	68.12	65.41
Mean	66.45	69.69	72.24		56.38	60.21	63.61	
L.S.D. at 5%								
120								
N <sub>0</sub>	40.36	47.96	52.53	46.95		80	100	120
N <sub>40</sub>	48.22	52.74	56.66	52.54	Spray (S)	0.35	0.10	0.16
N <sub>60</sub>	51.29	55.20	58.11	54.87	Nitrogen (N)	0.40	0.12	0.19
N <sub>80</sub>	54.28	57.58	58.76	56.87	S x N	NS	NS	NS
Mean	48.54	53.37	56.51					

Table 87. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on rate of photosynthesis ( $\mu\text{mol m}^{-2}\text{s}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	15.10	17.47	19.87	17.48	13.14	15.58
N <sub>40</sub>	18.24	20.96	24.07	21.09	16.16	19.13
N <sub>60</sub>	21.00	23.93	26.64	23.86	19.23	22.25
N <sub>80</sub>	23.14	25.73	28.32	25.73	22.15	24.90
Mean	19.37	22.02	24.73		17.67	20.47
L.S.D. at 5%						
			80		100	
Spray (S)			0.15		0.23	
Nitrogen (N)			0.18		0.26	
S x N			0.31		0.45	

Table 88 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stomatal conductance ( $\text{mol m}^{-2} \text{s}^{-1}$ ) of mustard (*Brassica juncea* L ) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	0.42	0.44	0.46	0.44	0.40	0.41	0.43	0.41
N <sub>40</sub>	0.45	0.47	0.49	0.47	0.43	0.45	0.47	0.45
N <sub>60</sub>	0.48	0.50	0.52	0.50	0.46	0.48	0.51	0.48
N <sub>80</sub>	0.51	0.53	0.55	0.53	0.50	0.52	0.54	0.52
Mean	0.46	0.48	0.50		0.45	0.47	0.49	
L S D. at 5%								

Table 89 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on internal carbon dioxide concentration of mustard (*Brassica juncea* L ) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	257 80	272 47	287 60	272 62	240 97	257 30	272 27	256 84
N <sub>40</sub>	291 60	310 50	333 53	311 88	272 50	294 90	315 53	294 31
N <sub>60</sub>	324 23	345 53	365 10	344 96	308 73	330 83	351 57	330 38
N <sub>80</sub>	351 37	368 50	384 63	368 17	342 23	361 23	377 13	360 20
Mean	306 25	324 25	342 72		291 11	311 07	329 13	
L S D at 5%								
					80                      100			
					Spray (S)            1 99            2 56			
					Nitrogen (N)        2 30            2 95			
					S x N                    3 99            5 12			

Table 90 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on transpiration rate ( $\text{kg m}^{-2} \text{ day}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80			100				
	Ethiel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	11.92	12.34	12.71	12.32	9.18	9.61	9.97	9.59
N <sub>40</sub>	12.89	13.25	13.60	13.24	10.60	10.97	11.32	10.96
N <sub>60</sub>	13.17	13.69	14.11	13.66	10.84	11.29	11.72	11.28
N <sub>80</sub>	13.81	14.23	14.59	14.21	11.38	11.74	12.07	11.73
Mean	12.95	13.38	13.75		10.50	10.90	11.27	
L S D at 5%								
80100								
Spray (S)0.040.03								
Nitrogen (N)0.040.03								
S x N0.080.06								



Table 91 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on carboxylation efficiency (%) of mustard (*Brassica juncea* L ) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)								
	80				100				
	Ethiel concentration (μL/L)								
	0	100	200	Mean	0	100	200	Mean	
N <sub>0</sub>	5.85	6.41	6.91	6.39	5.45	6.05	6.59	6.03	
N <sub>40</sub>	6.28	6.74	7.22	6.74	5.93	6.48	6.99	6.47	
N <sub>60</sub>	6.48	6.92	7.30	6.90	6.23	6.72	7.19	6.71	
N <sub>80</sub>	6.58	6.98	7.36	6.97	6.47	6.88	7.27	6.87	
Mean	6.29	6.76	7.20		6.02	6.54	7.01		
L S D at 5%									
				80					100
				Spray (S)					0.05
				Nitrogen (N)					0.06
				S x N					NS

Table 92. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on photosynthetic water use efficiency ( $\mu\text{mol mol}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	35.95	39.69	43.19	39.61	32.85	37.69
N <sub>40</sub>	40.35	44.59	49.12	44.69	37.58	42.82
N <sub>60</sub>	43.75	47.54	51.23	47.50	41.80	46.03
N <sub>80</sub>	45.37	48.86	51.49	48.57	44.30	47.59
Mean	41.35	45.17	48.76		39.13	43.53
L.S.D. at 5%						
	80			100		
	Spray (S)			0.34		
	Nitrogen (N)			0.39		
	S x N			0.67		
				NS		

Table 93. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on water use efficiency ( $\text{mg g}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	2.13	2.37	2.56	2.35	3.10	3.36
N <sub>40</sub>	2.61	2.80	3.00	2.80	3.47	3.58
N <sub>60</sub>	2.72	2.91	3.12	2.92	3.67	3.75
N <sub>80</sub>	2.87	3.06	3.22	3.05	3.93	4.08
Mean	2.58	2.79	2.97		3.54	3.77
L.S.D. at 5%						
80100						
Spray (S)0.010.01						
Nitrogen (N)0.010.01						
S x NNSNS						

Table 94. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on 1-aminocyclopropane-1-carboxylic acid (ACC) ( $\eta$  mol g fresh weight<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	17.27	19.77	22.19	19.74	19.56	22.32
N <sub>40</sub>	20.99	23.24	26.01	23.41	23.72	26.09
N <sub>60</sub>	24.50	26.99	29.41	26.96	26.92	30.15
N <sub>80</sub>	28.07	30.78	33.16	30.67	31.31	33.89
Mean	22.71	25.19	27.69		25.38	28.11
L.S.D. at 5%						
80100						
Spray (S)0.120.22						
Nitrogen (N)0.140.25						
S x NNS0.43						

Table 95. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on ACC oxidase ( $\eta$  L g<sup>-1</sup> fresh weight h<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80			100				
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	Mean		
N <sub>0</sub>	57.98	63.85	69.75	63.86	70.21	78.56	87.07	78.61
N <sub>40</sub>	80.05	80.01	89.75	83.27	95.28	105.44	114.02	104.91
N <sub>60</sub>	89.50	97.26	104.96	97.24	108.10	116.54	124.99	116.55
N <sub>80</sub>	104.11	111.90	119.46	111.83	121.37	130.80	141.02	131.06
Mean	82.91	88.26	95.98		98.74	107.84	116.77	
L.S.D. at 5%								
80100								
Spray (S)1.471.45								
Nitrogen (N)1.701.67								
S x N2.94NS								

Table 96. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on ethylene evolution ( $\eta\text{l h}^{-1} \text{ plant}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Sampling days (after sowing)						
Nitrogen (kg/ha)	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	9.27	11.15	12.02	10.81	11.98	13.12
N <sub>40</sub>	17.94	17.60	19.14	18.23	19.43	21.17
N <sub>60</sub>	21.97	24.00	25.83	23.93	25.22	26.98
N <sub>80</sub>	28.89	30.59	32.75	30.74	32.32	34.36
Mean	19.52	20.84	22.43		22.24	23.91
L.S.D. at 5%						
80100						
Spray (S)0.480.50						
Nitrogen (N)0.550.58						
S x NNSNS						

Table 97. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrate reductase activity ( $\mu\text{mol NO}_2^- \text{g}^{-1} \text{h}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)						.	
	80			100				
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100		200
N <sub>0</sub>	0.14	0.17	0.20	0.17	0.22	0.22	0.26	0.23
N <sub>40</sub>	0.40	0.40	0.42	0.40	0.44	0.46	0.49	0.46
N <sub>60</sub>	0.46	0.49	0.53	0.49	0.53	0.56	0.62	0.57
N <sub>80</sub>	0.58	0.62	0.64	0.61	0.67	0.70	0.73	0.70
Mean	0.39	0.42	0.45		0.46	0.49	0.53	
L.S.D. at 5%								
80100								
Spray (S)0.010.01								
Nitrogen (N)0.010.01								
S x NNSNS								

Table 98 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrogen content (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	2.15	2.30	2.43	2.29	1.97	2.20	2.37	2.18
N <sub>40</sub>	2.46	2.58	2.69	2.58	2.19	2.36	2.52	2.36
N <sub>60</sub>	2.65	2.76	2.88	2.76	2.33	2.49	2.63	2.48
N <sub>80</sub>	2.87	2.96	3.00	2.94	2.51	2.65	2.75	2.64
Mean	2.53	2.65	2.75		2.25	2.43	2.57	
L.S.D. at 5%								
120								
N <sub>0</sub>	1.72	1.92	2.07	1.90		80	100	120
N <sub>40</sub>	2.08	2.19	2.27	2.18	Spray (S)	0.01	0.01	0.01
N <sub>60</sub>	2.15	2.25	2.32	2.24	Nitrogen (N)	0.02	0.01	0.01
N <sub>80</sub>	2.21	2.29	2.35	2.28	S x N	0.03	NS	0.02
Mean	2.04	2.16	2.25					



Table 99 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrogen accumulation (mg plant<sup>-1</sup>) of mustard (*Brassica juncea* L ) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	544 33	672 00	791 67	669 33	561 67	712 33	846 67	706 89
N <sub>40</sub>	829 33	960 33	1099 67	963 11	806 67	953 00	1105 33	955 00
N <sub>60</sub>	947 67	1099 67	1266 00	1104 44	927 67	1093 00	1257 67	1092 78
N <sub>80</sub>	1140 00	1289 67	1411 67	1280 44	1123 67	1288 67	1434 67	1282 33
Mean	865 33	1005 42	1142 25		854 92	1011 75	1161 08	
120								
L S D at 5%								
N <sub>0</sub>	566 67	700 00	826 33	697 67		80	100	120
N <sub>40</sub>	849 67	970 33	1093 67	971 22	Spray (S)	6 50	7 40	4 90
N <sub>60</sub>	964 33	1096 67	1226 33	1095 78	Nitrogen (N)	7 50	8 54	5 66
N <sub>80</sub>	1121 67	1257 33	1375 33	1251 44	S x N	12 99	14 79	NS
Mean	875 58	1006 08	1130 42					

Table 100. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on number of pod plant<sup>-1</sup>, number of seeds pod<sup>-1</sup> and 1000 seed weight (g) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Pods plant <sup>-1</sup>			Seeds pod <sup>-1</sup>					
	Ethrel concentration (μL/L)								
	0	100	200	Mean	0	100	200	Mean	
N <sub>0</sub>	138.00	146.33	153.33	145.89	12.45	13.23	13.95	13.21	
N <sub>40</sub>	159.30	166.33	173.00	166.22	13.83	14.52	15.23	14.53	
N <sub>60</sub>	172.67	181.67	189.00	181.44	15.10	15.82	16.52	15.82	
N <sub>80</sub>	190.33	197.33	203.67	197.11	16.36	17.07	17.76	17.06	
Mean	165.07	172.92	179.75		14.44	15.16	15.86		
L.S.D. at 5%									
1000 seed weight									
N <sub>0</sub>	4.28	4.41	4.51	4.40	Pods plant <sup>-1</sup>		Seeds pod <sup>-1</sup>	1000 seed weight	
N <sub>40</sub>	4.46	4.52	4.56	4.51					
N <sub>60</sub>	4.61	4.62	4.63	4.62	Spray (S)		0.71	0.05	0.01
N <sub>80</sub>	4.70	4.69	4.69	4.69	Nitrogen (N)		0.82	0.06	0.01
Mean	4.51	4.56	4.60		S x N		NS	NS	NS

Table 101. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on seed yield ( $\text{q ha}^{-1}$ ), biological yield ( $\text{q ha}^{-1}$ ) and harvest index (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Seed yield				Biological yield			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	9.19	10.67	12.05	10.64	42.81	45.49	48.01	45.44
N <sub>40</sub>	12.27	13.65	15.00	13.64	52.80	55.49	58.36	55.55
N <sub>60</sub>	15.03	16.58	18.07	16.56	57.67	60.95	64.22	60.94
N <sub>80</sub>	18.29	19.74	21.20	19.74	65.27	68.31	71.28	68.29
Mean	13.70	15.16	16.58		54.64	57.56	60.47	
L.S D. at 5%								
Harvest index								
N <sub>0</sub>	21.46	23.44	25.10	23.33	Seed yield		Biological	Harvest
N <sub>40</sub>	23.24	24.60	25.70	24.51			yield	index
N <sub>60</sub>	26.06	27.20	28.13	27.13	Spray (S)		0.08	0.08
N <sub>80</sub>	28.02	28.90	29.74	28.89	Nitrogen (N)		0.09	0.09
Mean	24.69	26.04	27.17		S x N		NS	0.16

Table 102. Effect of ethrel spray at 60d (post flowering stage) on seed nitrogen content ( $\text{mg g}^{-1} \text{ plant}^{-1}$ ), nitrogen harvest index and nitrogen yield potential of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Seed nitrogen content				Nitrogen harvest index			
	Ethrel concentration ( $\mu\text{L/L}$ )							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	32.37	40.07	46.73	39.72	0.06	0.06	0.06	0.06
N <sub>40</sub>	43.37	50.07	56.50	49.98	0.05	0.05	0.05	0.05
N <sub>60</sub>	45.73	52.07	58.93	52.24	0.05	0.05	0.05	0.05
N <sub>80</sub>	50.30	58.50	65.07	57.96	0.05	0.05	0.05	0.05
Mean	42.94	50.18	56.81		0.05	0.05	0.05	
L.S.D. at 5%								
	Nitrogen yield merit							
N <sub>0</sub>	1.94	2.40	2.80	2.38	Seed nitrogen content	Nitrogen harvest index	Nitrogen yield potential	
N <sub>40</sub>	2.17	2.50	2.82	2.50				
N <sub>60</sub>	2.29	2.60	2.95	2.61	Spray (S)	NS	0.01	
N <sub>80</sub>	2.51	2.92	3.25	2.89	Nitrogen (N)	NS	0.01	
Mean	2.23	2.61	2.95		S x N	NS	0.02	

Table 103. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on oil yield (q ha<sup>-1</sup>) and oil content (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Oil yield				Oil content			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	3.27	3.94	4.60	3.94	35.55	36.92	38.21	36.89
N <sub>40</sub>	4.29	4.95	5.63	4.96	34.99	36.26	37.52	36.26
N <sub>60</sub>	5.12	5.85	6.61	5.86	34.07	35.31	36.60	35.33
N <sub>80</sub>	6.02	6.78	7.55	6.78	32.94	34.32	35.61	34.29
Mean	4.67	5.38	6.10		34.39	35.70	36.99	
L.S.D. at 5%								
Oil yield    Oil content								
Spray (S)		0.04		0.06				
Nitrogen (N)		0.04		NS				
S x N		NS		NS				

Table 104. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on acid value, iodine value and saponification value of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Acid value				Iodine value			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	3.58	3.90	4.21	3.90	100.12	102.40	104.07	102.19
N <sub>40</sub>	3.36	3.74	4.04	3.71	98.75	100.83	102.39	100.66
N <sub>60</sub>	3.03	3.41	3.76	3.40	97.07	98.89	100.30	98.75
N <sub>80</sub>	2.81	3.12	3.44	3.12	95.70	97.50	98.93	97.37
Mean	3.19	3.54	3.86		97.91	99.91	101.42	
L.S.D. at 5%								
Saponification value								
N <sub>0</sub>	143.19	146.15	149.47	146.27	Acid value		Iodine value	Saponifica- tion value
N <sub>40</sub>	149.91	152.95	155.53	152.80				
N <sub>60</sub>	152.87	156.42	159.29	156.19	Spray (S)		0.04	0.31
N <sub>80</sub>	156.77	159.83	162.73	159.78	Nitrogen (N)		NS	NS
Mean	150.69	153.84	156.76		S x N		NS	NS

Table 105. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant height (cm plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	58.33	62.83	68.23	63.13	61.60	68.93	75.73	68.76
N <sub>40</sub>	75.73	80.87	85.93	80.84	77.27	84.87	91.93	84.69
N <sub>60</sub>	83.63	90.13	96.10	89.96	86.73	94.13	101.70	94.19
N <sub>80</sub>	94.37	100.93	106.43	100.58	97.37	103.90	112.70	104.66
Mean	78.02	83.69	89.18		80.74	87.96	95.52	
L.S.D. at 5%								
120								
N <sub>0</sub>	71.20	75.17	81.03	75.80		80	100	120
N <sub>40</sub>	84.97	89.77	95.77	90.17	Spray (S)	0.61	0.82	0.95
N <sub>60</sub>	92.13	99.00	106.10	99.08	Nitrogen (N)	0.71	0.94	1.10
N <sub>80</sub>	103.87	110.27	117.10	110.41	S x N	1.23	NS	NS
Mean	88.04	93.55	100.00					





Table 107. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf area index of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80			100				
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	2.03	2.59	3.12	2.58	2.26	3.02	3.73	3.00
N <sub>40</sub>	2.99	3.55	4.08	3.54	3.26	4.00	4.75	4.01
N <sub>60</sub>	3.38	3.97	4.55	3.97	3.79	4.58	5.34	4.57
N <sub>80</sub>	3.89	4.48	5.06	4.48	4.41	5.17	4.94	5.17
Mean	3.07	3.65	4.20		3.43	4.19	4.32	
L.S.D. at 5%								
120								
N <sub>0</sub>	1.19	1.69	2.15	1.68		80	100	120
N <sub>40</sub>	1.82	2.28	2.77	2.29	Spray (S)	0.03	0.02	0.02
N <sub>60</sub>	2.36	2.87	3.35	2.86	Nitrogen (N)	0.03	0.03	0.02
N <sub>80</sub>	2.97	3.43	3.91	3.44	S x N	NS	NS	NS
Mean	2.08	2.57	3.04					

Table 108. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on specific leaf area ( $\text{cm}^2 \text{g}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	0	100	200	Mean	0	100
Ethrel concentration ( $\mu\text{L/L}$ )						
N <sub>0</sub>	248.50	264.67	276.83	263.33	359.56	387.00
N <sub>40</sub>	274.46	286.00	293.70	284.72	386.24	403.00
N <sub>60</sub>	2.86.93	294.33	302.96	294.74	397.95	412.33
N <sub>80</sub>	299.87	306.33	312.25	306.32	409.13	421.00
Mean	277.44	287.83	296.43		388.22	405.83
L.S.D. at 5%						
120						
N <sub>0</sub>	370.82	411.67	418.18	400.22		80
N <sub>40</sub>	390.95	429.33	449.39	423.22	Spray (S)	0.69
N <sub>60</sub>	419.55	440.67	457.13	439.12	Nitrogen (N)	0.79
N <sub>80</sub>	435.90	453.00	467.11	452.00	S x N	NS
Mean	404.30	433.67	447.95			NS

Table 109. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on specific leaf weight ( $\text{mg cm}^{-2}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	4.02	3.78	3.61	3.80	2.78	2.61
N <sub>40</sub>	3.64	3.50	3.40	3.51	2.59	2.49
N <sub>60</sub>	3.48	3.40	3.30	3.39	2.51	2.43
N <sub>80</sub>	3.33	3.26	3.20	3.26	2.44	2.38
Mean	3.62	3.48	3.38		2.58	2.39
L.S.D. at 5%						
N <sub>0</sub>	2.70	2.43	2.28	2.47	80	120
N <sub>40</sub>	2.56	2.33	2.22	2.37	Spray (S)	0.009
N <sub>60</sub>	2.38	2.27	2.19	2.28	Nitrogen (N)	0.011
N <sub>80</sub>	2.29	2.20	2.14	2.21	S x N	NS
Mean	2.48	2.31	2.21		NS	NS

Table 110. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	21.56	25.25	28.69	25.17	24.64	28.38	31.79	28.27
N <sub>40</sub>	27.65	31.06	34.41	31.04	30.78	34.10	37.66	34.18
N <sub>60</sub>	29.16	32.82	36.26	32.75	33.28	36.93	40.39	36.87
N <sub>80</sub>	31.03	34.49	37.96	34.49	36.13	39.56	43.02	39.57
Mean	27.35	30.90	34.33		31.21	34.74	38.22	
L.S.D. at 5%								
120								
N <sub>0</sub>	27.71	31.44	34.85	31.33		80	100	120
N <sub>40</sub>	34.03	37.21	40.81	37.35	Spray (S)	0.15	0.16	0.15
N <sub>60</sub>	38.47	42.11	45.48	42.02	Nitrogen (N)	0.17	0.18	0.18
N <sub>80</sub>	42.28	45.64	49.16	45.69	S x N	NS	NS	NS
Mean	35.62	39.10	42.57					

Table 111. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	6.55	7.83	9.02	7.80	5.02	6.23	7.38	6.21
N <sub>40</sub>	8.72	9.92	10.12	9.92	6.76	7.93	9.15	7.95
N <sub>60</sub>	9.444	10.79	12.03	10.75	7.63	8.88	10.09	8.87
N <sub>80</sub>	10.40	11.71	12.98	11.70	8.62	9.83	11.02	9.82
Mean	8.78	10.06	11.29		7.01	8.22	9.41	
L.S.D. at 5%								
120								
N <sub>0</sub>	2.57	3.29	3.92	3.26		80	100	120
N <sub>40</sub>	3.72	4.25	4.94	4.27	Spray (S)	0.06	0.05	0.03
N <sub>60</sub>	4.50	5.21	5.87	5.19	Nitrogen (N)	0.07	0.06	0.04
N <sub>80</sub>	5.46	6.04	6.70	6.07	S x N	NS	NS	0.07
Mean	4.04	4.70	5.36					

Table 112. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stem dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	13.62	15.41	17.05	15.36	16.10	17.61	18.93	17.55
N <sub>40</sub>	16.73	18.30	19.85	18.29	19.15	20.38	21.74	20.42
N <sub>60</sub>	17.14	18.75	20.27	18.73	20.02	21.39	22.70	21.37
N <sub>80</sub>	17.54	19.09	20.67	19.10	21.06	22.35	23.67	22.36
Mean	16.26	17.89	19.46		19.08	20.43	21.76	
L.S.D. at 5%								
120								
N <sub>0</sub>	19.40	21.14	22.76	21.10		80	100	120
N <sub>40</sub>	22.86	24.28	25.99	24.38	Spray (S)	0.06	0.08	0.08
N <sub>60</sub>	25.06	26.69	28.24	26.67	Nitrogen (N)	0.07	0.09	0.09
N <sub>80</sub>	26.46	28.04	29.71	28.07	S x N	0.13	NS	NS
Mean	23.45	25.04	26.67					

Table 113. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on pod dry weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	1.39	2.01	2.63	2.01	3.51	4.53	5.49	4.51
N <sub>40</sub>	2.31	2.84	3.44	2.86	4.87	5.79	6.77	5.81
N <sub>60</sub>	2.58	3.28	3.94	3.26	5.63	6.63	7.59	6.62
N <sub>80</sub>	3.08	3.69	4.31	3.70	6.44	7.38	8.33	7.39
Mean	2.34	2.95	3.58		5.12	6.08	7.44	
L.S.D. at 5%								
120								
N <sub>0</sub>	5.74	7.01	8.17	6.98		80	100	120
N <sub>40</sub>	7.55	8.65	9.89	8.70	Spray (S)	0.04	0.04	0.05
N <sub>60</sub>	8.91	10.20	11.37	10.16	Nitrogen (N)	0.04	0.05	0.06
N <sub>80</sub>	10.50	11.56	12.75	11.60	S x N	NS	NS	0.10
Mean	8.17	9.36	10.54					

Table 114. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)									
	80				100					
	Ethrel concentration (μL/L)									
	0	100	200	Mean	0	100	200	Mean		
N <sub>0</sub>	30.38	31.00	31.44	30.94	20.37	21.96	23.21	21.85		
N <sub>40</sub>	31.54	31.94	32.32	31.93	21.96	23.26	24.30	23.17		
N <sub>60</sub>	32.37	32.89	33.18	32.81	22.93	24.05	24.98	23.99		
N <sub>80</sub>	33.51	33.94	34.19	33.88	23.86	24.84	25.61	24.77		
Mean	31.95	32.44	32.78		22.28	23.53	24.52			
L.S.D. at 5%										
120										
N <sub>0</sub>	9.28	10.45	11.39	10.39	80				100	120
N <sub>40</sub>	10.93	11.42	12.10	11.43	Spray (S)				0.08	0.06
N <sub>60</sub>	11.70	12.38	12.91	12.36	Nitrogen (N)				0.09	0.07
N <sub>80</sub>	12.91	13.24	13.63	13.27	S x N				NS	0.12
Mean	11.20	11.87	12.51							



Table 115. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stem dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	63.17	61.03	59.43	61.21	65.34	62.06	59.55	62.32
N <sub>40</sub>	60.50	58.92	57.69	59.03	62.21	59.43	57.73	59.79
N <sub>60</sub>	58.78	57.12	55.90	57.27	60.16	57.92	56.20	58.09
N <sub>80</sub>	56.52	55.35	54.45	55.44	58.29	56.50	55.02	56.60
Mean	59.74	58.11	56.87		61.50	58.98	57.12	
L.S.D. at 5%								
	120							
N <sub>0</sub>	70.01	67.24	65.31	67.52		80	100	120
N <sub>40</sub>	67.18	65.30	63.68	65.39	Spray (S)	0.14	0.16	0.10
N <sub>60</sub>	65.14	63.39	62.09	63.54	Nitrogen (N)	0.16	0.18	0.12
N <sub>80</sub>	62.58	61.43	60.43	61.48	S x N	NS	NS	0.21
Mean	66.23	64.34	62.88					

Table 116. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on pod dry weight (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)										
	80				100						
	Ethrel concentration (μL/L)										
	0	100	200	Mean	0	100	200	Mean			
N <sub>0</sub>	6.45	7.96	9.17	7.86	14.24	15.97	17.27	15.83			
N <sub>40</sub>	8.35	9.14	10.00	9.16	15.82	16.97	17.98	16.92			
N <sub>60</sub>	8.85	9.98	10.85	9.89	17.52	17.94	18.79	18.08			
N <sub>80</sub>	9.923	10.71	11.35	10.66	17.82	18.67	19.36	18.62			
Mean	8.39	9.45	10.34		16.35	17.39	18.35				
L.S.D. at 5%											
120											
N <sub>0</sub>	20.71	22.30	23.44	22.15	80				100	120	
N <sub>40</sub>	22.19	23.26	24.23	23.23	Spray (S)				0.08	0.06	0.05
N <sub>60</sub>	23.16	24.22	25.00	24.13	Nitrogen (N)				0.09	0.07	0.06
N <sub>80</sub>	24.83	25.33	25.93	25.36	S x N				NS	0.12	0.11
Mean	22.72	23.78	24.65								

Table 117. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf fresh weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	23.84	29.24	34.35	29.14	16.26	21.40	26.12	21.26
N <sub>40</sub>	33.56	38.66	43.94	38.72	24.11	28.94	34.14	29.06
N <sub>60</sub>	37.30	43.02	48.34	42.89	28.10	33.25	33.38	33.24
N <sub>80</sub>	41.34	47.15	52.74	47.08	32.13	36.87	42.16	37.06
Mean	34.01	39.52	44.84		25.15	30.12	35.20	
L.S.D. at 5%								
120								
N <sub>0</sub>	6.96	9.69	12.13	9.59		80	100	120
N <sub>40</sub>	11.05	13.55	16.27	13.62	Spray (S)	0.26	0.24	0.12
N <sub>60</sub>	14.07	16.95	19.76	16.93	Nitrogen (N)	0.30	0.28	0.13
N <sub>80</sub>	17.66	20.16	23.02	20.28	S x N	NS	NS	0.23
Mean	12.43	15.09	17.80					

Table 118. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf turgid weight (g plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	46.46	52.46	57.71	52.21	37.93	43.63	48.66	43.41
N <sub>40</sub>	58.70	63.98	69.30	63.99	48.36	53.55	59.01	53.64
N <sub>60</sub>	62.29	68.47	74.08	68.28	53.41	59.05	64.47	58.98
N <sub>80</sub>	66.66	72.58	78.27	72.50	58.53	63.24	69.13	63.63
Mean	58.53	64.37	69.84		49.56	54.87	60.32	
L.S.D. at 5%								
	120							
N <sub>0</sub>	19.23	23.83	27.77	23.61		80	100	120
N <sub>40</sub>	26.07	29.96	34.28	30.10	Spray (S)	0.41	0.41	0.24
N <sub>60</sub>	31.73	36.24	40.32	36.10	Nitrogen (N)	0.47	0.48	0.27
N <sub>80</sub>	37.55	41.09	45.16	41.27	S x N	NS	NS	0.48
Mean	28.65	32.78	36.88					

Table 119. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on leaf relative water content (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	43.40	47.98	52.02	47.80	34.00	40.56	45.40	39.99
N <sub>40</sub>	49.70	53.16	56.41	53.09	41.71	46.06	50.12	45.96
N <sub>60</sub>	52.71	55.88	58.52	55.70	44.71	48.58	52.02	48.44
N <sub>80</sub>	54.97	58.23	60.46	57.89	47.10	50.69	53.59	50.46
Mean	50.19	53.81	56.85		41.88	46.47	50.28	
L.S.D. at 5%								
	120							
N <sub>0</sub>	26.35	31.16	34.28	30.60		80	100	120
N <sub>40</sub>	32.70	36.17	38.62	35.83	Spray (S)	0.12	0.20	0.12
N <sub>60</sub>	35.14	37.84	40.32	37.77	Nitrogen (N)	0.14	0.23	0.14
N <sub>80</sub>	38.02	40.27	42.43	40.24	S x N	0.24	NS	0.24
Mean	33.05	36.36	38.91					

Table 120. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on rate of photosynthesis ( $\mu\text{mol m}^{-2}\text{s}^{-1}$ ) of mustard (*Brassica juncea* L ) cultivar Alankai grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	13.14	15.60	17.94	15.56	11.62	14.04	16.62	14.09
N <sub>40</sub>	16.19	19.06	22.14	19.13	14.45	17.23	20.38	17.35
N <sub>60</sub>	19.12	22.15	25.07	22.11	17.34	20.29	23.22	20.28
N <sub>80</sub>	21.89	24.71	27.21	24.60	20.19	23.26	26.14	23.20
Mean	17.58	20.38	23.099		15.90	18.71	21.59	
L S D. at 5%								

Table 121. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on stomatal conductance ( $\text{mol m}^{-2}\text{s}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80			100				
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	0.40	0.41	0.43	0.41	0.37	0.39	0.41	0.39
N <sub>40</sub>	0.43	0.45	0.47	0.45	0.41	0.43	0.45	0.43
N <sub>60</sub>	0.47	0.49	0.51	0.49	0.44	0.46	0.49	0.46
N <sub>80</sub>	0.50	0.52	0.54	0.52	0.48	0.51	0.53	0.50
Mean	0.45	0.47	0.49		0.42	0.45	0.47	
L.S.D. at 5%								

Table 122. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on internal carbon dioxide concentration of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	242.67•	259.23	274.13	258.68	228.30	244.93	263.13	245.46
N <sub>40</sub>	275.03	296.93	321.27	297.74	264.73	286.17	307.50	286.13
N <sub>60</sub>	312.33	333.70	354.20	333.41	298.80	319.17	340.93	319.63
N <sub>80</sub>	343.23	360.80	377.33	360.46	332.17	351.23	369.93	351.11
Mean	293.32	312.67	331.73		281.00	300.38	320.38	
L.S.D. at 5%								
				80		100		
Spray (S)				2.52		2.52		
Nitrogen (N)				2.91		2.91		
S x N				5.05		NS		



Table 123. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on transpiration rate ( $\text{kg m}^{-2} \text{ day}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	10.98	11.38	11.72	11.36	8.39	8.90
N <sub>40</sub>	11.59	11.93	12.34	11.95	9.54	9.99
N <sub>60</sub>	11.92	12.39	12.78	12.36	9.877	10.32
N <sub>80</sub>	11.65	12.72	13.07	12.71	10.07	10.51
Mean	11.71	12.10	12.48		9.47	10.38
L.S.D. at 5%						
	80			100		
	Spray (S)			0.03		
	Nitrogen (N)			0.04		
	S x N			NS		
				0.05		

Table 124. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on carboxylation efficiency (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	5.41	6.01	6.54	5.99	5.09	5.73	6.32	5.71
N <sub>40</sub>	5.89	6.42	6.89	6.40	5.46	6.02	6.63	6.04
N <sub>60</sub>	6.12	6.63	7.08	6.61	5.80	6.36	6.81	6.32
N <sub>80</sub>	6.38	6.83	7.21	6.81	6.08	6.62	7.07	6.59
Mean	5.95	6.47	6.93		5.61	6.18	6.71	
L.S.D. at 5%								
<hr/>								
<div>80100</div>								
<hr/>								
Spray (S)0.060.04								
Nitrogen (N)0.070.04								
S x NNS0.07								

Table 125 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on photosynthetic water use efficiency ( $\mu\text{mol mol}^{-1}$ ) of mustard (*Brassica juncea* L ) cultivar Alankai grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80			100				
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	32 85	38 06	41 72	37 54	31 38	35 98	40 54	35 97
N <sub>40</sub>	37 65	42 05	47 11	42 27	35 24	40 07	45 29	40 20
N <sub>60</sub>	40 68	45 51	49 16	45 12	39 41	43 80	47 39	43 53
N <sub>80</sub>	43 78	47 44	50 39	47 20	42 06	45 90	49 32	45 76
Mean	38 74	43 26	47 09		37 02	41 44	45 63	
L S D at 5%								
			80			100		
			Spray (S)			0 34		
			Nitrogen (N)			0 39		
			S x N			0 67		

Table 126. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on plant water use efficiency ( $\text{mg g}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	1.96	2.22	2.45	2.21	2.94	3.17
N <sub>40</sub>	2.38	2.60	2.79	2.59	3.23	3.42
N <sub>60</sub>	2.45	2.65	2.84	2.65	3.37	3.56
N <sub>80</sub>	2.66	2.71	2.90	2.76	3.59	3.76
Mean	2.36	2.54	2.74		3.28	3.67
L.S.D. at 5%						

Table 127. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on l-aminocyclopropane-1-carboxylic acid (ACC) ( $\eta$  mol g fresh weight<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	20.63	23.04	25.81	23.16	23.58	26.06	28.61	26.08
N <sub>40</sub>	24.23	26.74	29.40	26.79	28.04	30.63	33.03	30.57
N <sub>60</sub>	27.76	30.80	34.26	30.94	31.60	34.90	37.30	34.60
N <sub>80</sub>	32.80	35.76	38.56	35.70	36.43	39.12	41.98	39.18
Mean	26.36	29.08	32.01		29.91	32.68	35.23	
L.S.D. at 5%								
80100								
Spray (S)0.250.18								
Nitrogen (N)0.290.20								
S x N0.510.35								

Table 128. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on ACC oxidase ( $\eta$  l g<sup>-1</sup> fresh weight h<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	71.40	78.29	83.80	77.84	82.56	85.22	97.49	88.42
N <sub>40</sub>	94.01	100.68	109.39	101.36	106.65	109.13	121.26	112.35
N <sub>60</sub>	109.35	115.35	122.63	115.78	116.94	125.12	134.72	125.60
N <sub>80</sub>	119.69	129.46	137.08	128.83	131.26	142.18	154.71	142.72
Mean	98.68	105.95	113.23		109.35	115.41	127.05	
L.S.D. at 5%								
80100								
Spray (S)1.291.89								
Nitrogen (N)1.492.19								
S x NNS3.79								

Table 129. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on ethylene evolution ( $\eta\text{l h}^{-1} \text{ plant}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	11.65	13.08	13.93	12.89	14.58	16.18	17.37	16.05
N <sub>40</sub>	18.83	20.05	22.58	20.49	24.97	24.35	26.45	25.26
N <sub>60</sub>	25.05	27.45	29.81	27.43	28.89	31.61	34.06	31.52
N <sub>80</sub>	32.40	35.41	37.65	35.15	37.27	39.06	40.76	39.03
Mean	21.98	24.00	25.99		26.43	27.80	29.66	
L.S.D. at 5%								

Table 130. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrate reductase activity ( $\mu\text{ mol NO}_2^- \text{ g}^{-1} \text{ h}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80 and 100 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)					
	80			100		
	Ethrel concentration (μL/L)					
	0	100	200	Mean	0	Mean
N <sub>0</sub>	0.15	0.17	0.20	0.17	0.12	0.13
N <sub>40</sub>	0.40	0.40	0.43	0.40	0.32	0.35
N <sub>60</sub>	0.46	0.50	0.54	0.50	0.40	0.43
N <sub>80</sub>	0.58	0.62	0.65	0.62	0.52	0.55
Mean	0.39	0.42	0.45		0.33	0.39
L.S.D. at 5%						
			80	100		
Spray (S)			0.01	0.01		
Nitrogen (N)			0.01	0.01		
S x N			NS	NS	NS	



Table 131 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrogen content (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	1.93	2.15	2.31	2.13	1.57	1.95	2.22	1.91
N <sub>40</sub>	2.27	2.42	2.57	2.42	1.99	2.26	2.49	2.25
N <sub>60</sub>	2.46	2.61	2.75	2.61	2.12	2.36	2.57	2.35
N <sub>80</sub>	2.74	2.82	2.87	2.81	2.23	2.47	2.68	2.46
Mean	2.35	2.50	2.62		1.98	2.26	2.49	
L.S.D. at 5%								
120								
N <sub>0</sub>	1.55	1.76	1.93	1.76		80	100	120
N <sub>40</sub>	1.72	1.86	2.01	1.86	Spray (S)	0.012	0.008	0.012
N <sub>60</sub>	1.81	1.97	2.12	1.97	Nitrogen (N)	0.013	0.009	0.014
N <sub>80</sub>	1.99	2.12	2.22	2.11	S x N	0.023	NS	NS
Mean	1.77	1.93	2.07					

Table 132. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on nitrogen accumulation (mg plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen at 80, 100 and 120 DAS

Nitrogen (kg/ha)	Sampling days (after sowing)										
	80				100						
	Ethrel concentration (μL/L)										
	0	100	200	Mean	0	100	200	Mean			
N <sub>0</sub>	415.67	541.67	661.67	539.67	387.00	552.33	707.33	548.89			
N <sub>40</sub>	627.33	750.67	885.00	754.33	612.00	771.67	937.00	773.56			
N <sub>60</sub>	719.00	857.33	996.67	857.67	704.33	872.67	1038.33	871.78			
N <sub>80</sub>	850.33	972.33	1091.33	971.33	817.33	977.00	1145.67	980.00			
Mean	653.08	780.50	909.67		630.17	793.42	957.08				
L.S.D. at 5%											
120											
N <sub>0</sub>	429.33	554.00	672.00	551.89	80				100	120	
N <sub>40</sub>	585.00	693.00	819.67	699.22	Spray (S)				5.61	4.48	6.21
N <sub>60</sub>	696.33	829.33	962.33	829.33	Nitrogen (N)				6.48	5.17	7.17
N <sub>80</sub>	841.67	967.67	1094.00	967.78	S x N				11.23	NS	12.42
Mean	638.08	761.08	887.00								

Table 133. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on number of pods plant<sup>-1</sup>, number of seeds pod<sup>-1</sup> and 1000 seed weight (g) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Pods plant <sup>-1</sup>				Seeds pod <sup>-1</sup>			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	122.00	129.33	137.00	129.44	11.14	11.87	12.50	11.84
N <sub>40</sub>	137.00	144.33	152.00	144.44	12.13	12.69	13.31	12.71
N <sub>60</sub>	149.33	156.67	164.00	156.67	13.30	13.80	14.35	13.82
N <sub>80</sub>	159.67	166.67	174.67	167.00	14.00	14.87	15.44	14.77
Mean	142.00	149.25	156.92		12.64	13.31	13.90	
L.S.D. at 5%								
1000 seed weight								
N <sub>0</sub>	4.13	4.32	4.45	4.30	Pods plant <sup>-1</sup>		Seeds pod <sup>-1</sup>	1000 seed weight
N <sub>40</sub>	4.37	4.48	4.55	4.47				
N <sub>60</sub>	4.49	4.57	4.61	4.56	Spray (S)		0.54	0.04
N <sub>80</sub>	4.49	4.66	4.66	4.60	Nitrogen (N)		0.63	0.05
Mean	4.37	4.51	4.57		S x N		NS	0.08
								NS

Table 134 Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on seed yield (q ha<sup>-1</sup>), biological yield (q ha<sup>-1</sup>) and harvest index (%) of mustard (*Brassica juncea* L ) cultivar Alankai grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Seed yield				Biological yield			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	7.01	8.29	9.53	8.28	35.47	39.30	42.80	39.19
N <sub>40</sub>	90.08	10.26	11.51	10.28	43.28	46.51	50.27	46.69
N <sub>60</sub>	11.15	12.36	13.56	12.36	48.84	52.64	56.10	52.53
N <sub>80</sub>	12.55	14.43	15.71	14.23	53.60	57.05	60.70	57.12
Mean	9.95	11.34	12.58		45.30	48.88	52.47	
L S.D. at 5%								
Harvest index								
N <sub>0</sub>	19.76	21.10	22.27	21.04	Seed yield		Biological	Harvest
N <sub>40</sub>	20.98	22.05	22.90	21.98			yield	index
N <sub>60</sub>	22.83	23.48	24.17	23.49	Spray (S)	0.29	0.20	0.08
N <sub>80</sub>	23.41	25.29	25.88	24.86	Nitrogen (N)	0.34	0.23	0.09
Mean	21.74	22.98	23.80		S x N	NS	NS	0.15

Table 135. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on seed nitrogen content (mg g<sup>-1</sup> plant<sup>-1</sup>), nitrogen harvest index and nitrogen yield merit of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Seed nitrogen content				Nitrogen harvest index			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	24.73	31.27	37.07	31.02	0.06	0.06	0.06	0.06
N <sub>40</sub>	33.37	38.77	45.23	39.12	0.05	0.05	0.05	0.05
N <sub>60</sub>	37.27	42.83	49.20	43.10	0.05	0.05	0.05	0.05
N <sub>80</sub>	42.00	47.70	53.80	47.83	0.05	0.05	0.05	0.05
Mean	34.34	40.14	45.33		0.05	0.05	0.05	
L.S.D., at 5%								
Nitrogen yield merit								
N <sub>0</sub>	1.48	1.88	2.22	1.86	Seed nitrogen content	Nitrogen harvest index	Nitrogen yield merit	
N <sub>40</sub>	1.67	1.94	2.26	1.96				
N <sub>60</sub>	1.86	2.14	2.46	2.15	Spray (S)	NS	0.01	
N <sub>80</sub>	2.10	2.38	2.69	2.39	Nitrogen (N)	NS	0.01	
Mean	1.78	2.08	2.41		S x N	NS	0.03	

Table 136. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on oil yield ( $q\ ha^{-1}$ ) and oil content (%) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Oil yield				Oil content			
	Ethrel concentration ( $\mu L/L$ )							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	2.34	2.87	3.42	2.88	33.39	34.65	35.92	34.65
N <sub>40</sub>	2.96	3.48	4.04	3.49	32.64	33.90	35.08	33.88
N <sub>60</sub>	3.51	4.05	4.61	4.06	31.52	32.80	34.03	32.78
N <sub>80</sub>	3.84	4.60	5.20	4.55	30.61	31.91	33.11	31.87
Mean	3.16	3.75	4.32		32.04	33.31	34.53	
L.S.D. at 5%								
		Oil yield		Oil content				
Spray (S)		0.02		0.07				
Nitrogen (N)		0.03		NS				
S x N		NS		NS				

Table 137. Effect of ethrel spray at 60d after sowing (DAS, flowering stage) on acid value, iodine value and saponification value of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Acid value				Iodine value			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	2.69	3.01	3.32	3.01	97.13	99.14	100.95	99.08
N <sub>40</sub>	2.56	2.89	3.39	2.95	95.93	98.06	99.47	97.82
N <sub>60</sub>	2.44	2.78	3.06	2.76	94.18	96.20	97.97	96.11
N <sub>80</sub>	2.29	2.62	2.89	2.60	92.70	94.73	95.96	94.53
Mean	2.50	2.82	3.17		95.04	97.03	98.59	
L.S.D. at 5%								
Saponification value								
N <sub>0</sub>	138.96	141.96	145.17	142.03	Acid value	Iodine value	Saponifica-	
N <sub>40</sub>	144.97	147.82	150.52	147.77			tion value	
N <sub>60</sub>	148.58	151.33	154.58	151.49	Spray (S)	0.10	0.09	0.30
N <sub>80</sub>	151.81	154.72	153.61	154.71	Nitrogen (N)	NS	NS	NS
Mean	146.08	148.96	151.97		S x N	NS	NS	NS

Table 138 Pooled analysis of plant dry weight in Experiment 3 and 4

Sampling stage (days after sowing)																
80100120																
Nitrogen	Ethrel concentration (μL/L)															
	0100200Mean0100200Mean															
	IrrigatedNon-IrrigatedIrrigatedNon-Irrigated															
N <sub>0</sub>	25 35	29 18	32 59	29 04	28 45	32 34	35 68	32 16	32 84	36 39	39 81	36 35				
N <sub>40</sub>	33 68	37 22	40 92	37 27	36 78	40 33	43 99	40 37	40 84	44 39	48 09	44 44				
N <sub>60</sub>	35 81	39 90	43 97	39 89	39 77	43 83	47 87	43 71	44 73	48 76	52 77	48 75				
N <sub>80</sub>	39 67	43 53	47 03	43 41	44 77	48 63	52 12	48 51	50 82	54 65	58 45	54 64				
N <sub>0</sub>	21 56	25 25	28 69	24 95	24 64	28 38	31 79	28 27	27 71	31 44	34 84	31 33				
N <sub>40</sub>	27 65	31 06	34 41	31 04	30 78	34 10	37 66	34 18	34 03	37 21	40 81	37 35				
N <sub>60</sub>	29 16	32 82	36 26	32 75	33 28	36 93	40 39	36 87	38 47	42 11	45 48	42 02				
N <sub>80</sub>	31 03	34 49	37 96	34 49	36 13	39 56	43 02	39 57	42 28	45 64	49 16	45 69				
Mean	30 49	34 18	37 73	34 49	34 32	38 01	41 56		38 96	42 57	46 18					
L S D at 5%																
80100120																
Irrigation (A)NS																











Table 143. Pooled analysis of carboxylation efficiency in Experiment 3 and 4

Nitrogen	Sampling stage (days after sowing)							
	80				100			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
		Irrigated				Irrigated		
N <sub>0</sub>	5.85	6.41	6.91	6.39	5.45	6.05	6.59	6.03
N <sub>40</sub>	6.25	6.74	7.22	6.74	5.93	6.48	6.99	6.47
N <sub>60</sub>	6.48	6.92	7.30	6.90	6.23	6.72	7.19	6.71
N <sub>80</sub>	6.58	6.98	7.36	6.97	6.47	6.88	7.27	6.87
		Non-Irrigated				Non-Irrigated		
N <sub>0</sub>	5.41	6.01	6.54	5.99	5.09	5.73	6.32	5.71
N <sub>40</sub>	5.89	6.42	6.89	5.40	5.46	6.02	6.63	6.04
N <sub>60</sub>	6.12	6.63	7.08	6.61	5.80	6.36	6.81	6.32
N <sub>80</sub>	6.38	6.83	7.21	6.81	6.08	6.62	7.70	6.59
Mean	6.12	6.62	7.06		5.81	6.36	6.86	
L.S.D. at 5%								
							80	100
						Irrigation (A)	NS	NS
						Nitrogen (B)	0.04	0.04
						A X B	NS	NS
						Spray (C)	NS	NS
						A X C	NS	NS
						B X C	NS	NS
						A X B X C	NS	NS

Table 144 Pooled analysis of photosynthetic water use efficiency in Experiment 3 and 4

Nitrogen	Sampling stage (days after sowing)						
	80				100		
	Ethrel concentration (μL/L)						Mean
	0	100	200	Mean	0	100	
N <sub>0</sub>	35.95	39.69	43.19	38.61	32.85	37.69	37.42
N <sub>40</sub>	40.35	44.59	49.12	44.69	37.58	42.82	42.45
N <sub>60</sub>	43.75	47.54	51.23	47.50	41.80	46.03	44.62
N <sub>80</sub>	45.37	48.86	51.49	48.57	44.30	47.59	47.56
	Irrigated				Irrigated		
	Non-Irrigated				Non-Irrigated		
N <sub>0</sub>	32.85	38.06	41.72	37.54	31.38	35.98	35.97
N <sub>40</sub>	37.65	42.05	47.11	42.27	35.24	40.07	40.20
N <sub>60</sub>	40.68	45.51	49.16	45.12	39.41	43.80	43.53
N <sub>80</sub>	43.78	47.44	50.39	47.20	42.06	45.90	45.76
Mean	40.05	44.22			38.08	42.48	46.44
L.S.D. at 5%							
	80				100		
	Irrigation (A)				NS		
	Nitrogen (B)				0.34		
	A X B				NS		
	Spray (C)				0.0002		
	A X C				NS		
	B X C				0.006		
	A X B X C				NS		



Table 146 Pooled analysis of pod number, seed yield and oil yield in Experiment 3 and 4

Nitrogen	Pod number				Seed yield				Oil yield			
					Ethrel concentration ( $\mu\text{I/L}$ )							
	0	100	200	Mean	0	100	200	Mean	0	100	200	Mean
N <sub>0</sub>	138 00	146 33	153 33	145 89	9 19	10 67	12 05	10 64	3 27	3 94	4 60	3 94
N <sub>40</sub>	159 30	166 33	173 00	166 22	12 27	13 65	15 00	13 64	4 29	4 95	5 63	4 96
N <sub>60</sub>	172 67	181 67	189 00	181 44	15 03	16 58	18 07	16 56	5 12	5 85	6 61	5 86
N <sub>80</sub>	190 33	197 33	203 67	197 11	18 29	19 74	21 20	19 74	6 02	6 78	7 55	6 78
	Non-Irrigated				Non-Irrigated				Non-Irrigated			
N <sub>0</sub>	122 00	129 33	137 00	129 44	7 01	8 29	9 53	8 28	2 34	2 87	3 42	2 88
N <sub>40</sub>	137 00	144 33	152 00	144 44	9 08	10 26	11 51	10 28	2 96	3 48	4 04	3 49
N <sub>60</sub>	149 33	156 67	164 00	156 67	11 15	12 36	13 56	12 36	3 51	4 05	4 61	4 06
N <sub>80</sub>	159 67	166 67	174 67	167 00	12 55	14 43	15 71	14 23	3 84	4 60	5 20	4 55
Mean	153 54	161 08	168 33		11 82	13 25	14 58		3 92	4 57	5 21	
L S D at 5%												
					Pod number		Seed yield		Oil yield			
Irrigation (A)					NS		NS		NS		NS	
Nitrogen (B)					0 44		0 19		0 02		0 02	
A X B					NS		NS		NS		NS	
Spray (C)					0 005		0 0001		NS		NS	
A X C					NS		NS		NS		NS	
B X C					NS		NS		NS		NS	
A X B X C					NS		NS		NS		NS	



Table 147. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on leaf area plant<sup>-1</sup> (cm<sup>2</sup>) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80, 100 and 120 DAS

Treatment	Sampling stage (days after sowing)				
	80				
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	3473.67	3512.33	3493.00	4041.67	4150.33
E <sub>200</sub>	3748.33	3850.67	3795.50	4251.67	4353.33
S <sub>1</sub>	3383.33	3406.67	3395.00	3843.33	3973.00
Mean	3535.11	3589.67		4045.56	4158.89

Treatment	120				
	L.S.D. at 5%				
	Non-irrigated	Irrigated	Mean	Irrigation (I)	Spray (S)
E <sub>0</sub>	2667.67	2676.67	2672.17	NS	NS
E <sub>200</sub>	2840.00	2951.67	2895.84	NS	NS
S <sub>1</sub>	2504.00	2654.33	2579.17	NS	NS
Mean	2670.56	2760.89		NS	NS

Table 148. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on plant dry weight (g/plant) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80, 100 and 120 DAS

Treatment	Sampling stage (days after sowing)					
	80			100		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated	Mean
E <sub>0</sub>	35.20	42.33	38.76	40.20	47.84	44.02
E <sub>200</sub>	36.10	43.22	39.66	41.20	49.08	45.14
S <sub>1</sub>	33.49	41.17	37.33	38.25	46.08	42.17
Mean	34.93	42.24		39.88	47.67	

Treatment	L.S.D. at 5%					
	120			80		
	Non-irrigated	Irrigated	Mean	Irrigation (I)	Spray (S)	I x S
E <sub>0</sub>	45.26	53.79	49.53	NS	0.29	NS
E <sub>200</sub>	46.39	55.32	50.86	NS	0.20	NS
S <sub>1</sub>	42.70	51.58	47.14	NS	0.25	NS
Mean	44.79	53.56		NS	NS	NS

Table 149. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on rate of photosynthesis ( $\mu$  mol m<sup>-2</sup> s<sup>-1</sup>) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Non-irrigated	Irrigated	Mean
E <sub>0</sub>	24.03	25.32	22.40	24.30	23.35
E <sub>200</sub>	25.32	26.55	24.21	25.51	24.86
S <sub>1</sub>	23.20	24.77	21.63	23.59	22.61
Mean	24.19	25.55	22.75	24.47	

L.S.D. at 5%

	80	100
Irrigation (I)	NS	NS
Spray (S)	0.22	0.20
I x S	NS	NS

Table 150 Effect of ethiel (E, 200 $\mu$ L/L) or silver thiosulphate (S, 1mM) at 60d after sowing (DAS, flowering stage) on stomatal conductance (mol m<sup>-2</sup> s<sup>-1</sup>) of mustard (*Brassica juncea* L ) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)			
	80		100	
	Non-irrigated	Irrigated	Non-irrigated	Irrigated
E <sub>0</sub>	0 51	0 52	0 49	0 51
E <sub>200</sub>	0 53	0 54	0 52	0 53
S <sub>1</sub>	0 50	0 51	0 48	0 50
Mean	0 51	0 52	0 50	0 51

L S D at 5%			
		80	100
Irrigation (I)		NS	NS
Spray (S)		0 007	0 008
I x S		NS	NS

Table 151. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on internal CO<sub>2</sub> concentration ( $\mu$  mol mol<sup>-1</sup>) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Non-irrigated	Irrigated	Mean
E <sub>0</sub>	354.60	361.60	342.77	354.37	348.57
E <sub>200</sub>	368.10	376.07	361.07	368.67	364.87
S <sub>1</sub>	346.67	355.83	335.23	345.10	340.17
Mean	356.46	364.50	346.36	356.04	

L.S.D. at 5%

	80	100
Irrigation (I)	NS	NS
Spray (S)	13.74	13.57
I x S	NS	NS

Table 152. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on transpiration rate (kg m<sup>-2</sup> day<sup>-1</sup>) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Non-irrigated	Irrigated	Mean
E <sub>0</sub>	12.72	14.11	10.52	11.69	11.10
E <sub>200</sub>	12.85	14.25	10.63	11.74	11.19
S <sub>1</sub>	12.57	13.86	10.42	11.54	10.98
Mean	12.71	14.07	10.52	11.66	

L.S.D. at 5%				
		80	100	
Irrigation (I)		NS	NS	NS
Spray (S)		0.05	0.02	
I x S		NS	NS	NS

Table 153. Effect of ethrel (E; 200µL/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on carboxylation efficiency (%) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	6.78	7.00	6.89	6.54	6.86
E <sub>200</sub>	6.88	7.06	6.97	6.70	6.92
S <sub>1</sub>	6.69	6.96	6.83	6.45	6.83
Mean	6.78	7.01		6.56	6.87

L.S.D. at 5%

	80	100
Irrigation (I)	NS	NS
Spray (S)	0.03	0.04
I x S	NS	NS

Table 154. Effect of ethrel (E; 200µL/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on photosynthetic water use efficiency ( $\mu\text{ mol mol}^{-1}$ ) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		Mean
	Non-irrigated	Irrigated	Non-irrigated	Irrigated	
E <sub>0</sub>	46.82	48.70	45.41	47.34	46.38
E <sub>200</sub>	47.78	49.38	46.86	48.44	47.65
S <sub>1</sub>	46.09	48.25	44.76	47.18	45.97
Mean	46.90	48.78	45.67	47.65	

L.S.D. at 5%				
		80	100	
Irrigation (I)		NS	NS	NS
Spray (S)		0.42	0.42	
I x S		NS	NS	NS



Table 155. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on plant water use efficiency (mg g<sup>-1</sup>) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)			
	80		100	
	Non-irrigated	Irrigated	Non-irrigated	Irrigated
E <sub>0</sub>	2.76	3.00	3.82	4.09
E <sub>200</sub>	2.81	3.04	3.88	4.18
S <sub>1</sub>	2.68	2.97	3.67	3.99
Mean	2.75	3.00	3.79	4.09

L.S.D. at 5%

	80	100
Irrigation (I)	NS	NS
Spray (S)	0.02	0.02
I x S	NS	NS

Table 156. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on 1-aminocyclopropane-1-carboxylic acid content (ACC) ( $\eta$  mol g<sup>-1</sup> fresh weight) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	30.86	25.88	28.37	36.17	30.30
E <sub>200</sub>	35.21	29.79	32.50	40.16	35.52
S <sub>1</sub>	26.46	22.09	24.28	31.43	26.33
Mean	30.84	25.92		35.92	30.72

L.S.D. at 5%

	80	100
Irrigation (I)	NS	NS
Spray (S)	4.03	4.36
I x S	NS	NS

Table 157. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on ACC oxidase ( $\eta$  L g<sup>-1</sup> fresh weight h<sup>-1</sup>) of mustard (*Brassica juncea* L ) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	124.00	108.13	116.07	135.49	125.71
E <sub>200</sub>	132.97	117.50	125.23	149.53	134.88
S <sub>1</sub>	109.09	91.93	100.51	121.88	111.15
Mean	122.02	105.85		135.63	123.91

L.S.D. at 5%

	80	100
Irrigation (I)	NS	NS
Spray (S)	8.25	10.93
I x S	NS	NS

Table 158. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S, 1mM) at 60d after sowing (DAS, flowering stage) on ethylene evolution ( $\eta$ l h<sup>-1</sup> plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	31.88	27.66	29.77	35.88	32.14
E <sub>200</sub>	36.92	31.17	34.05	39.99	35.55
S <sub>1</sub>	25.64	21.92	23.78	30.29	26.07
Mean	31.48	26.92		35.39	31.25

L.S.D. at 5%				
		80	100	
Irrigation (I)		NS	NS	NS
Spray (S)		2.97	2.83	
I x S		NS	NS	NS

Table 159 Effect of ethrel (E, 200 $\mu$ L/L) or silver thiosulphate (S, 1mM) at 60d after sowing (DAS, flowering stage) on nitrate reductase activity ( $\mu$  mol NO<sub>2</sub><sup>-</sup> g<sup>-1</sup> h<sup>-1</sup>) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80 and 100 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	0.59	0.68	0.63	0.52	0.59
E <sub>200</sub>	0.64	0.73	0.69	0.57	0.65
S <sub>1</sub>	0.56	0.64	0.60	0.49	0.56
Mean	0.60	0.68		0.53	0.60

L S D at 5%				
		80	100	
Irrigation (I)		NS	NS	NS
Spray (S)		0.013	0.014	
I x S		NS	NS	NS

Table 160 Effect of ethrel (E, 200µL/L) or silver thiosulphate (S, 1mM) at 60d after sowing (DAS, flowering stage) on N content (%) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80, 100 and 120 DAS

Treatment	Sampling stage (days after sowing)				
	80		100		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	2.81	2.92	2.87	2.46	2.63
E <sub>200</sub>	2.84	3.01	2.93	2.52	2.69
S <sub>1</sub>	2.73	2.89	2.81	2.42	2.57
Mean	2.79	2.94		2.47	2.63

Treatment	L.S.D. at 5%				
	120		80		
			Irrigation (I)	Spray (S)	I x S
E <sub>0</sub>	2.11	2.28	2.20	NS	NS
E <sub>200</sub>	2.17	2.31	2.24	NS	NS
S <sub>1</sub>	2.05	2.26	2.16	0.02	0.01
Mean	2.11	2.28		NS	NS

Table 161. Effect of ethrel (E; 200µL/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on N accumulation (mg plant<sup>-1</sup>) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions at 80, 100 and 120 DAS

Treatment	Sampling stage (days after sowing)				
	80				
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	988.00	1237.00	1112.50	987.67	1258.00
E <sub>200</sub>	1025.33	1300.67	1163.00	1038.33	1318.67
S <sub>1</sub>	913.00	1189.67	1051.33	925.67	1184.33
Mean	975.44	1242.44		983.89	1253.67

Treatment	L.S.D at 5%				
	120				
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated
E <sub>0</sub>	956.67	1228.67	1092.67	80	100
E <sub>200</sub>	1008.33	1277.67	1143.00	Irrigation (I)	NS
S <sub>1</sub>	875.33	1167.33	1021.33	Spray (S)	8.51
Mean	946.78	1224.56		I x S	NS

Table 162 Effect of ethrel (E, 200µL/L) or silver thiosulphate (S, 1mM) at 60d after sowing (DAS, flowering stage) on number of pods plant<sup>-1</sup>, number of seeds pod<sup>-1</sup> and 1000 seed weight (g) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions

Treatment	Seeds pod <sup>-1</sup>			
	Pods plant <sup>-1</sup>		Seeds pod <sup>-1</sup>	
	Non-irrigated	Irrigated	Mean	Mean
E <sub>0</sub>	165.00	195.00	180.00	15.86
E <sub>200</sub>	169.67	198.67	184.17	16.03
S <sub>1</sub>	160.00	189.67	174.83	15.31
Mean	164.89	194.44	14.64	16.83

	L.S.D at 5%			
	1000 seed weight		Seeds pod <sup>-1</sup>	
	Non-irrigated	Irrigated	Mean	Mean
E <sub>0</sub>	4.65	4.68	4.67	1000 seed weight
E <sub>200</sub>	4.68	4.74	4.71	NS
S <sub>1</sub>	4.63	4.65	4.64	NS
Mean	4.65	4.69	4.67	NS



Table 163 Effect of ethrel (E, 200µL/L) or silver thiosulphate (S, 1mM) at 60d after sowing (DAS, flowering stage) on seed yield (q ha<sup>-1</sup>), biological yield (q ha<sup>-1</sup>) and harvest index (%) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions

Treatment	Seed yield			Biological yield		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated	Mean
E <sub>0</sub>	14 19	19 32	16 75	56 58	67 36	61 97
E <sub>200</sub>	14 81	20 17	17 49	57 99	69 15	63 57
S <sub>1</sub>	13 16	18 07	15 61	53 37	64 48	58 93
Mean	14 05	19 18		55 98	67 00	

L S D at 5%					
Harvest index			Seed yield		
E <sub>0</sub>	25 07	28 68	26 88		
E <sub>200</sub>	25 53	29 17	27 35		
S <sub>1</sub>	24.65	28 02	26 33		
Mean	25 09	28 62			

Irrigation (I)			NS	NS	NS
Spray (S)			0 12	0 28	0 10
I x S			NS	NS	NS

Table 164. Effect of ethrel (E; 200 $\mu$ L/L) or silver thiosulphate (S, 1mM) at 60d after sowing (DAS, flowering stage) on seed nitrogen content (mg g<sup>-1</sup> seed<sup>-1</sup>), nitrogen harvest index and nitrogen yield potential of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions

. Treatment	Seed nitrogen content			Nitrogen harvest index		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated	Mean
E <sub>0</sub>	47.20	56.60	51.90	0.05	0.05	0.05
E <sub>200</sub>	49.07	59.17	54.12	0.05	0.05	0.05
S <sub>1</sub>	45.17	55.17	50.17	0.05	0.05	0.05
Mean	47.14	56.98		0.05	0.05	

L.S.D. at 5%						
Nitrogen yield potential						
E <sub>0</sub>	2.36	2.83	2.60	Seed		
				nitrogen	Nitrogen	Nitrogen
E <sub>200</sub>	2.45	2.96	2.70	content	harvest	yield
S <sub>1</sub>	2.26	2.76	2.51	potential	index	potential
Mean	2.36	2.85		NS	NS	NS
				NS	NS	NS
				0.39	NS	0.02
				NS	NS	NS
				I x S		

Table 165. Effect of ethrel (E; 200µL/L) or silver thiosulphate (S; 1mM) at 60d after sowing (DAS, flowering stage) on oil yield (q ha<sup>-1</sup>) and oil content (%) of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions

Treatment	Oil yield		Oil content	
	Non-irrigated	Irrigated	Mean	Mean
E <sub>0</sub>	4.46	6.55	5.50	31.46
E <sub>200</sub>	4.74	6.96	5.85	33.89
S <sub>1</sub>	3.97	5.82	4.89	34.53
Mean	4.39	6.44	31.23	32.67

L.S.D. at 5%

	Oil yield	Oil content
Irrigation (I)	NS	NS
Spray (S)	0.05	0.17
I x S	NS	NS

Table 166 Effect of ethiel (E, 200 $\mu$ L/L) or silver thiosulphate (S, 1mM) at 60d after sowing (DAS, flowering stage) on acid value, iodine value and saponification value of mustard (*Brassica juncea* L.) grown with basal 80kg N/ha under irrigated and non-irrigated conditions

Treatment	Acid value			Iodine value		
	Non-irrigated	Irrigated	Mean	Non-irrigated	Irrigated	Mean
E <sub>0</sub>	2 52	3 00	2 76	93 86	96 72	95 29
E <sub>200</sub>	2 69	3 24	2 97	94 89	97 97	96 43
S <sub>1</sub>	2 14	2 66	2 40	92 63	95 34	93 98
Mean	2 45	2 97		93 79	96 68	

	Saponification value		L S D at 5%		
	Irrigation (I)	Acid value	Iodine value	Saponification value	
E <sub>0</sub>	153 45	158 44	155 95		
E <sub>200</sub>	156 06	160 53	158 30		
S <sub>1</sub>	151 54	155 23	153 39		
Mean	153 68	158 07			

Table 167. Linear regression values of plant characteristics in Experiment 1 at 80 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.994**	1.000							
Specific leaf weight	0.968**	0.947**	1.000						
Specific leaf area	0.973**	0.951**	0.996**	1.000					
Leaf dry weight	0.997**	0.996**	0.951**	0.955**	1.000				
Plant N content	0.865**	0.860**	0.897**	0.866**	0.866**	1.000			
Pod number	0.973**	0.960**	0.976**	0.972**	0.965**	0.908**	1.000		
Seed number	0.900**	0.877**	0.851**	0.850**	0.912**	0.837**	0.893**	1.000	
Seed yield	0.965**	0.946**	0.935**	0.936**	0.967**	0.876**	0.969**	0.974**	1.000

Table 168. Linear regression values of plant characteristics in Experiment 1 at 100 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Paramcter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.997**	1.000							
Specific leaf weight	0.943**	0.932**	1.000						
Specific leaf area	0.954**	0.945**	0.998**	1.000					
Leaf dry weight	0.996**	0.995**	0.911**	0.924**	1.000				
Plant N content	0.848**	0.848**	0.866**	0.877**	0.831**	1.000			
Pod number	0.961**	0.964**	0.953**	0.965**	0.944**	0.919**	1.000		
Seed number	0.969**	0.972**	0.946**	0.962**	0.986**	0.795**	0.893**	1.000	
Seed yield	0.995**	0.998**	0.923**	0.938**	0.994**	0.861**	0.969**	0.974**	1.000

Table 169. Linear regression values of plant characteristics in Experiment 1 at 120 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.998**	1.000							
Specific leaf weight	0.914**	0.900**	1.000						
Specific leaf area	0.895**	0.878**	0.992**	1.000					
Leaf dry weight	0.996**	0.997**	0.880**	0.858**	1.000				
Plant N content	0.828**	0.817**	0.941**	0.952**	0.791**	1.000			
Pod number	0.940**	0.937**	0.961**	0.956**	0.917**	0.953**	1.000		
Seed number	0.983**	0.990**	0.843**	0.813**	0.993**	0.758*	0.893**	1.000	
Seed yield	0.992**	0.993**	0.929**	0.910**	0.985**	0.869**	0.969**	0.974**	1.000

Table 170. Linear regression values of plant characteristics in Experiment 2 at 80 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.995**	1.000							
Specific leaf weight	0.929**	0.902**	1.000						
Specific leaf area	0.933**	0.904**	0.999**	1.000					
Leaf dry weight	0.994**	0.998**	0.889**	0.893**	1.000				
Plant N content	0.678*	0.665*	0.770**	0.771**	0.649*	1.000			
Pod number	0.996**	0.991**	0.910**	0.916**	0.995**	0.677*	1.000		
Seed number	0.923**	0.916**	0.771**	0.780**	0.939**	0.510 <sup>NS</sup>	0.940**	1.000	
Seed yield	0.991**	0.985**	0.891**	0.898**	0.992**	0.654*	0.996**	0.960**	1.000



Table 171. Linear regression values of plant characteristics in Experiment 2 at 100 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.996**	1.000							
Specific leaf weight	0.890**	0.861**	1.000						
Specific leaf area	0.865**	0.832**	0.995**	1.000					
Leaf dry weight	0.996**	0.998**	0.849**	0.820**	1.000				
Plant N content	0.839**	0.821**	0.929**	0.915**	0.808**	1.000			
Pod number	0.994**	0.985**	0.923**	0.899**	0.984**	0.863**	1.000		
Seed number	0.956**	0.946**	0.789**	0.762*	0.961**	0.738*	0.940**	1.000	
Seed yield	0.995**	0.985**	0.909**	0.883**	0.986**	0.848**	0.996**	0.960**	1.000

Table 172 Linear regression values of plant characteristics in Experiment 2 at 120 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1 000								
Plant dry weight	0.995**	1.000							
Specific leaf weight	0.884**	0.851**	1.000						
Specific leaf area	0.896**	0.864**	0.996**	1.000					
Leaf dry weight	0.996**	0.998**	0.844**	0.857**	1 000				
Plant N content	0.944**	0.940**	0.908**	0.912**	0.935**	1 000			
Pod number	0.990**	0.978**	0.923**	0.936**	0.977**	0.960**	1 000		
Seed number	0.968**	0.966**	0.782**	0.801**	0.974**	0.867**	0.940**	1.000	
Seed yield	0.996**	0.986**	0.907**	0.920**	0.986**	0.947**	0.996**	0.960**	1.000

Table 173. Linear regression values of plant characteristics in Experiment 3 at 80. DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.870**	1.000							
Specific leaf weight	0.889**	0.992**	1.000						
Specific leaf area	0.902**	0.992**	0.998**	1.000					
Leaf dry weight	0.916**	0.985**	0.981**	0.988**	1.000				
Plant N content	0.942**	0.973**	0.977**	0.986**	0.995**	1.000			
Pod number	0.927**	0.962**	0.964**	0.973**	0.989**	0.992**	1.000		
Seed number	0.932**	0.935**	0.941**	0.954**	0.975**	0.983**	0.996**	1.000	
Seed yield	0.934**	0.929**	0.933**	0.948**	0.974**	0.981**	0.993**	0.999**	1.000

Table 174. Linear regression values of plant characteristics in Experiment 3 at 100 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.983**	1.000							
Specific leaf weight	0.990**	0.990**	1.000						
Specific leaf area	0.994**	0.988**	0.997**	1.000					
Leaf dry weight	0.999**	0.986**	0.991**	0.994**	1.000				
Plant N content	0.995**	0.973**	0.985**	0.993**	0.992**	1.000			
Pod number	0.997**	0.986**	0.989**	0.991**	0.998**	0.989**	1.000		
Seed number	0.994**	0.968**	0.978**	0.983**	0.993**	0.989**	0.996**	1.000	
Seed yield	0.994**	0.964**	0.975**	0.983**	0.991**	0.991**	0.993**	0.999**	1.000

Table 175. Linear regression values of plant characteristics in Experiment 3 at 120 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.990**	1.000							
Specific leaf weight	0.971**	0.982**	1.000						
Specific leaf area	0.981**	0.987**	0.998**	1.000					
Leaf dry weight	0.999**	0.992**	0.973**	0.983**	1.000				
Plant N content	0.830**	0.879**	0.931**	0.915**	0.836**	1.000			
Pod number	0.993**	0.995**	0.979**	0.985**	0.995**	0.863**	1.000		
Seed number	0.993**	0.986**	0.962**	0.971**	0.994**	0.822**	0.996**	1.000	
Seed yield	0.996**	0.984**	0.957**	0.968**	0.996**	0.807**	0.993**	0.999**	1.000

Table 176. Linear regression values of plant characteristics in Experiment 4 at 80 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.978**	1.000							
Specific leaf weight	0.996**	0.984**	1.000						
Specific leaf area	0.998**	0.973**	0.998**	1.000					
Leaf dry weight	0.998**	0.989**	0.996**	0.994**	1.000				
Plant N content	0.986**	0.941**	0.982**	0.990**	0.976**	1.000			
Pod number	0.981**	0.937**	0.975**	0.984**	0.971**	0.986**	1.000		
Seed number	0.937**	0.862**	0.924**	0.942**	0.916**	0.962**	0.981**	1.000	
Seed yield	0.963**	0.897**	0.950**	0.965**	0.946**	0.974**	0.989**	0.989**	1.000

Table 177. Linear regression values of plant characteristics in Experiment 4 at 100 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.994**	1.000							
Specific leaf weight	0.995**	0.993**	1.000						
Specific leaf area	0.998**	0.993**	0.997**	1.000					
Leaf dry weight	0.999**	0.997**	0.996**	0.998**	1.000				
Plant N content	0.977**	0.991**	0.986**	0.981**	0.982**	1.000			
Pod number	0.992**	0.981**	0.976**	0.986**	0.990**	0.950**	1.000		
Seed number	0.958**	0.932**	0.930**	0.946**	0.951**	0.881**	0.981**	1.000	
Seed yield	0.977**	0.954**	0.957**	0.970**	0.971**	0.914**	0.989**	0.989**	1.000

Table 178. Linear regression values of plant characteristics in Experiment 4 at 120 DAS (\* significant at  $P < 0.05$ , \*\* significant at  $P < 0.01$ )

Parameter	Leaf area	Plant dry weight	Specific leaf weight	Specific leaf area	Leaf dry weight	Plant N content	Pod number	Seed number	Seed yield
Leaf area	1.000								
Plant dry weight	0.987**	1.000							
Specific leaf weight	0.986**	0.987**	1.000						
Specific leaf area	0.990**	0.986**	0.999**	1.000					
Leaf dry weight	0.999**	0.991**	0.988**	0.990**	1.000				
Plant N content	0.978**	0.940**	0.960**	0.967**	0.973**	1.000			
Pod number	0.993**	0.999**	0.989**	0.989**	0.995**	0.950**	1.000		
Seed number	0.990**	0.975**	0.961**	0.964**	0.989**	0.963**	0.981**	1.000	
Seed yield	0.993**	0.984**	0.979**	0.983**	0.992**	0.967**	0.989**	0.989**	1.000



Table 179. Effect of ethrel spray at 60d (post flowering stage) on nitrogen uptake efficiency ( $\text{kg ha}^{-1}$ ), nitrogen utilization efficiency ( $\text{kg ha}^{-1}$ ) and nitrogen use efficiency ( $\text{kg ha}^{-1}$ ) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	...	Nitrogen uptake efficiency				Nitrogen utilization efficiency			
		Ethrel concentration (μL/L)							
		0	100	200	Mean	0	100	200	Mean
N <sub>40</sub>	0.43	0.48	0.55	0.49	11.56	11.26	10.97	11.26	
N <sub>60</sub>	0.44	0.51	0.57	0.51	12.47	12.09	11.79	12.12	
N <sub>80</sub>	0.48	0.54	0.59	0.54	13.04	12.56	12.33	12.64	
Mean	0.45	0.51	0.57		12.36	11.97	11.70		
L.S.D. at 5%									
Nitrogen use efficiency									
N <sub>40</sub>	4.93	5.44	6.00	5.46	Nitrogen uptake efficiency				Nitrogen use efficiency
N <sub>60</sub>	5.53	6.13	6.72	6.12	Nitrogen uptake efficiency				Nitrogen use efficiency
N <sub>80</sub>	6.26	6.83	7.27	6.79	Spray (S)				0.046
Mean	5.57	6.13	6.66		Nitrogen (N)				0.046
					S x N				0.080

Table 180. Effect of ethrel spray at 60d (post flowering stage) on nitrogen uptake efficiency (kg ha<sup>-1</sup>), nitrogen utilization efficiency (kg ha<sup>-1</sup>) and nitrogen use efficiency (kg ha<sup>-1</sup>) of mustard (*Brassica juncea* L.) cultivar Alankar grown with four basal levels of nitrogen

Nitrogen (kg/ha)	Nitrogen uptake efficiency				Nitrogen utilization efficiency			
	Ethrel concentration (μL/L)							
	0	100	200	Mean	0	100	200	Mean
N <sub>40</sub>	0.29	0.34	0.40	0.34	12.41	11.84	11.25	11.84
N <sub>60</sub>	0.32	0.38	0.44	0.38	12.81	11.92	11.28	12.00
N <sub>80</sub>	0.36	0.41	0.46	0.41	11.93	11.92	11.48	11.78
Mean	0.32	0.38	0.43		12.38	11.89	11.34	
L.S.D. at 5%								
Nitrogen use efficiency								
N <sub>40</sub>	3.60	4.02	4.56	4.06	Nitrogen uptake efficiency	Nitrogen utilization efficiency	Nitrogen use efficiency	
N <sub>60</sub>	4.06	4.49	4.93	4.49				
N <sub>80</sub>	4.25	4.89	5.28	4.81	Spray (S)	0.004	0.374	0.127
Mean	3.97	4.47	4.92		Nitrogen (N)	0.004	NS	0.127
					S x N	0.007	NS	NS